

Affine and Vector Spaces

LA1♦1. Suppose ℓ_1 and ℓ_2 are skew lines in the space \mathbb{R}^3 . Is it true that lines PQ , where $P \in \ell_1, Q \in \ell_2$, sweep the whole space?

LA1♦2. Find a basis of the vector space $V = \{p(x) \in \mathbb{R}_4[x] : p'(5) = 0\}$.

LA1♦3. Find a dimension and a basis of the vector space

- (a) of all symmetric matrices $A \in \text{Mat}_n(\mathbb{R})$.
- (b) of all skew-symmetric matrices $A \in \text{Mat}_n(\mathbb{R})$.
- (c) $\mathfrak{sl}_n(\mathbb{R}) = \{A \in \text{Mat}_n(\mathbb{R}) : \text{tr } A = 0\}$.

LA1♦4. Give an example of a finite dimensional space V and three its pairwise transversal subspaces U, W, T (that is, intersecting only at the origin) such that $\dim U + \dim W + \dim T = \dim V$, but $U + W + T \neq V$.

LA1♦5. Suppose $\dim(U + V) = \dim(U \cap V) + 1$ for some two vector subspaces $U, V \subset \mathbb{R}^n$. Is it true that $U + V$ is equal to one of the subspaces U, V and $U \cap V$ is equal to another?

LA1♦6. Is it possible that the intersection of the positive orthant

$$\{(x_1, x_2, x_3, x_4) : x_1, x_2, x_3, x_4 \geq 0\} \subset \mathbb{R}^4$$

with some 2-dimensional plane is a square?

LA1♦7. Let V be a vector space of dimension n over the finite field \mathbb{F}_q of q elements. How many

- (a) vectors
 - (b) bases
 - (c) k -dimensional subspaces
- are there in V ?

LA1♦8. Let V be an affine space of dimension n over the finite field \mathbb{F}_q of q elements. How many k -dimensional affine subspaces are there in V ?

LA1♦9. For which $c \in \mathbb{R}$ the hyperplane $\sum x_j = c$ intersects with

- (a) the three dimensional cube $I^3 = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : |x_j| \leq 1 \ \forall j\}$?
- (b) the four dimensional cube $I^4 = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : |x_j| \leq 1 \ \forall j\}$?

Draw all the polygons and polyhedra respectively that are cut from the cube by such hyperplanes.

LA1♦10. Prove that the vector space of all continuous functions on \mathbb{R} is infinite dimensional.