

If you have left from the schedule please reach out. Your doctor is here to help.

Clinic Phone _____

Your medication taper

This plan slowly lowers your medication over time. Going slowly is what makes it work. Most people do best when each reduction is small and they stay at each step long enough to settle before the next one.

Some symptoms during a taper are normal and usually pass. If a step feels too hard, that is not a failure — it usually means staying at that step longer, not going back up. Do not change your dose on your own. Talk to your doctor first.

Never stop a benzodiazepine suddenly. Stopping abruptly can be dangerous.

Getting a pill organizer

A weekly pill organizer makes this much easier and much safer. You are taking your medicine three times a day, so look for:

- Search for a **"3-times-a-day pill organizer"**. These usually divide each day into morning, noon, and evening slots.

Filling your organizer each week

1. Find the row in the table below whose date range includes the coming week.
2. That row tells you exactly what to take, and at what times of day.
3. Fill the whole week at once, on the same day each week — Sunday works well for most people.
4. Check the dates before you fill. Some weeks the dose changes, and it is easy to fill from the wrong row.
5. If a tablet needs to be halved, use a pill cutter from any pharmacy. Ask your pharmacist to show you.
6. Keep the rest of the medicine in its original bottle, out of reach of children and pets.

Reminder apps

A phone reminder helps a lot, especially on days the dose changes. In your phone's app store, search for "medication reminder", "pill reminder", or "medisafe". Set one alarm for each time of day you take a dose. Your phone's built-in Clock or Reminders app works fine too.

Your schedule

Take the medicines listed for each date range. Do not skip ahead.

DATES	HOW LONG	MORNING	MIDDAY	NIGHT
8/11/26 – 8/17/26	1 week	1 × 2mg lorazepam	1 × 2mg lorazepam	1 × 2mg lorazepam
8/18/26 – 8/24/26	1 week	1 × 2mg lorazepam	1 × 2mg lorazepam	$\frac{1}{2}$ × 2mg lorazepam + 1 × 10mg diazepam
8/25/26 – 8/31/26	1 week	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	1 × 2mg lorazepam	1 × 1mg lorazepam + 1 × 10mg diazepam
9/1/26 – 9/7/26	1 week	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	1 × 2mg lorazepam	$\frac{1}{2}$ × 1mg lorazepam + $1\frac{1}{2}$ × 10mg diazepam
9/8/26 – 9/14/26	1 week	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	$\frac{1}{2}$ × 1mg lorazepam + $1\frac{1}{2}$ × 10mg diazepam
9/15/26 – 9/28/26	2 weeks	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	2 × 10mg diazepam
9/29/26 – 10/12/26	2 weeks	1 × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	$1\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	2 × 10mg diazepam
10/13/26 – 10/26/26	2 weeks	1 × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	1 × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	2 × 10mg diazepam
10/27/26 – 11/9/26	2 weeks	$\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	1 × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	2 × 10mg diazepam
11/10/26 – 11/23/26	2 weeks	$\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	$\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	2 × 10mg diazepam
11/24/26 – 12/7/26	2 weeks	$\frac{1}{2}$ × 10mg diazepam	$\frac{1}{2}$ × 1mg lorazepam + $\frac{1}{2}$ × 10mg diazepam	2 × 10mg diazepam

12/8/26 – 12/21/26	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	$\frac{1}{2} \times 10\text{mg}$ diazepam	2 $\times$ 10mg diazepam
12/22/26 – 1/4/27	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	$\frac{1}{2} \times 10\text{mg}$ diazepam	1 $\times$ 10mg diazepam + 4 $\times$ 2mg diazepam
1/5/27 – 1/18/27	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	$\frac{1}{2} \times 10\text{mg}$ diazepam	1 $\times$ 10mg diazepam + 3 $\times$ 2mg diazepam
1/19/27 – 2/1/27	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	$\frac{1}{2} \times 10\text{mg}$ diazepam	1 $\times$ 10mg diazepam + 2 $\times$ 2mg diazepam
2/2/27 – 2/15/27	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	$\frac{1}{2} \times 10\text{mg}$ diazepam	1 $\times$ 10mg diazepam + 1 $\times$ 2mg diazepam
2/16/27 – 3/1/27	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	$\frac{1}{2} \times 10\text{mg}$ diazepam	1 $\times$ 10mg diazepam
3/2/27 – 3/15/27	2 weeks	$\frac{1}{2} \times 10\text{mg}$ diazepam	2 $\times$ 2mg diazepam	1 $\times$ 10mg diazepam
3/16/27 – 3/29/27	2 weeks	2 $\times$ 2mg diazepam	2 $\times$ 2mg diazepam	1 $\times$ 10mg diazepam
3/30/27 – 4/12/27	2 weeks	2 $\times$ 2mg diazepam	1 $\frac{1}{2}$ $\times$ 2mg diazepam	1 $\times$ 10mg diazepam
4/13/27 – 4/26/27	2 weeks	1 $\frac{1}{2}$ $\times$ 2mg diazepam	1 $\frac{1}{2}$ $\times$ 2mg diazepam	1 $\times$ 10mg diazepam
4/27/27 – 5/10/27	2 weeks	1 $\frac{1}{2}$ $\times$ 2mg diazepam	1 $\times$ 2mg diazepam	1 $\times$ 10mg diazepam
5/11/27 – 5/24/27	2 weeks	1 $\times$ 2mg diazepam	1 $\times$ 2mg diazepam	1 $\times$ 10mg diazepam
5/25/27 – 6/7/27	2 weeks	1 $\times$ 2mg diazepam	$\frac{1}{2} \times 2\text{mg}$ diazepam	1 $\times$ 10mg diazepam
6/8/27 – 6/21/27	2 weeks	$\frac{1}{2} \times 2\text{mg}$ diazepam	$\frac{1}{2} \times 2\text{mg}$ diazepam	1 $\times$ 10mg diazepam
6/22/27 – 7/5/27	2 weeks	$\frac{1}{2} \times 2\text{mg}$ diazepam	—	1 $\times$ 10mg diazepam

7/6/27 – 7/19/27	2 weeks	—	—	1 × 10mg diazepam
7/20/27 – 8/2/27	2 weeks	—	—	½ × 10mg diazepam + 2 × 2mg diazepam
8/3/27 – 8/16/27	2 weeks	—	—	4 × 2mg diazepam
8/17/27 – 8/30/27	2 weeks	—	—	½ × 10mg diazepam + 1 × 2mg diazepam
8/31/27 – 9/13/27	2 weeks	—	—	3 × 2mg diazepam
9/14/27 – 9/27/27	2 weeks	—	—	½ × 10mg diazepam
9/28/27 – 10/11/27	2 weeks	—	—	2 × 2mg diazepam
10/12/27 – 10/25/27	2 weeks	—	—	1½ × 2mg diazepam
10/26/27 – 11/8/27	2 weeks	—	—	1 × 2mg diazepam
11/9/27 – 11/22/27	2 weeks	—	—	½ × 2mg diazepam
11/23/27	—	Finished. No further doses.		

½ means half a tablet. Use a pill cutter.

Dose sequences are cited from Ashton, C.H., *Benzodiazepines: How They Work and How to Withdraw* (2002). **This schedule was generated by this software and was not produced, reviewed, or endorsed by Professor Ashton, her estate, or the Benzodiazepine Information Coalition.** Step durations are set by the prescriber and may differ from those published. Not reviewed by a clinician. For informational and research purposes only — not medical advice, not a prescription, and not a substitute for professional judgement. Verify every dose independently before use. Provided AS IS, without warranty of any kind; the author accepts no liability arising from its use. Do not start, stop, or change any medication except as directed by your prescriber.

benzo-taper v1.0 (engine v2) · Code 6D2G-2W3S-LVJ8-ZP (contains PHI) · Generated Tue, Aug 11, 2026

Dispensing breakdown

Quantities are grouped into **2-week fills**. A fill usually spans more than one step, so quantities are totalled across the whole fill rather than per step. No fill requires more than two tablet strengths of any one drug.

Prescription 1 — written 8/11/26

6 fills (initial fill plus 5 refills, 70 days of refills). A new prescription is required after this one: 5-refill limit reached.

Fill 1 — 8/11/26 to 8/24/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	2 mg	39 tablets
diazepam	10 mg	7 tablets

Patient instructions during this fill:

- **8/11/26 – 8/17/26:** Take one tablet of 2mg lorazepam in the morning; one tablet of 2mg lorazepam at midday; one tablet of 2mg lorazepam at night. [Schedule 3, starting dosage — Ashton 2002]
- **8/18/26 – 8/24/26:** Take one tablet of 2mg lorazepam in the morning; one tablet of 2mg lorazepam at midday; half a tablet of 2mg lorazepam and one tablet of 10mg diazepam at night. [Schedule 3, Stage 1 — Ashton 2002]

Fill 2 — 8/25/26 to 9/7/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	2 mg	14 tablets
lorazepam	1 mg	32 tablets
diazepam	10 mg	25 tablets

Patient instructions during this fill:

- **8/25/26 – 8/31/26:** Take one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one tablet of 2mg lorazepam at midday; one tablet of 1mg lorazepam and one tablet of 10mg diazepam at night. [Schedule 3, Stage 2 — Ashton 2002]
- **9/1/26 – 9/7/26:** Take one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one tablet of 2mg lorazepam at midday; half a tablet of 1mg lorazepam and one and a half tablets of 10mg diazepam at night. [Schedule 3, Stage 3 — Ashton 2002]

Fill 3 — 9/8/26 to 9/21/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	1 mg	46 tablets
diazepam	10 mg	39 tablets

Patient instructions during this fill:

- **9/8/26 – 9/14/26:** Take one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam at midday; half a tablet of 1mg lorazepam and one and a half tablets of 10mg diazepam at night. [Schedule 3, Stage 4 — Ashton 2002]
- **9/15/26 – 9/28/26:** Take one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 5 — Ashton 2002]

Fill 4 — 9/22/26 to 10/5/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	1 mg	39 tablets
diazepam	10 mg	42 tablets

Patient instructions during this fill:

- **9/15/26 – 9/28/26:** Take one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 5 — Ashton 2002]
- **9/29/26 – 10/12/26:** Take one tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 6 — Ashton 2002]

Fill 5 — 10/6/26 to 10/19/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	1 mg	32 tablets
diazepam	10 mg	42 tablets

Patient instructions during this fill:

- **9/29/26 – 10/12/26:** Take one tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one and a half tablets of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 6 — Ashton 2002]
- **10/13/26 – 10/26/26:** Take one tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 7 — Ashton 2002]

Fill 6 — 10/20/26 to 11/2/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	1 mg	25 tablets
diazepam	10 mg	42 tablets

Patient instructions during this fill:

- **10/13/26 – 10/26/26:** Take one tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 7 — Ashton 2002]
- **10/27/26 – 11/9/26:** Take half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 8 — Ashton 2002]

Prescription 2 — written 11/3/26

6 fills (initial fill plus 5 refills, 70 days of refills). A new prescription is required after this one: 5-refill limit reached.

Fill 7 — 11/3/26 to 11/16/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	1 mg	18 tablets
diazepam	10 mg	42 tablets

Patient instructions during this fill:

- **10/27/26 – 11/9/26:** Take half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; one tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 8 — Ashton 2002]
- **11/10/26 – 11/23/26:** Take half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 9 — Ashton 2002]

Fill 8 — 11/17/26 to 11/30/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
lorazepam	1 mg	11 tablets
diazepam	10 mg	42 tablets

Patient instructions during this fill:

- **11/10/26 – 11/23/26:** Take half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam in the morning; half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 9 — Ashton 2002]
- **11/24/26 – 12/7/26:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 10 — Ashton 2002]

Fill 9 — 12/1/26 to 12/14/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	42 tablets
lorazepam	1 mg	4 tablets

Patient instructions during this fill:

- **11/24/26 – 12/7/26:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 1mg lorazepam and half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 10 — Ashton 2002]
- **12/8/26 – 12/21/26:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 11 — Ashton 2002]

Fill 10 — 12/15/26 to 12/28/26 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	35 tablets
diazepam	2 mg	28 tablets

Patient instructions during this fill:

- **12/8/26 – 12/21/26:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; two tablets of 10mg diazepam at night. [Schedule 3, Stage 11 — Ashton 2002]
- **12/22/26 – 1/4/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and four tablets of 2mg diazepam at night. [Schedule 3, Stage 12 — Ashton 2002]

Fill 11 — 12/29/26 to 1/11/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	28 tablets
diazepam	2 mg	49 tablets

Patient instructions during this fill:

- **12/22/26 – 1/4/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and four tablets of 2mg diazepam at night. [Schedule 3, Stage 12 — Ashton 2002]
- **1/5/27 – 1/18/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and three tablets of 2mg diazepam at night. [Schedule 3, Stage 13 — Ashton 2002]

Fill 12 — 1/12/27 to 1/25/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	28 tablets
diazepam	2 mg	35 tablets

Patient instructions during this fill:

- **1/5/27 – 1/18/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and three tablets of 2mg diazepam at night. [Schedule 3, Stage 13 — Ashton 2002]
- **1/19/27 – 2/1/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and two tablets of 2mg diazepam at night. [Schedule 3, Stage 14 — Ashton 2002]

Prescription 3 — written 1/26/27

6 fills (initial fill plus 5 refills, 70 days of refills). A new prescription is required after this one: 5-refill limit reached.

Fill 13 — 1/26/27 to 2/8/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	28 tablets
diazepam	2 mg	21 tablets

Patient instructions during this fill:

- **1/19/27 – 2/1/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and two tablets of 2mg diazepam at night. [Schedule 3, Stage 14 — Ashton 2002]
- **2/2/27 – 2/15/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and one tablet of 2mg diazepam at night. [Schedule 3, Stage 15 — Ashton 2002]

Fill 14 — 2/9/27 to 2/22/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	28 tablets
diazepam	2 mg	7 tablets

Patient instructions during this fill:

- **2/2/27 – 2/15/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam and one tablet of 2mg diazepam at night. [Schedule 3, Stage 15 — Ashton 2002]
- **2/16/27 – 3/1/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 16 — Ashton 2002]

Fill 15 — 2/23/27 to 3/8/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	25 tablets
diazepam	2 mg	14 tablets

Patient instructions during this fill:

- **2/16/27 – 3/1/27:** Take half a tablet of 10mg diazepam in the morning; half a tablet of 10mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 16 — Ashton 2002]
- **3/2/27 – 3/15/27:** Take half a tablet of 10mg diazepam in the morning; two tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 17 — Ashton 2002]

Fill 16 — 3/9/27 to 3/22/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	18 tablets
diazepam	2 mg	42 tablets

Patient instructions during this fill:

- **3/2/27 – 3/15/27:** Take half a tablet of 10mg diazepam in the morning; two tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 17 — Ashton 2002]
- **3/16/27 – 3/29/27:** Take two tablets of 2mg diazepam in the morning; two tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 18 — Ashton 2002]

Fill 17 — 3/23/27 to 4/5/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	53 tablets

Patient instructions during this fill:

- **3/16/27 – 3/29/27:** Take two tablets of 2mg diazepam in the morning; two tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 18 — Ashton 2002]
- **3/30/27 – 4/12/27:** Take two tablets of 2mg diazepam in the morning; one and a half tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 19 — Ashton 2002]

Fill 18 — 4/6/27 to 4/19/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	46 tablets

Patient instructions during this fill:

- **3/30/27 – 4/12/27:** Take two tablets of 2mg diazepam in the morning; one and a half tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 19 — Ashton 2002]
- **4/13/27 – 4/26/27:** Take one and a half tablets of 2mg diazepam in the morning; one and a half tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 20 — Ashton 2002]

Prescription 4 — written 4/20/27

6 fills (initial fill plus 5 refills, 70 days of refills). A new prescription is required after this one: 5-refill limit reached.

Fill 19 — 4/20/27 to 5/3/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	39 tablets

Patient instructions during this fill:

- **4/13/27 – 4/26/27:** Take one and a half tablets of 2mg diazepam in the morning; one and a half tablets of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 20 — Ashton 2002]
- **4/27/27 – 5/10/27:** Take one and a half tablets of 2mg diazepam in the morning; one tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 21 — Ashton 2002]

Fill 20 — 5/4/27 to 5/17/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	32 tablets

Patient instructions during this fill:

- **4/27/27 – 5/10/27:** Take one and a half tablets of 2mg diazepam in the morning; one tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 21 — Ashton 2002]
- **5/11/27 – 5/24/27:** Take one tablet of 2mg diazepam in the morning; one tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 22 — Ashton 2002]

Fill 21 — 5/18/27 to 5/31/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	25 tablets

Patient instructions during this fill:

- **5/11/27 – 5/24/27:** Take one tablet of 2mg diazepam in the morning; one tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 22 — Ashton 2002]
- **5/25/27 – 6/7/27:** Take one tablet of 2mg diazepam in the morning; half a tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 23 — Ashton 2002]

Fill 22 — 6/1/27 to 6/14/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	18 tablets

Patient instructions during this fill:

- **5/25/27 – 6/7/27:** Take one tablet of 2mg diazepam in the morning; half a tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 23 — Ashton 2002]
- **6/8/27 – 6/21/27:** Take half a tablet of 2mg diazepam in the morning; half a tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 24 — Ashton 2002]

Fill 23 — 6/15/27 to 6/28/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	11 tablets

Patient instructions during this fill:

- **6/8/27 – 6/21/27:** Take half a tablet of 2mg diazepam in the morning; half a tablet of 2mg diazepam at midday; one tablet of 10mg diazepam at night. [Schedule 3, Stage 24 — Ashton 2002]
- **6/22/27 – 7/5/27:** Take half a tablet of 2mg diazepam in the morning; one tablet of 10mg diazepam at night. [Schedule 3, Stage 25 — Ashton 2002]

Fill 24 — 6/29/27 to 7/12/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	14 tablets
diazepam	2 mg	4 tablets

Patient instructions during this fill:

- **6/22/27 – 7/5/27:** Take half a tablet of 2mg diazepam in the morning; one tablet of 10mg diazepam at night. [Schedule 3, Stage 25 — Ashton 2002]
- **7/6/27 – 7/19/27:** Take one tablet of 10mg diazepam at night. [Schedule 3, Stage 26 — Ashton 2002]

Prescription 5 — written 7/13/27

6 fills (initial fill plus 5 refills, 70 days of refills). A new prescription is required after this one: 5-refill limit reached.

Fill 25 — 7/13/27 to 7/26/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	11 tablets
diazepam	2 mg	14 tablets

Patient instructions during this fill:

- **7/6/27 – 7/19/27:** Take one tablet of 10mg diazepam at night. [Schedule 3, Stage 26 — Ashton 2002]
- **7/20/27 – 8/2/27:** Take half a tablet of 10mg diazepam and two tablets of 2mg diazepam at night. [Schedule 3, Stage 27 — Ashton 2002]

Fill 26 — 7/27/27 to 8/9/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	4 tablets
diazepam	2 mg	42 tablets

Patient instructions during this fill:

- **7/20/27 – 8/2/27:** Take half a tablet of 10mg diazepam and two tablets of 2mg diazepam at night. [Schedule 3, Stage 27 — Ashton 2002]
- **8/3/27 – 8/16/27:** Take four tablets of 2mg diazepam at night. [Schedule 3, Stage 28 — Ashton 2002]

Fill 27 — 8/10/27 to 8/23/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	4 tablets
diazepam	2 mg	35 tablets

Patient instructions during this fill:

- **8/3/27 – 8/16/27:** Take four tablets of 2mg diazepam at night. [Schedule 3, Stage 28 — Ashton 2002]
- **8/17/27 – 8/30/27:** Take half a tablet of 10mg diazepam and one tablet of 2mg diazepam at night. [Schedule 3, Stage 29 — Ashton 2002]

Fill 28 — 8/24/27 to 9/6/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	4 tablets
diazepam	2 mg	28 tablets

Patient instructions during this fill:

- **8/17/27 – 8/30/27:** Take half a tablet of 10mg diazepam and one tablet of 2mg diazepam at night. [Schedule 3, Stage 29 — Ashton 2002]
- **8/31/27 – 9/13/27:** Take three tablets of 2mg diazepam at night. [Schedule 3, Stage 30 — Ashton 2002]

Fill 29 — 9/7/27 to 9/20/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	4 tablets
diazepam	2 mg	21 tablets

Patient instructions during this fill:

- **8/31/27 – 9/13/27:** Take three tablets of 2mg diazepam at night. [Schedule 3, Stage 30 — Ashton 2002]
- **9/14/27 – 9/27/27:** Take half a tablet of 10mg diazepam at night. [Schedule 3, Stage 31 — Ashton 2002]

Fill 30 — 9/21/27 to 10/4/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	10 mg	4 tablets
diazepam	2 mg	14 tablets

Patient instructions during this fill:

- **9/14/27 – 9/27/27:** Take half a tablet of 10mg diazepam at night. [Schedule 3, Stage 31 — Ashton 2002]
- **9/28/27 – 10/11/27:** Take two tablets of 2mg diazepam at night. [Schedule 3, Stage 32 — Ashton 2002]

Prescription 6 — written 10/5/27

4 fills (initial fill plus 3 refills, 35 days of refills). This prescription covers the remainder of the schedule.

Fill 31 — 10/5/27 to 10/18/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	2 mg	25 tablets

Patient instructions during this fill:

- **9/28/27 – 10/11/27:** Take two tablets of 2mg diazepam at night. [Schedule 3, Stage 32 — Ashton 2002]
- **10/12/27 – 10/25/27:** Take one and a half tablets of 2mg diazepam at night. [Schedule 3, Stage 33 — Ashton 2002]

Fill 32 — 10/19/27 to 11/1/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	2 mg	18 tablets

Patient instructions during this fill:

- **10/12/27 – 10/25/27:** Take one and a half tablets of 2mg diazepam at night. [Schedule 3, Stage 33 — Ashton 2002]
- **10/26/27 – 11/8/27:** Take one tablet of 2mg diazepam at night. [Schedule 3, Stage 34 — Ashton 2002]

Fill 33 — 11/2/27 to 11/15/27 (14 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	2 mg	11 tablets

Patient instructions during this fill:

- **10/26/27 – 11/8/27:** Take one tablet of 2mg diazepam at night. [Schedule 3, Stage 34 — Ashton 2002]
- **11/9/27 – 11/22/27:** Take half a tablet of 2mg diazepam at night. [Schedule 3, Stage 35 — Ashton 2002]

Fill 34 — 11/16/27 to 11/22/27 (7 days)

MEDICATION	STRENGTH	QUANTITY PER FILL
diazepam	2 mg	4 tablets

Patient instructions during this fill:

- **11/9/27 – 11/22/27:** Take half a tablet of 2mg diazepam at night. [Schedule 3, Stage 35 — Ashton 2002]

Prescription grouping applies California Schedule IV limits: at most five refills, refills capped at 120 days combined, and expiry six months after the date written. Whichever limit binds first ends the prescription. Verify against current state rules.

Dose sequences are cited from Ashton, C.H., *Benzodiazepines: How They Work and How to Withdraw* (2002). **This schedule was generated by this software and was not produced, reviewed, or endorsed by Professor Ashton, her estate, or the Benzodiazepine Information Coalition.** Step durations are set by the prescriber and may differ from those published. Not reviewed by a clinician. For informational and research purposes only — not medical advice, not a prescription, and not a substitute for professional judgement. Verify every dose independently before use. Provided AS IS, without warranty of any kind; the author accepts no liability arising from its use. Do not start, stop, or change any medication except as directed by your prescriber.

benzo-taper v1.0 (engine v2) · Code 6D2G-2W3S-LVJ8-ZP (contains PHI) · Generated Tue, Aug 11, 2026

NOT A
PRESCRIPTION
PRESCRIBER
REVIEW REQUIRED

Schedule code

This code regenerates this exact schedule. Enter it in the "Enter schedule code here" box on the first screen to return to it, adjust it, or reprint it.

6D2G-2W3S-LVJ8-ZP

CONTAINS PHI This code encodes the patient's dosing and taper dates. Handle it with the same care as any other identifiable clinical information. Do not post it publicly or send it over unsecured channels.

- Locked to engine version 2. A future version will refuse it rather than regenerate something different.
- Checksummed — a mistyped character is rejected outright, never silently accepted as a different schedule.
- The characters 0, O, 1 and I are never used, so they cannot be confused.

Schedule summary

MEDICATIONS	Lorazepam
STARTING DOSE	60mg diazepam-equivalent
START DATE	Tue, Aug 11, 2026
TOTAL DURATION	67 weeks (1.3 years)
DOSE SEQUENCES	36 steps reproduce dose sequences from published schedules; 0 generated by this tool
DURATIONS	Set by the prescriber
CITED SOURCE ENTERS AT	Schedule 3, stage starting dosage (Ashton 2002)

Dose sequences are cited from Ashton, C.H., *Benzodiazepines: How They Work and How to Withdraw* (2002). **This schedule was generated by this software and was not produced, reviewed, or endorsed by Professor Ashton, her estate, or the Benzodiazepine Information Coalition.** Step durations are set by the prescriber and may differ from those published. Not reviewed by a clinician. For informational and research purposes only — not medical advice, not a prescription, and not a substitute for professional judgement. Verify every dose independently before use. Provided AS IS, without warranty of any kind; the author accepts no liability arising from its use. Do not start, stop, or change any medication except as directed by your prescriber.