

Rough numbers at a fixed distance with prescribed Liouville signs

Jizhou Guo

mitsuha2021b@gmail.com

October 2026

Abstract

Let λ be the Liouville function and let $h \geq 2$ be a fixed even integer. We show that there is an absolute constant $\gamma > 0$ such that, uniformly for $2 \leq z \leq \exp((\log X)^{1/3750})$, each of the four sign patterns of $(\lambda(n), \lambda(n+h))$ occurs for $\frac{1}{4}XV_h(z)(1 + O_h((\log X)^{-\gamma}))$ integers $n \leq X$ such that n and $n+h$ have no prime factor up to z ; here $XV_h(z)$ is the main term for the number of such n . Consequently each pattern occurs for $\gg_h X/(\log X)^{1/1875}$ integers $n \leq X$ with n and $n+h$ squarefree, free of prime factors up to $\exp((\log X)^{1/3750})$, and with at most $(1 - 1/4000)\log \log X$ prime factors. We prove corresponding statements for the weights $r^{\omega_z(n)}$ on the small prime factors and $r^{\Omega(n)}$ on all prime factors, in each case with a squarefreeness restriction at the small primes. The proofs adapt the graph argument of OpenAI's proof of the two-point Chowla conjecture, keeping the sieve density inside the operator.

2020 Mathematics Subject Classification. Primary 11N37; Secondary 11N36, 11N25, 11P32.

1 Introduction

All logarithms are natural. The function $\Omega(n)$ counts prime factors with multiplicity, $\omega(n)$ counts distinct prime factors, and $\lambda(n) = (-1)^{\Omega(n)}$. Put $P^-(1) = \infty$ and write $P^-(n)$ for the least prime factor otherwise. Throughout the paper $h \geq 2$ is a fixed even integer. Define

$$R_z(n) = \mathbf{1}_{P^-(n) > z}, \quad V_h(z) = \prod_{p \leq z} \left(1 - \frac{\nu_h(p)}{p}\right), \quad \nu_h(p) = \begin{cases} 1, & p \mid h, \\ 2, & p \nmid h. \end{cases} \quad (1.1)$$

Sums and counts over $n \leq X$ always concern positive integers.

Theorem 1.1. *There is an absolute constant $\gamma > 0$ such that, for each fixed even $h \geq 2$ and every sufficiently large X , uniformly for $2 \leq z \leq \exp((\log X)^{1/3750})$ and $(s, t) \in \{\pm 1\}^2$,*

$$\#\{n \leq X : R_z(n)R_z(n+h) = 1, (\lambda(n), \lambda(n+h)) = (s, t)\} = \frac{1}{4}XV_h(z)(1 + O_h((\log X)^{-\gamma})). \quad (1.2)$$

Theorem 1.2. *For each fixed even $h \geq 2$ and each sign pattern there are $\gg_h X/(\log X)^{1/1875}$ integers $n \leq X$ such that n and $n+h$ are squarefree, have no prime factor up to $\exp((\log X)^{1/3750})$, have that sign pattern, and satisfy*

$$\Omega(n), \Omega(n+h) \leq (1 - 1/4000)\log \log X. \quad (1.3)$$

This holds for every sufficiently large X .

For $z \geq 2$ let

$$S_z(n) = \prod_{p \leq z} \mathbf{1}_{p^2 \nmid n}, \quad \omega_z(n) = \#\{p \leq z : p \mid n\}, \quad b_z(n) = S_z(n)r^{\omega_z(n)}. \quad (1.4)$$

Theorem 1.3. *There is an absolute constant $\gamma > 0$, which may be the same as in Theorem 1.1, such that for each fixed even $h \geq 2$, each fixed $0 < r < 1$, and every sufficiently large X , uniformly for $2 \leq z \leq \exp((\log X)^{1/3750})$ and $(s, t) \in \{\pm 1\}^2$,*

$$\sum_{\substack{n \leq X \\ (\lambda(n), \lambda(n+h)) = (s, t)}} b_z(n)b_z(n+h) = \frac{1}{4}XV_{r,h}^{\text{sf}}(z)(1 + O_{h,r}((\log X)^{-\gamma})), \quad (1.5)$$

where $V_{r,h}^{\text{sf}}(z) = \prod_{p \leq z} v_{r,h}(p)$ and

$$v_{r,h}(p) = \begin{cases} 1 - 2/p + 2r(p-1)/p^2, & p \nmid h, \\ 1 - 1/p + r^2(p-1)/p^2, & p^2 \mid h, \\ 1 - 1/p + r^2(p-2)/p^2, & p \mid h, p^2 \nmid h. \end{cases} \quad (1.6)$$

The threshold and implied constant may depend on h, r ; r is fixed before X .

The constants for the following theorem are separate from those used in Theorems 1.1 and 1.3. Let C_3 be the absolute graph constant in [20, (8.7), p. 41], with the same permissible choice for the two-layer operators of Proposition 9.1. Put $C_\Omega = \max(4, C_3)$ and fix

$$W_\Omega = 16eC_\Omega^2, \quad \alpha_\Omega = \frac{1}{1200W_\Omega}, \quad \delta_\Omega = \sqrt{\alpha_\Omega/216}, \quad r = 1 - \delta_\Omega, \quad \gamma_\Omega = \alpha_\Omega/3600. \quad (1.7)$$

Theorem 1.4. *With the parameters (1.7) and $B(n) = r^{\Omega(n)}S_y(n)$, for each fixed even $h \geq 2$, every sufficiently large X , $2 \leq y \leq \exp((\log X)^{1/216})$, and every nonzero $(e_1, e_2) \in \{0, 1\}^2$,*

$$\left| \sum_{n \leq X} \lambda(n)^{e_1} \lambda(n+h)^{e_2} B(n)B(n+h) \right| \ll_h X(\log X)^{-2\delta_\Omega - \gamma_\Omega}. \quad (1.8)$$

In the same range the unsigned mass satisfies

$$\sum_{n \leq X} B(n)B(n+h) \asymp_h X(\log X)^{-2\delta_\Omega}. \quad (1.9)$$

The parameter y imposes squarefreeness at small primes; it imposes no roughness restriction.

Remark 1.5. Put $\rho = 1/3750$. For every fixed $0 < \kappa \leq \rho$ and $0 < \epsilon < \kappa$, for every sufficiently large X , every sign pattern occurs for at least $c_{h,\kappa,\epsilon}X/(\log X)^{2\kappa}$ integers $n \leq X$ with both endpoints squarefree and

$$\Omega(n), \Omega(n+h) \leq (1 - \kappa + \epsilon) \log \log X. \quad (1.10)$$

For $\kappa = \rho$ both endpoints may additionally be required to have no prime factor up to $\exp((\log X)^\rho)$. For $\kappa < \rho$ this additional condition is not part of the assertion. All parameters are fixed before X .

Mertens' theorem gives $V_h(z) \asymp_h (\log z)^{-2}$; see Lemma 2.1. At the upper endpoint of Theorem 1.1, the density is of order $(\log X)^{-2/3750}$. An absolute cancellation estimate must therefore save more than this density to give a relative asymptotic. The bound on Ω in Theorem 1.2 grows with X .

OpenAI's paper of 24 September 2026 [20, Theorem 1.1 and Proposition 2.1, pp. 3, 7] gives an absolute power-of-logarithm saving for Liouville correlations in fixed affine forms and fixed progressions, including nonunit classes. For a fixed z , finite inclusion–exclusion over the primes up to z , together with classical one-point cancellation, gives each rough sign pattern a quarter of the rough-pair mass. Its Theorem 1.2 (p. 4) gives qualitative ordinary cancellation for bounded multiplicative functions when an original factor is uniformly nonpretentious, also with fixed progression restrictions.

Goldston, Graham, Pintz and Yıldırım (18 March 2008) [7, Theorems 6 and 9, pp. 5, 7] prove that $\Omega(n) = \Omega(n+2) = B$ infinitely often for each integer $B \geq 5$, and for each fixed even h that $\Omega(n) = \Omega(n+h) = A$ infinitely often when $A \geq 4$ if $4 \mid h$, or $A \geq 5$ if $h \equiv 2 \pmod{4}$. Helfgott and Radziwiłł (11 March 2021; version 2, 13 April 2021) [11, Corollaries 1.4, 1.7–1.8, pp. 5–7] give correlations with restrictions on Ω in logarithmic averages and at almost all scales. Matomäki, Radziwiłł and Tao (4 September 2015; version 2, 23 September 2015) [17, Theorems 1.6 and 1.9, pp. 2–3] prove positive lower natural density for all eight Liouville patterns of length three and all nine adjacent Möbius patterns.

Weighted two-point sums of the present kind were formulated as a conjecture by Preobrazhenskii and Preobrazhenskaya [24, Conjecture 2, p. 3]. For $y \geq 2$ let $\omega^-(n, y)$ and $\omega^+(n, y)$ be the numbers of prime divisors of n that are less than y and at least y , and on squarefree n put $\tau^\pm(n) = \tau_{\kappa_1, \kappa_2}^\pm(n, y) = \kappa_1^{\omega^-(n, y)} \kappa_2^{\omega^+(n, y)}$. Their conjecture states that, for fixed $0 \leq \kappa_1 < \kappa_2$ and $y = \exp((\log x)^\delta)$ with some fixed $0 < \delta < 1$,

$$\sum_{N/2 \leq n \leq N} \mu(n+h_1)\mu(n+h_2)\tau^\pm(n+h_1)\tau^\pm(n+h_2) = o\left(\sum_{N/2 \leq n \leq N} \tau^\pm(n+h_1)\tau^\pm(n+h_2)\right).$$

They remark that the case $\kappa_1 = \kappa_2 = 1$ is the two-point Chowla conjecture for μ , and state that for $0 \leq \kappa_1 < \kappa_2$ the conjecture essentially implies the twin prime conjecture; their Theorem 4 includes a logarithmically averaged two-point estimate for λ with a sieve weight. The weights R_z and b_z of Theorems 1.1 and 1.3 have this shape with $\kappa_2 = 1$ and $\kappa_1 = 0$, respectively $\kappa_1 = r$. The two theorems are stated for λ , impose squarefreeness only at the primes up to z , and hold for $z \leq \exp((\log X)^{1/3750})$.

For the truncated functions $\lambda_y(n) = (-1)^{\Omega_y(n)}$ and $\mu_y(n) = \mu^2(n)(-1)^{\omega_y(n)}$, where Ω_y and ω_y count only the prime factors up to y , Daboussi and Sárközy [5, Corollary 2] prove $|\sum_{n \leq x} \lambda_y(n)\lambda_y(n+1)| \ll x((\log y)^{-9} + \exp(-u/8))$ for $3 \leq y \leq x$ and $u = \log x / \log y$, and Mangerel [15, Theorem 1.2(a)] proves, for each fixed a and in a range of $\beta = \log x / \log y$ specified there, $\sum_{n \leq x} \mu_y(n)\mu_y(n+a) \ll x(1/\log \log y + \exp(-\frac{1}{21}\beta \log \beta))$. Mangerel's introduction attributes the first estimate of this type, with $y \leq (\log x)^{2+o(1)}$, to Cassaigne, Ferenczi, Mauduit, Rivat and Sárközy [3]. These results concern the prime factors up to y ; Theorem 1.1 concerns integers without such prime factors.

Without sieve conditions, Tao [26, Theorem 1.2] proved the logarithmically averaged two-point Chowla conjecture, Tao and Teräväinen [27, Corollary 1.14] proved that $\sum_{n \leq X} \lambda(n+h_1)\lambda(n+h_2) = o(X)$ for X outside a set of logarithmic density zero, and Pilatte [22, Theorem 1.1] proved $\sum_{n \leq x} \lambda(n)\lambda(n+1)/n \ll (\log x)^{1-c}$ with an absolute $c > 0$. Tao and Teräväinen [28, Theorem 3.1] give a quantitative two-point estimate for 1-bounded multiplicative functions outside an exceptional set of scales, and deduce an asymptotic formula for consecutive smooth numbers at those scales [28,

Theorem 1.8]; a second paper of the release containing [O] proves the joint Dickman law for the largest prime factors of n and $n + 1$ in natural density [21, Theorem 1.1]. Charamaras and Richter [4, Theorem A] prove asymptotic independence of $\Omega(n)$ and $\Omega(n + 1)$ along logarithmic averages, and Hughes [14, Corollary 7.1] gives logarithmic frequency $1/4$ for each sign pair of λ at two affine forms in every window $(x/\omega, x]$ with ω above an explicit threshold.

A Lean 4 repository [13] contains formal statements at specific small sieve levels: for every z with $\prod_{p \leq z} p \leq 548$ (that is, $z \leq 10$) and every admissible residue class modulo that product, infinitely many n with $n(n + 2)$ free of prime factors up to z and $\Omega(n(n + 2))$ odd (5 September 2026); and, when the product is at most 2310, a lower bound for the logarithmic mass of such n in each admissible class (14 September 2026), described there as an instance of [26]. The sifted sum $\sum_{n \leq N, P^-(n) > D, P^-(n+2) > D} \lambda(n)\lambda(n + 2)$ with $D = N^{1/8}$ is considered in [18, Section 9], in a reduction of the twin prime conjecture to a hypothesis on $\lambda(n)\lambda(n + 2)$ in arithmetic progressions.

The author’s papers [9, 10], both dated 9 October 2026, treat, respectively, sums of two rough numbers with prescribed Liouville signs and shifted primes with a prescribed parity of the number of prime factors. The paper [8], dated 8 October 2026, treats Goldbach-type representations with prescribed Liouville signs. The present argument concerns a fixed distance between the two integers.

What is used. Write [O] for [20], taken at repository commit

adc7f1241b42e322a6451854ab7e4b4c146bf78a.

We use [O, Lemma 5.5, p. 26], the arithmetic and combinatorial statements [O, Lemmas 6.1, 6.3–6.4, 7.1–7.2, pp. 27, 30–37], and the finite-dimensional transfer [O, Lemma 8.1, p. 39] as stated, checking their hypotheses in Proposition 5.2. For the padding weight $16^{\omega(u)}$, vertex weight $17^{\omega_{\mathcal{Q}}}$, rough-target cutoff, balanced soft weight and numerical signs in the full- Ω argument, the original statements of [O, Lemma 2.2, Lemmas 4.1–4.2, 4.4–4.6, 5.2–5.3, 5.6, 6.2, 6.5, 7.3, Theorem 5.4, Lemma 8.2 and Propositions 8.3–8.4] do not serve as direct instances. We give the required versions in Sections 2–6, 7 and 9, following those proofs with their page references. The comparison of [O, Lemma 3.1 and Corollaries 3.2–3.3, pp. 10–12] is replaced by Section 3. The classical inputs are the Matomäki–Radziwiłł–Tao short exponential-sum and distance bounds [16, Theorem 1.7 and (1.12), p. 6], the fundamental lemma [6, Theorem 3.6(a), p. 38], Liouville Bombieri–Vinogradov over all residues [23, (1.5), p. 2], Shiu’s positive mean bound [12, p. 1], the Nair–Tenenbaum upper theorem [12, Theorem 1, pp. 1–2], bounded independence [1, 2, 25], Halász in the form [29, Chapter III.4], and the fixed-modulus prime number theorem and Mertens’ theorem [19]. The zero-free half-plane $\Re s > 7/8$ is not used in any of the four theorems. The classical zero-free region enters Theorem 1.4, as well as the other signed graph estimates, through the cited MRT distance bound.

The roughness of the target vertex is put into the padding probability, with row cutoff KW_1/L . Different paddings retain three local densities; equal paddings require a separate diagonal term with exponent $24/17$. Closed walks are rewritten using arrival indicators before the padding sum. The resulting truncation event is a depth-three Boolean circuit; bounded independence compares it with the independent residue model. For soft weights, $\sqrt{b}$ is placed in the operator and $\sqrt{b}$ in the test vector.

The shift h is fixed throughout; the statements have no exceptional set of X or of scales. All absolute constants in the theorems and their thresholds $X_0(h)$ (or $X_0(h, r)$ for the fixed soft weight) are ineffective. No statement here prescribes a fixed number of prime factors or asserts primality. One remaining two-point estimate toward a fixed number of prime factors is $|\sum_{n \leq X} \lambda(n)\lambda(n + h)J_{\sigma}(n)J_{\sigma}(n + h)| \leq (1 - 2\epsilon) \sum_{n \leq X} J_{\sigma}(n)J_{\sigma}(n + h)$, where $J_{\sigma}(n) = \prod_{p|n} (1 - p^{-\sigma})$, $\sigma = c/\log X$, $0 < \epsilon \leq 1/4$ and c is fixed sufficiently large in terms of ϵ .

Section 2 specifies the inputs and graph data. Section 3 proves finite comparison. Sections 4–6 prove Theorem 1.1. Section 7 treats the balanced soft weight. Section 8 proves Theorem 1.2 and Remark 1.5. Sections 9–11 prove Theorem 1.4.

2 Notation, classical inputs and graph data

Write $e(\theta) = \exp(2\pi i\theta)$, τ_k for the k -fold divisor function, $\tau = \tau_2$, and $P(z) = \prod_{p \leq z} p$. Let μ be the Möbius function, φ Euler's totient, and ζ the Riemann zeta function. Write $P^+(n)$ for the largest prime factor, with $P^+(1) = 1$. Powers with exponent zero mean one, also at $r = 0$. In an independent residue model there is one uniform coordinate R_s modulo each s in a pairwise coprime family. Write $\mathbb{P}_*$ and $\mathbb{E}_*$ for probability and expectation under this product law, and $\mathbb{P}_I$ for the law induced by a uniformly chosen integer in a finite interval I of consecutive integers. Every translated site uses the same coordinates with the indicated offsets. Translated sites are not independent. Write $\rho(B)$ for the spectral radius of a finite matrix B ; the Hilbert–Schmidt norm is unnormalized. For a finite prime set $\mathcal{T}$, put $\omega_{\mathcal{T}}(n) = \sum_{p \in \mathcal{T}} \mathbf{1}_{p|n}$. The same divisibility notation is used for a residue site. At integer or residue sites, the rough mask $R_z(n)$ means $\prod_{p \leq z} \mathbf{1}_{p \nmid n}$.

Lemma 2.1 (Rough densities and exceptional primes). *Let $W_1(z) = \prod_{p \leq z} (1 - 1/p)$ and $v = \log(2z)$. Then*

$$\begin{aligned} W_1(z) &\asymp v^{-1}, & V_h(z) &\asymp_h W_1(z)^2, \\ V_h(z) &\geq c_* W_1(z)^2, & W_1(z)/V_h(z) &\ll v, \\ c_* &= \prod_{p > 2} \left(1 - \frac{1}{(p-1)^2}\right) > 0. \end{aligned} \tag{2.1}$$

For a nonzero integer Δ ,

$$\sum_{p|\Delta} \frac{1}{p} \leq \log \log \log(3|\Delta|) + O(1), \quad \prod_{\substack{p > 3 \\ p|\Delta}} (1 - 3/p)^{-1} \ll (\log \log(3|\Delta|))^6. \tag{2.2}$$

Proof. Mertens' theorem [19, Section 2.2] gives the assertion about W_1 . For $p \nmid h$, $(1 - 2/p)/(1 - 1/p)^2 = 1 - 1/(p-1)^2$; for $p \mid h$ the ratio is $(1 - 1/p)^{-1}$. Since $2 \mid h$, the factor at 2 is positive and all other exceptional factors form a fixed finite product. This proves every comparison in (2.1). For (2.2), separate primes at $\log(3|\Delta|)$. The smaller ones contribute at most $\log \log \log(3|\Delta|) + O(1)$. There are at most $\log(3|\Delta|)/\log \log(3|\Delta|)$ larger prime divisors, and their reciprocal sum is bounded. Finally $-\log(1 - 3/p) \leq 6/p$ for $p \geq 5$. Exponentiation gives the product bound. $\square$

Input 2.2 (Fundamental lemma). *For a sieve problem with multiplicative densities $0 \leq g(p) < 1$ satisfying the product condition of dimension at most two, there are upper and lower divisor weights $\xi_d^\pm$, $|\xi_d^\pm| \leq 1$, supported on squarefree $d \mid P(z)$, $d \leq D$, such that the corresponding divisor sums bound the sifted indicator above and below and*

$$\left| \sum_d \xi_d^\pm g(d) - \prod_{p \leq z} (1 - g(p)) \right| \leq E(s) \prod_{p \leq z} (1 - g(p)), \quad s = \frac{\log D}{\log z}, \quad E(s) \ll e^{-cs} \tag{2.3}$$

for sufficiently large s , when $2 \leq z \leq D^{1/2}$. The constants are uniform for subsets of the forbidden roots $0, -h$. This is [6, Theorem 3.6(a), p. 38; probability-space formulation, pp. 1–3]; its sharper error is $\exp(-s \log s + s \log_3 s + O(s))$. Indeed $g(p) \leq \nu_h(p)/p$, and the associated product condition is bounded by the dimension-two product. In particular its constants can be absolute for even h .

Lemma 2.3 (Maximal all-residue Liouville bound). *For fixed $\vartheta < 1/2$ and every $A > 0$,*

$$\sum_{d \leq T^\vartheta} \max_{a \bmod d} \max_{y \leq T} \left| \sum_{\substack{n \leq y \\ n \equiv a \pmod{d}}} \lambda(n) \right| \ll_{A, \vartheta} T(\log T)^{-A}. \quad (2.4)$$

The maximum includes nonunit classes. For $D \leq X^{1/4}$, with E_d denoting this maximum at $T = 2X$,

$$\sum_{d \leq D} \mu^2(d) \nu_h(d) E_d \leq \left(\sum_{d \leq D} E_d \right)^{1/2} \left(\sum_{d \leq D} \mu^2(d) 4^{\omega(d)} E_d \right)^{1/2} \ll_{h, B} X(\log X)^{-B} \quad (2.5)$$

for every $B > 0$.

Proof. The bound without the maximum over y is [23, (1.5), p. 2]. Fix $\vartheta < \vartheta' < 1/2$ and an integer $m > A + 3$. Use $O(\log^m T)$ cutoffs with spacing at most $T/\log^m T$. At grid points $y \geq T/(2\log^m T)$ one has $T^\vartheta \leq y^{\vartheta'}$ and $\log y \asymp \log T$. The cited estimate with saving $A + m + 3$, summed over the grid, is $O(T \log^{-A-2} T)$. Interpolation and the initial small interval cost $O(T \log^{1-m} T + T^\vartheta)$, by the progression term count $T/(d \log^m T) + 1$. This proves (2.4). Since $\nu_h(d) \leq 2^{\omega(d)}$ on squarefree integers, Cauchy–Schwarz gives the inequality in (2.5). Also $E_d \leq 2X/d + 1$ and

$$\sum_{d \leq D} \frac{4^{\omega(d)}}{d} \leq (1 + \log D)^4, \quad \sum_{d \leq D} 4^{\omega(d)} \ll D(1 + \log D)^3. \quad (2.6)$$

These follow from $4^{\omega(d)} \leq \tau_4(d)$ by summing its four divisor variables. The second square in (2.5) is $O(X \log^4(2X))$; arbitrary A in (2.4) finishes the proof. Replacing n by $n + h$ costs two prefixes, once $X \geq h$. $\square$

Lemma 2.4 (Unsigned and singly signed rough sums). *For each fixed $0 < \rho_* < 1$, uniformly for $2 \leq z \leq \exp((\log X)^{\rho_*})$,*

$$\sum_{n \leq X} R_z(n) R_z(n + h) = X V_h(z) + O_{h, \rho_*, B}(X \log^{-B} X), \quad (2.7)$$

$$\sum_{n \leq X} \lambda(n) R_z(n) R_z(n + h), \quad \sum_{n \leq X} \lambda(n + h) R_z(n) R_z(n + h) \ll_{h, \rho_*, B} X \log^{-B} X. \quad (2.8)$$

Both assertions hold for every fixed $B > 0$.

Proof. Put $D = X^{1/4}$, $g_h(d) = \nu_h(d)/d$ on squarefree d , and $w^\pm(n) = \sum_{d|n(n+h)} \xi_d^\pm$. CRT gives

$$\sum_{n \leq X} \mathbf{1}_{d|n(n+h)} = X g_h(d) + O(\nu_h(d)). \quad (2.9)$$

Since $\sum_{d \leq D} \mu^2(d) \nu_h(d) \ll D \log(2D)$, Input 2.2 yields

$$\sum_{n \leq X} R_z(n) R_z(n + h) = X V_h(z) + O(E(s) X V_h(z) + D \log(2D)), \quad (2.10)$$

$$\sum_{n \leq X} |w^+(n) - R_z(n) R_z(n + h)| \leq 2E(s) X V_h(z) + O(D \log(2D)), \quad (2.11)$$

$$s \geq \frac{1}{4} (\log X)^{1-\rho_*}.$$

For large X , $z \leq D^{1/2}$, $g_h(p) < 1$ because h is even, and $E(s)$ beats every logarithmic power. The second bound follows from $w^+ \geq R_z R_z \geq w^-$. It is an L^1 bound and may be multiplied by either Liouville sign. Expanding w^+ and applying (2.5) proves the two signed bounds. $\square$

Input 2.5 (Short Fourier means and distance). *For a multiplicative g with $|g| \leq 1$, $10 \leq D \leq Y$, and $Q = \min((\log Y)^{1/125}, (\log D)^5)$, define*

$$M(g; Y, Q) = \min_{\substack{q \leq Q, \chi \bmod q \\ |t| \leq Y}} \sum_{p \leq Y} \frac{1 - \Re(g(p) \overline{\chi(p)} p^{-it})}{p}. \quad (2.12)$$

Then

$$\sup_{\theta} \frac{1}{Y} \int_0^Y \left| \sum_{x \leq n \leq x+D} g(n) e(\theta n) \right| dx \ll D \left(e^{-M(g; Y, Q)/20} + \frac{\log \log D}{\log D} + (\log Y)^{-1/700} \right). \quad (2.13)$$

Moreover in this character and height range

$$M(\lambda; Y, Q) \geq \frac{1}{4} \log \log Y - O(1). \quad (2.14)$$

These are [16, Theorem 1.7 and (1.12), p. 6], with a conservative choice of the constant in (1.12). For the distance assertion one can apply (1.12) with auxiliary long and short parameters both Y and its coefficient bound 1, giving the distance at prime cutoff $10Y$ for every $q \leq Q$ and $|t| \leq Y$. Removing the primes in $(Y, 10Y]$ costs at most $2 \sum_{Y < p \leq 10Y} 1/p = O(1)$, giving (2.14). The supremum in (2.13) is outside the origin integral. The distance bound in the cited version is deduced from the classical Vinogradov–Korobov zero-free region.

Definition 2.6 (Labels, bins and vertex cuts). Fix $W \geq 10$, $J = \lfloor \alpha \log L \rfloor \geq 1$, with $\alpha = 1/(1200W)$, and $0 < \eta \leq 1$. Let $H_0 = \exp(L^{199/200})$, $Y_i = L^{199/200} e^{6W_i}$, and

$$\mathcal{P}_i = \{p \equiv 1 \pmod{5} : Y_{i-1} \leq \log p < Y_i, p \nmid h\}, \quad \mathcal{P} = \bigcup_{i=1}^J \mathcal{P}_i, \quad \mathcal{D} = \{p_1 \cdots p_J : p_i \in \mathcal{P}_i\}. \quad (2.15)$$

At a small-prime cutoff z (or y) below H_0 take

$$\begin{aligned} \mathcal{Q} &= \{p : z < p \leq e^L, p \not\equiv 1 \pmod{5}, p \nmid h\}, \\ a(u) &= a^{\omega(u)}, \quad w(n) = (1+a)^{\omega_{\mathcal{Q}}(n)}, \quad S = \prod_{p \in \mathcal{Q}} (1+a/p). \end{aligned} \quad (2.16)$$

where $a = 4$ or 16 , and $\mathcal{Q}^{\text{sf}}$ is the finite set of squarefree products of primes in $\mathcal{Q}$, including 1. Set

$$\mathcal{S}_j = \{(d, u) \in \mathcal{D} \times \mathcal{Q}^{\text{sf}} : \omega(u) \leq 100 \log L, j\eta \leq \log(du) < (j+1)\eta\}, \quad T_j = X e^{j\eta}, \quad j\eta \leq 100L. \quad (2.17)$$

Write $V_i = \sum_{p \in \mathcal{P}_i} 1/p$, $V_* = \prod_i V_i$, $S_0 = \sum_j \sum_{(d,u) \in \mathcal{S}_j} a(u)/(du)$, and $k = \lfloor L \rfloor$, $\ell_0 = \lfloor L^{1/10} \rfloor$, $M = \lceil e^{103L} \rceil$. The exterior projection E restricts to $\omega_{\mathcal{P}}(n) \leq 6WJ$.

Proposition 2.7 (Label supply and retained mass). *With the fixed data above, for sufficiently large L ,*

$$W \leq V_i \leq 2W, \quad \log d < 2L, \quad \frac{1}{2} S V_* \leq S_0 \leq S V_*, \quad X e^{-\eta} < T_j/(du) \leq X. \quad (2.18)$$

Proof. Since $6WJ \leq (1/200)\log L$, the supplies lie below e^L . The fixed-modulus prime number theorem [20, (2.2)–(2.3), pp. 8–9] gives $V_i = (3/2)W + o(1)$ uniformly in i ; removing the fixed divisors of h has bounded, eventually vanishing effect. The geometric sum of Y_i is less than $L/(1 - e^{-6W}) < 2L$. Under the law $a(u)/(uS)$, a prime occurs with probability $a/(p+a)$. For $a = 4$ the expected logarithmic size and degree are at most $4L$ and $4\log L$. For $a = 16$ the bounds $17L$ and $17\log L$ suffice for large L , also after removing primes up to z . Markov excludes $\log u > 98L$ and $\omega(u) > 100\log L$ with probabilities totaling at most $4/98 + 4/100 < 1/2$ or $17/98 + 17/100 < 1/2$, respectively. Every remaining u belongs to a bin with every d . This repeats the two exclusions in [20, Lemma 2.2, pp. 8–9], with the present harmonic law proved explicitly. Dropping restrictions gives the upper bound. The last assertion follows directly from (2.17). $\square$

Definition 2.8 (Rare vertices). A signed numerical step is $D_i = \epsilon_i h d_i u_i$, $\epsilon_i \in \{\pm 1\}$, $(d_i, u_i) \in \mathcal{S}_j$. A path is positive when $d_i u_i$ divides its departure at each step; this also holds at the arrival. A forward prohibited word has length $3 \leq m \leq \ell_0$, unequal consecutive whole tuples, interval-shaped occurrences of every centre prime, and a prime $p \mid d_1$, $p \nmid d_m$, such that

$$p \mid \sum_{i=b}^m D_i \quad \text{for some } 1 < b < m. \quad (2.19)$$

It is minimal if no shorter contiguous subword is forward prohibited in either orientation. Let $\mathcal{B}_j$ be the sites where a positive minimal prohibited word starts, and $\sigma_j = \mathbf{1}_{n \notin \mathcal{B}_j}$. These are the definitions of [20, Definition 5.1, pp. 21–22] with the present label supply.

Proposition 2.9 (Rare-site estimate for the changed padding). *For either padding weight in (2.16),*

$$\mathbb{P}_*(n \in \mathcal{B}_j) \leq \exp(-\tfrac{1}{2}L^{199/200}). \quad (2.20)$$

Its positivity event is a disjunction of at most $\exp(O(\ell_0 L))$ conjunctions, each with $O(\ell_0 \log L)$ residue equalities.

Proof. Follow [20, Lemmas 5.2–5.3, pp. 22–23]. A word of at most $R \leq 4L$ steps has $O(R \log L)$ prime slots, so its equality patterns and polynomially many records per slot number $\exp(O(R \log^2 L))$. Each distinct prime contributes one reciprocal and harmonic mass $O(\log L)$. Now $La(u) \leq L^{1+100\log a}$; this changes only the absolute constant in that count. In a prohibited word, a centre prime z' entering on the last transition has coefficient $\epsilon_m h u_m d_m / z'$ in (2.19). The controlling prime p is absent from d_m , divides neither h nor any padding, and so that coefficient is a unit modulo p , independent of z' . With the other labels fixed the reciprocal sum for z' is bounded by

$$\sum_{\substack{H_0 \leq n \leq e^L \\ n \equiv c \pmod{p}}} \frac{1}{n} \leq \frac{1}{H_0} + \frac{L}{p} \leq \frac{1+L}{H_0}. \quad (2.21)$$

The remaining positive reciprocal sum is $\exp(O(\ell_0 \log^2 L))$. Thus the probability is at most $(1 + L)H_0^{-1} \exp(O(\ell_0 \log^2 L))$, giving (2.20). Numerical pairs have $du \leq e^{101L}$ and are uniquely recovered from their product by disjoint supports. Preselect the numerical prohibited and minimal words; positivity alone is a conjunction of the stated size. This proves the Boolean description and translation covariance. No Liouville values enter this event. $\square$

3 Finite residue comparison

Lemma 3.1 (Shallow CNF comparison). *Fix an integer $\ell \geq 1$. Let the coordinate moduli be pairwise coprime integers between 2 and e^L , all coprime to ℓ , with the coordinate modulo ℓ added if $\ell > 1$. An event given by a residue CNF with at most e^{L^3} clauses, each of width at most L , has*

$$|\mathbb{P}_I(E) - \mathbb{P}_*(E)| \leq e^{-L^6} \quad (|I| \geq e^{L^{18/2}}). \quad (3.1)$$

The literals may be residue equalities or their negations, with repetitions and arbitrary offsets. An expansion of such events with total absolute coefficient mass e^{CL^4} has error at most e^{-L^5} . The threshold depends only on ℓ, C .

Proof. Bazzi's bounded-independence theorem [1], in the form of [25, Theorem 1, p. 2], bounds the error for an m -clause CNF under exactly t -wise uniform bits by $m^{C_0} e^{-c_0 \sqrt{t}}$; complementation gives CNFs from DNFs. We give the reduction to that precise hypothesis.

Put $B_0 = \lceil L^8 \rceil$. Adjoin an independent U_s uniform on $[0, 1)$ at every coordinate, and encode $K_s = \lfloor 2^{B_0}(R_s + U_s)/s \rfloor$ in B_0 bits. Decode by

$$\hat{R}_s = \left\lfloor \frac{s(K_s + 1/2)}{2^{B_0}} \right\rfloor, \quad \mathbb{P}(\exists s : \hat{R}_s \neq R_s) \leq 2^{1-B_0} \left(\ell + \sum_s s \right) \leq e^{3L} 2^{-B_0}. \quad (3.2)$$

Conditional on any residue, only its two boundary dyadic cells can cause failure, of probability at most $2s2^{-B_0}$. Under the product law all encoded bits are independent and unbiased. The monotone decoder makes every residue preimage an interval of binary integers. An interval has a CNF with at most $2B_0$ clauses, by excluding each forbidden initial differing bit at each boundary. Its complement has at most B_0^2 clauses after distributing the two one-sided CNFs. Thus a residue literal has at most $2B_0^2$ clauses. Distributing a clause of width at most L gives a Boolean CNF with

$$m \leq e^{L^3} (2B_0^2)^L \leq e^{L^4}, \quad n_b \leq e^{2L} \quad (3.3)$$

clauses and bits, respectively, for large L .

Take $t = \lceil L^{13} \rceil$. At most t selected bits read at most t coordinates, whose product is at most ℓe^{tL} . CRT counts each residue with error $O(1)$, so their total variation from uniform is at most that product divided by $|I|$, and hence at most $\Delta = e^{2tL-L^{18/2}}$. Jitter and projection do not increase this distance. If f is the full bit-law density relative to the uniform cube, its Walsh coefficients satisfy $|\hat{f}(S)| \leq 2\Delta$ for $1 \leq |S| \leq t$. Consequently

$$q := \sum_{1 \leq |S| \leq t} |\hat{f}(S)| \leq 2\Delta(t+1)n_b^t \leq e^{-L^{17}}. \quad (3.4)$$

Let $H = \sum_{1 \leq |S| \leq t} \hat{f}(S) \chi_S$ and $a_0 = e^{-2L^6}$. Since $|H| \leq q \leq a_0$,

$$g = \frac{f - H + a_0}{1 + a_0} \geq 0, \quad \mathbb{E}g = 1, \quad d_{\text{TV}}(f, g) \leq \frac{q + a_0 \mathbb{E}|1 - f|}{2(1 + a_0)} \leq 3a_0/2. \quad (3.5)$$

All nonconstant Walsh coefficients of order at most t vanish for g , so Fourier inversion on each marginal makes g exactly t -wise uniform. Bazzi's error is at most $\exp(C_0 L^4 - c_0 L^{13/2}) \leq a_0$. Adding the correction and decoding errors gives $2e^{3L} 2^{-B_0} + 5a_0/2 \leq e^{-L^6}$. Multiplication by e^{CL^4} gives the scalar assertion. The argument used only the coordinate count, sizes and CRT, so it applies to prime-square coordinates as well as primes. $\square$

Proposition 3.2 (Encoding the graph events). *For padding $4^{\omega(u)}$, any padding-prime subset as above, and any fixed-power choices of η, K , every comparison used in positive deletions, the signed trace, and the truncated testing norm is a scalar expansion covered by Lemma 3.1, of total absolute coefficient mass $\exp(O(L^4))$. Small-prime rough indicators or prime-square negative tests may be adjoined.*

Proof. This replaces [20, Lemma 3.1 and Corollaries 3.2–3.3, pp. 10–12] for the present objects. Follow the event expansions of [20, Lemma 4.6, pp. 19–21; Theorem 5.4, pp. 24–25; Proposition 8.4, pp. 41–42]. A numerical witness is preselected by its numerical arithmetic and minimality conditions. Its positivity is a conjunction of $O(L^{1/10} \log L) < L$ equalities. There are $\exp(O(L^{11/10}))$ witnesses at one site; their absence is a CNF. Conjoining these tests at at most $2k \leq 2L$ visited sites still gives fewer than e^{L^3} clauses.

An exact active- $\mathcal{Q}$ state is a conjunction of all presences and all absences, with unit-width clauses. After the cutoff $\omega_{\mathcal{Q}} \leq \lfloor 400 \log L \rfloor$, there are $\exp(O(L \log L))$ possible states at a site. In each, denominators and the original padding cutoff are scalars. A degree condition $\omega_{\mathcal{T}} \leq b$, $b = O(\log L)$, excludes each simultaneous $(b+1)$ -subset and is a CNF of $\exp(O(L \log L))$ clauses. Roughness and fixed divisibilities add unit clauses.

In a trace, fix a signed numerical word and the two external copy indices. Enumerate its states and expand its $2kJ$ centred factors into indicators and scalars, retaining their signs. The resulting event is the conjunction just described. There are $\exp(O(kL \log L))$ states, at most 2^{2kJ} centred expansions, scalar sizes $\exp(O(k \log L))$, and $\exp(O(kL))$ numerical words and external indices. Their total mass is $\exp(O(L^2 \log L)) \leq \exp(O(L^4))$. This comparison precedes absolute values of word expectations.

For deletion, expand the positive centred majorant and split the four failed cuts at both endpoints by the exact union bound. A large-degree or witness event B may be treated by

$$\mathbf{1}_{H \cap B} = \mathbf{1}_H - \mathbf{1}_{H \cap B^c}. \quad (3.6)$$

Both right-hand events have the required CNFs. This avoids an expansion over intersections of arbitrary witness families. Eligible pairs number $e^{O(L)}$ and exact states cost $e^{O(L \log L)}$, within the same scalar bound. Finally the truncated norm is expanded by its exact active state, of coefficient $5^{|\mathcal{Q}_n|}$, at most a fixed power of L . Its $M \leq e^{104L}$ block positions preserve the bound. Apply Lemma 3.1 to all expansions. None reads the full Ω weight or a Liouville value. $\square$

Proposition 3.3 (Rough-target threshold comparison). *For padding $16^{\omega(u)}$ and $w = 17^{\omega_{\mathcal{Q}}}$, replace the vertex degree cutoff by $\lfloor 18 \log L \rfloor$. The norm, positive deletion and signed-trace comparisons with the rough-target threshold defined in (4.1) have per-event error $e^{-L^{10}}$ and aggregate error e^{-L^9} on any origin interval of length at least $e^{L^{625}/2}$. These estimates are uniform in $z \leq e^{L^{1/6}}$, bin position and numerical word data, for fixed h .*

Proof. On an exact active state there are at most $N = 2^{18 \log L} = L^{18 \log 2} < L^{13}$ possible padding divisors. A weighted sum of their target rough indicators exceeds its threshold exactly when some subset of candidates whose scalar weights exceed the threshold has all its targets rough. Negating this statement gives an AND over those subsets of an OR of positive residue equalities: one selected target has a small prime divisor. There are at most 2^N clauses, each of width at most $N\pi(z)$. Both orientations, all $O(L)$ visited sites, exact states, no-witness predicates and endpoint masks give a residue CNF whose logarithmic size is $O(L^{18 \log 2}) + O(L^{1/6}) + O(L \log L)$. Use (3.6) for positive bad-event costs. The subsets are internal to this circuit, rather than scalar summands.

Use the monotone decoder (3.2) with $B_0 = \lceil L^{15} \rceil$. Each binary interval has a DNF with at most $2B_0$ dyadic-prefix conjunctions, by following its two boundary paths in the binary tree and

taking the included subtrees. Its complement has $O(B_0)$ such conjunctions too. Substitution into the residue CNF and flattening adjacent OR levels gives AND–OR–AND, of depth three and size at most $e^{L^{14}}$, including input occurrences. Decoding error is charged per coordinate: one correct decode settles every offset test at that coordinate.

Braverman’s exact independence bound [2, Corollary 2, p. 1] for depth d , size m and error ϵ_0 is

$$3 \cdot 60^{d+3} (\log m)^{(d+1)(d+3)} \{5 \log(15m/\epsilon_0)\}^{d(d+3)}. \quad (3.7)$$

At $d = 3$, $m \leq e^{L^{14}}$ and $\epsilon_0 = e^{-L^{11}}$, this is $O(L^{588})$. Equivalently the Main Theorem there may be applied with

$$s_B = \lceil 100L^{14} \rceil, \quad 0.82^{s_B}(15m) \leq e^{-L^{11}} \quad (3.8)$$

for large L ; the same exponent follows independently of the logarithm-base convention. Take $t = \lceil L^{589} \rceil$. At most t bits involve moduli of product $e^{O(tL)}$, and the total bit count is at most e^{2L} . CRT and the low-order Walsh coefficient count therefore cost $\exp(-L^{625}/2 + O(L^{590})) \leq e^{-L^{12}}$. Apply the nonnegative correction (3.5) with $a_0 = e^{-L^{11}}$ to obtain exact t -wise uniformity. Its error is $O(e^{-L^{11}})$; adding Braverman’s error and the coordinate decoding error $e^{3L}2^{-L^{15}}$ gives $e^{-L^{10}}$. The states, words and factors $L16^{\omega(u)}$ retain the scalar mass $e^{O(L^4)}$ proved above. Thus their aggregate error is e^{-L^9} . The large threshold is not covered merely by Lemma 3.1. $\square$

4 Rough-target padding and its second moment

Use $a = 16$ in Definition 2.6, and suppose $2 \leq z \leq e^{L^{1/6}}$. For a numerical d whose primes exceed z and are outside $\mathcal{Q}$, $\log d \leq 2L$, define

$$\rho_{j,z}^{\pm}(n, d) = \frac{1}{w(n)} \sum_{\substack{u|n, u \in \mathcal{Q}^{\text{sf}} \\ j\eta \leq \log(du) < (j+1)\eta}} 16^{\omega(u)} R_z(n \pm hdu), \quad \rho_{j,z} = \max(\rho_{j,z}^+, \rho_{j,z}^-). \quad (4.1)$$

The sum here includes all padding divisors in the bin, before the eligible degree restriction.

Proposition 4.1 (Three-density second moment). *For $0 < \eta \leq 1$ and the bins $j\eta \leq 100L$, uniformly in z, η, d ,*

$$\mathbb{E}_* \left[w(n) R_z(n) \sum_j \rho_{j,z}(n, d)^2 \right] \ll_h \frac{SW_1^3(1 + \log L)^6}{L}. \quad (4.2)$$

Deleting a row with $\rho_{j,z} > KW_1/L$, after the low- $\mathcal{Q}$ cutoff, costs, after summing bins and labels and normalizing by SV_ , at most*

$$C_h W_1^2 \frac{2^J(1 + \log L)^6}{K}. \quad (4.3)$$

Proof. Following the conditional-divisor experiment of [20, Lemma 4.5, pp. 17–18], tilt by w/S . Each $\mathcal{Q}$ prime is then active with probability $17/(p+16)$, and conditional on activity is selected in a divisor with probability $16/17$. Two conditionally independent selected divisors u_1, u_2 give the factor

$$1 - \frac{32}{17(p+16)}(1 - \cos(t \log p)) \quad (4.4)$$

for their log-ratio characteristic function. It is nonnegative for every real t ; the harmonic estimate below is used only for $|t| \leq 2$. Fixed-modulus PNT and partial summation, as in [20, (4.11)–(4.13), pp. 17–18], give

$$0 \leq \phi_z(t) \ll_h \left(\frac{1+v|t|}{1+L|t|} \right)^{24/17}, \quad v = \log(2z) \leq 2L^{1/6}, \quad |t| \leq 2. \quad (4.5)$$

Indeed the harmonic sum of $1 - \cos(t \log p)$ on these padding primes is $\frac{3}{4} \log((1+L|t|)/(1+v|t|)) + O_h(1)$; replacing $p+16$ by p changes it by $O(1)$. The coefficient $(32/17)(3/4)$ is $24/17$. The nonnegative triangular Fourier kernel

$$\kappa(x) = \left(\frac{\sin x}{x} \right)^2 = \frac{1}{4} \int_{-2}^2 (2-|t|) e^{itx} dt, \quad \mathbf{1}_{[-1,1]}(x) \leq \kappa(x)/(\sin 1)^2 \quad (4.6)$$

bounds bin collision. If ρ_j denotes the unmasked bin probability, it gives

$$\mathbb{E}^{\text{tilt}} \sum_j \rho_j^2 \ll_h L^{-1} + (v/L)^{24/17} \ll_h L^{-1}, \quad \mathbb{E}^{\text{tilt}} F = \mathbb{E}_*(wF)/S. \quad (4.7)$$

The exact equality probability is

$$D_Q = \prod_{p \in Q} \left(1 - \frac{32}{17(p+16)} \right) \ll_h (v/L)^{24/17}. \quad (4.8)$$

Fix an orientation and condition on the active padding configuration and u_1, u_2 . For equal paddings, the two small-prime sites have mean $V_h(z) \ll_h W_1^2$, since du is a unit at every small prime. For different eligible paddings, $0 < |u_1 - u_2| \leq e^{101L}$. At each $p > 3$ not dividing $h(u_1 - u_2)$, the three offsets $0, \pm hdu_1, \pm hdu_2$ are distinct modulo p . Dropping the other factors bounds their common roughness probability by

$$\prod_{3 < p \leq z} (1 - 3/p) \prod_{\substack{p > 3 \\ p | h(u_1 - u_2)}} (1 - 3/p)^{-1} \ll_h W_1^3 (1 + \log L)^6. \quad (4.9)$$

Use Lemma 2.1; if the forbidden roots cover a whole small field, the probability is zero. Apply this bound before enlarging the unmasked collision sum, so the difference bound remains valid. Enlarge only the diagonal to all equal divisor choices. Since $\max(x, y)^2 \leq x^2 + y^2$, the left side of (4.2) is bounded by a constant times

$$S \left\{ \frac{W_1^3 (1 + \log L)^6}{L} + W_1^2 (v/L)^{24/17} \right\}, \quad \frac{W_1^2 (v/L)^{24/17}}{W_1^3/L} \ll_h v^{41/17} L^{-7/17} \ll_h L^{-1/102}. \quad (4.10)$$

The last exponent is $(1/6)(41/17) - 7/17 = -1/102$. This proves (4.2).

On a bad row, the positive source edge mass $wR_z \rho_{j,z}^+$ is at most $LwR_z \rho_{j,z}^2/(KW_1)$. The centre majorant $a_d^+(n) = \prod_{p|d} (\mathbf{1}_{p|n} + 1/p)$ has mean $2^J/d$ independently of these coordinates. Sum d to obtain $2^J V_*$ and use (4.2). Reversing the orientation proves the arrival bound. This gives (4.3), without independence between translated sites. $\square$

5 The modified graph moment and spectral estimate

For the rough operator let

$$\tau_j(n, d) = \mathbf{1}_{\omega_Q(n) \leq \lfloor 18 \log L \rfloor} \mathbf{1}_{\rho_{j,z}(n, d) \leq KW_1/L} \sigma_j(n), \quad c_d(n) = \prod_{p|d} (\mathbf{1}_{p|n} - 1/p), \quad a_d(n) = |c_d(n)|. \quad (5.1)$$

On M consecutive sites define a symmetric matrix, for $m = n + hdu$ with $(d, u) \in \mathcal{S}_j$, by

$$A_d(n, m) = \frac{L16^{\omega(u)}}{\sqrt{w(n)w(m)}} \mathbf{1}_{u|n} c_d(n) \tau_j(n, d) \tau_j(m, d) R_z(n) R_z(m), \quad A_d(m, n) = A_d(n, m). \quad (5.2)$$

Other entries vanish. On the direct sum of d -indexed copies set $H_{d,d'} = \mathbf{1}_{d \neq d'} A_{d'}$. The $\mathcal{P}$ -degree restriction occurs only in E .

For endpoint functions f, g and a numerical sign $\epsilon(du)$ define the centred sums

$$C_j(f, g) = \frac{1}{T_j} \sum_{n \leq T_j} \sum_{(d, u) \in \mathcal{S}_j} a(u) \mathbf{1}_{u|n} c_d(n) \epsilon(du) f(n) g(n + hdu). \quad (5.3)$$

Their retained versions $C_j^\circ(f, g)$ insert $\chi_j(n, d) \chi_j(n + hdu, d)$, where $\chi_j = \tau_j \mathbf{1}_{\omega_P \leq 6WJ}$. For the rough case take $f = g = \lambda R_z$, $a = 16$ and $\epsilon = 1$. For the soft and full weights the endpoint functions, numerical sign, and vertex cuts are specified below.

Input 5.1 (Finite-dimensional transfer, [20, Lemma 8.1, p. 39]). *Let B_i be self-adjoint operators on a finite-dimensional complex Hilbert space, P an orthogonal projection, and $\mathcal{H}_{i,j} = \mathbf{1}_{i \neq j} B_j$. If $a, b \geq 0$, $\|B_i\| \leq a$, and $\sum_i \|B_i v\|^2 \leq b^2 \|v\|^2$ for $v = Pv$, then*

$$\left\| P \left(\sum_i B_i \right) P \right\| \leq 3 \max(a, b, \rho(\mathcal{H})). \quad (5.4)$$

Neither different B_i nor P need commute.

Proposition 5.2 (Moment with the rough density). *There are absolute constants C_2, C , fixed before W , for padding 16, with the following property. For each fixed even $h \geq 2$ and fixed $W \geq 10$, take X sufficiently large in terms of h, W . Use the operator (5.2), $\alpha = 1/(1200W)$, and*

$$L = (\log X)^{1/625}, \quad 2 \leq z \leq e^{L^{1/6}}, \quad J = \lfloor \alpha \log L \rfloor, \quad \eta = L^{-\alpha/100}, \quad K = L^{\alpha \log 2 + \alpha/100}. \quad (5.5)$$

Then

$$\mathbb{E}_* \|H^k\|_{\text{HS}}^2 \leq \{KW_1(C_2 \sqrt{W})^J\}^{2k}. \quad (5.6)$$

The same bound holds for finite origin intervals of length at least $e^{L^{625}/2}$, after one absolute enlargement of C_2 . Outside a fraction at most e^{-2k} of those origins,

$$\left\| E \left(\sum_d A_d \right) E \right\| \leq KW_1(C \sqrt{W})^J. \quad (5.7)$$

Proof. We check the proof of [20, Theorem 5.4, pp. 24–38] in its order. Write $\mathbf{b} = KW_1$. Expanding $\|H^k\|_{\text{HS}}^2$ joins two paths with common endpoints and reverses one. The result is a closed word of $2k$ steps with unequal consecutive whole tuples within each half. There are at most $|\mathcal{D}|^2 M \leq e^{108L}$

external choices, as in [O, (5.4), p. 24]; there is no additional copy choice per step. Proposition 3.3 compares this signed expansion before taking absolute values.

On a closed word the product of the two endpoint indicators is the product of the arrival indicators, counting visits with multiplicity, because $R_z^2 = R_z$. Its square-root denominators multiply to one w at each departure. Make these identities before dropping closure. The contribution now has the form [O, (5.6)–(5.7), p. 25], with $R \geq 0$ independent of all $\mathcal{P}$ residue draws, and $G = \prod_i \sigma_j(n_i)$. The new masks and cutoff may depend on numerical centre labels, but not on their residue draws; they remain in R . Only G contains the rare-witness predicates. This verifies the decisive hypothesis of [O, Lemma 5.5, p. 26]. A centre prime appearing in exactly one step is a singleton. At every nonsingleton occurrence expand the centred factor into its indicator (lit) or its scalar $-1/p$ (unlit). Lit consistency requires all the lit departure residues of one prime to agree. Force the common residue for each lit nonsingleton. For each subset H of the singleton set $\mathcal{U}$, also force the residue at the sole occurrence of each singleton in H , leaving the other coordinates at their original draws. Write G_H for G on this hybrid and define

$$\Delta G = \sum_{H \subset \mathcal{U}} (-1)^{S_1 - |H|} G_H, \quad S_1 = |\mathcal{U}|. \quad (5.8)$$

In particular, at a singleton prime,

$$\mathbb{E}[(\mathbf{1}_{Z=a} - 1/p)F(Z)] = p^{-1}\mathbb{E}[F(a) - F(Z)]. \quad (5.9)$$

Integrating all singleton coordinates gives the same mixed difference ΔG , bounded by 2^{S_1} , and the reciprocal factor

$$b_{\mathcal{L}} = \prod_{p \text{ singleton}} p^{-1} \prod_{p \text{ nonsingleton}} p^{-u_p - \mathbf{1}_{\ell_p > 0}}, \quad |\mathbb{E}(\text{designated term})| \leq \mathbf{1}_{\text{lit consistency}} b_{\mathcal{L}} \mathbb{E}[R|\Delta G|]. \quad (5.10)$$

Write $\mathcal{L}$ for the lit/unlit designation. Here ℓ_p, u_p count lit and unlit occurrences, and $U = \sum_{p \text{ nonsingleton}} u_p$ is the total number of unlit nonsingleton occurrences. Lit tests must force the same residue at their departures. This is a direct application of the numbered source lemma, with its independence condition verified.

Next follow the three discarded classes. Let $\mathcal{M}_a$ be the crude factor in the proof of [O, Lemma 5.2, p. 22] and the discussion preceding Lemma 5.6 (p. 26), with padding coefficient $a^{\omega(u)}$. It retains the padding divisibilities, but drops denominators, cutoffs and added small-prime masks. Pointwise, for the non- $\mathcal{P}$ factor of any fixed numerical word,

$$0 \leq R_{\text{mask}} \leq \mathcal{M}_{16} := \prod_{i=1}^{2k} L 16^{\omega(u_i)} \mathbf{1}_{u_i | n_i}, \quad (5.11)$$

$$\mathcal{M}_{16} = 4^{\sum_i \omega(u_i)} \mathcal{M}_4 \leq \exp(200k(\log 4) \log L) \mathcal{M}_4 = \exp(O(L \log L)) \mathcal{M}_4.$$

Here $w \geq 1$ and $\omega(u_i) \leq 100 \log L$. The remaining lit-consistency and reciprocal factors are identical. Thus replacing 4 by 16 introduces only this additional factor in the positive majorants of [O, Lemmas 5.6, 6.2, 6.5, pp. 26, 28–29, 31–33]. Their crude enumeration still costs $\exp(O(L \log^2 L))$, with all padding divisibilities retained. The added small-prime draws integrate to at most one after their masks are dropped, rather than requiring a sum over their configurations. For $U > L^{1/50}$ unlit occurrences, at least $U/2$ extra reciprocal powers remain. They give

$$\exp(O(L \log^2 L)) H_0^{-U/2} \leq \exp(O(L \log^2 L) - \frac{1}{2} L^{203/200}), \quad (5.12)$$

as in [O, Lemma 5.6, p. 26].

For high rank use [O, Lemma 6.1, pp. 27–28] as stated. In one column let z_i be the abstract label at step i , with numerical prime value p_{z_i} , and let $(\mathbf{e}_z)$ be the coordinate basis indexed by its abstract labels. Put

$$t_i = D_i/p_{z_i}, \quad v(i) = \sum_{a < i} t_a \mathbf{e}_{z_a}.$$

For two lit occurrences i, i' of a label z , its control/difference pair is $(\mathbf{e}_z, v(i) - v(i'))$. Joint independence of a family means independence of all its displayed vectors over $\mathbb{R}$. The coefficients t_i are independent of the numerical prime values in that column. A jointly independent family of $r_0 = \lfloor L^{1/50} \rfloor$ such pairs has an invertible minor over $\mathbb{R}$ after deleting its control columns. The independent reciprocal prime law has largest atom at most $(V_i H_0)^{-1}$. The differences in the minor are bounded by $2khe^{103L+1} = e^{O(L)}$, since eligible paddings still obey $u \leq e^{100L+1}$. These are precisely the atom, independence and size hypotheses of Lemma 6.1; no invertibility modulo a control prime is required. Its square-root probability saving is $\exp(O(r_0 \log L)) H_0^{-r_0/2}$. Multiplying the crude enumeration gives $\exp(-cL^{203/200})$. This redoes the application in [O, Lemma 6.2, pp. 28–29] for the new fixed padding coefficient.

For $S_1 > L^{1/4}$ singletons, [O, Lemma 6.3, p. 30] applies: $\Delta G \neq 0$ forces $T = \lfloor L^{1/12} \rfloor$ positive witnesses in a single hybrid, with distinct private marked singletons, and recorded total length at most $2k + \ell_0 T \leq 4L$. Its proof uses only the witness supports and the mixed difference, unchanged here. [O, Lemma 6.4, pp. 31–32] supplies either an active private label with a unit coefficient in an internal congruence, or an active label with a nonzero private-label prefix contribution. The prohibited-word and minimality definitions are exactly Definition 2.8, so its statement applies as given. To repeat [O, Lemma 6.5, pp. 32–33], order witnesses by attachment. If at least $T/8$ have an active label absent from all earlier witnesses, use their internal congruences and eliminate in reverse order; if absent from all later witnesses, reverse that order. Otherwise more than $3T/4$ witnesses have all their active labels in both an earlier and a later witness. For private labels whose main occurrence has not passed their attachment, compare with an earlier witness; the main segment omits that label, and privacy removes it from the partner witness. For labels already passed, compare with a later witness and reverse the attachment order. One group supplies at least $\lfloor T/8 \rfloor$ relations. Each relation is independent of all later selected variables and its selected coefficient is a unit. Use (2.21) successively. The residue tests for new witness primes are unchanged in every hybrid, so no hybrid-choice cost is incurred. All records still cost $\exp(O(L \log^2 L))$. The resulting bound is

$$\exp(O(L \log^2 L)) \left(\frac{1+L}{H_0} \right)^{\lfloor T/8 \rfloor} \leq e^{-L^{21/20}}, \quad T \log H_0 \asymp L^{647/600}. \quad (5.13)$$

Thus all three exceptional classes have total $\exp(-cL^{1+\epsilon_1})$ for some absolute $\epsilon_1 > 0$. The added small-prime draws were dropped in these positive bounds, with no sum over their configurations.

After the three classes are discarded, [O, Lemmas 7.1–7.2, pp. 34–37] apply to the remaining words as stated. Perfect positions are lit nonsingletons; their blocks have length at most ℓ_0 , have no prohibited witness, and have unequal consecutive tuples within each half. Their interval-shaped labels and nonzero subinterval displacements follow from Lemma 7.1. The bounds $S_1 \leq L^{1/4}$, $U \leq L^{1/50}$ and rank below r_0 are exactly the hypotheses of Lemma 7.2. Its forest universe has $e^{O(Jk)}$ combined patterns uniformly over all numerical labels and padding coefficients. Added masks only select a subset of this universe.

At this point repeat [O, Lemma 7.3 and (7.5), p. 37], changing its row constant. Fix the non- $\mathcal{P}$

environment, numerical tuples d_i and orientations $\epsilon_i \in \{\pm 1\}$. The coefficient at a departure n is

$$c_i(n, u) = \frac{La(u)}{w(n)} \mathbf{1}_{u|n} \mathbf{1}_{\text{bin}} R_z(n + \epsilon_i h d_i u) \cdot \mathbf{1}_{\rho_{j,z}(n, d_i) \leq KW_1/L}, \quad a(u) = 16^{\omega(u)}. \quad (5.14)$$

The retained oriented row satisfies

$$\sum_u \frac{L16^{\omega(u)}}{w(n)} \mathbf{1}_{u|n} \mathbf{1}_{\text{bin}} R_z(n \pm hdu) \mathbf{1}_{\rho_{j,z}(n, d) \leq KW_1/L} \leq KW_1 = \mathfrak{b}. \quad (5.15)$$

On a closed word the product of the endpoint rough indicators equals the product of the arrival indicators, with visit multiplicities. This rewriting has already been made before closure is dropped. Retain the arrival rough factor in every c_i while dropping block and the other endpoint restrictions. Set $F_{m+1} = 1$ and recursively define

$$F_i(n) = \sum_u c_i(n, u) F_{i+1}(n + \epsilon_i h d_i u) \leq KW_1 \|F_{i+1}\|_\infty, \quad (5.16)$$

$$F_i(n) \leq \mathfrak{b}^{m-i+1}, \quad F_1(n) \leq (KW_1)^m, \quad m = 2k.$$

The eligible-degree condition may be retained or dropped in this positive row bound; either choice gives the same upper bound. Backward induction therefore gives padding mass at most $(KW_1)^{2k}$. This works for every $\mathfrak{b} \geq 0$, including $\mathfrak{b} < 1$, and survives singleton overwrites.

There are at most $Jk + S_1/2$ distinct centre labels. As in [O, (7.6)–(7.7), p. 38], their remaining reciprocal sum and the forest count give

$$\mathfrak{b}^{2k} e^{C_4 Jk} (2W)^{Jk + L^{1/4}/2}. \quad (5.17)$$

Signs, designations, 2^{S_1} and the external e^{108L} are absorbed in $e^{C_4 Jk}$. For fixed W , $(2W)^{L^{1/4}/2} \leq e^k$ eventually. The logarithm of the resulting main bound is at least $-O_W(L \log L)$ because $W_1 \gg L^{-1/6}$. Both the exceptional classes and finite-comparison error e^{-L^9} are smaller; one absolute enlargement of C_2 , before choosing W , absorbs them. This proves (5.6).

Finally redo [O, Lemma 8.2 and Proposition 8.3, pp. 40–41]. The absolute centred factor a_d is constant along a d -edge. For an outgoing increasing edge from n to $n + hdu$, the weighted row reads $R_z(n + hdu)$ and is bounded by $L\rho_{j,z}^+(n, d)$. For an incoming edge, read it from its larger endpoint n to $n - hdu$; $u | n$ is equivalent to divisibility at the lower endpoint, and the row is bounded by $L\rho_{j,z}^-(n, d)$. Since $\rho_{j,z} = \max(\rho_{j,z}^+, \rho_{j,z}^-)$, each orientation contributes at most $KW_1 a_d(n)$. Weighted Schur with weight $\sqrt{w}$ and (5.15) gives

$$\|A_d\| \leq 2\mathfrak{b}, \quad \|A_d v\|^2 \leq 4\mathfrak{b}^2 \sum_n a_d(n)^2 |v_n|^2, \quad \sum_d \|A_d v\|^2 \leq 4\mathfrak{b}^2 (8W)^J \|v\|^2 \quad (v = Ev). \quad (5.18)$$

Indeed $\sum_d a_d(n)^2 \leq \prod_i (\omega_{\mathcal{P}_i}(n) + V_i) \leq (8W)^J$ on E , by the arithmetic–geometric mean inequality. No commutation with E is used. Since $\rho(H)^{2k} \leq \|H^k\|_{\text{HS}}^2$, Markov gives $\rho(H) \leq e\mathfrak{b}(C_2\sqrt{W})^J$ outside e^{-2k} of the origins. Input 5.1 now gives (5.7) with, for example, $C \geq 3 \max(2, 2\sqrt{8}, eC_2)$. All hypotheses have nonnegative row constants; none requires $\mathfrak{b} \geq 1$. $\square$

6 Centering and the hard rough theorem

Proposition 6.1 (Polynomial on rough shifts). *Fix $a < 199/200$, and suppose every element of $\mathcal{Z} \subset [D, 2D)$ avoids the primes up to e^{L^a} , where $H_0/2 \leq D \leq e^{2L}$. For arbitrary $|c_b| = 1$, put*

$$\mathcal{B}(\theta) = \sum_{b \in \mathcal{Z}} \frac{c_b e(hb\theta)}{b}. \quad \|\mathcal{B}\|_\infty \ll L^{-a}, \quad \int_0^1 |\mathcal{B}(\theta)|^4 d\theta \ll D^{-1} L^{-4a}. \quad (6.1)$$

Proof. We repeat [20, Lemma 4.2, pp. 13–14] with the changed cutoff e^{L^a} . For the sup norm, count one rough variable in $[D, 2D)$. Orthogonality in the fourth moment imposes $b_1 + b_2 = b_3 + b_4$; after discarding the unimodular coefficients and using weights at most D^{-4} , count the four forms $x_1, x_2, x_3, x_1 + x_2 - x_3$ in a three-dimensional box of side D . The last form is positive there. Their bad-prime union densities are $1/p$ and $4/p + O(p^{-2})$, respectively. Every pair of the four defining hyperplanes is independent, including at 2, and the vector $(1, 1, 1)$ avoids all four; their union density is strictly below one.

Use even inclusion–exclusion of order $r_0 = 2\lceil 500 \log L \rceil$. For each selected prime set of product q , CRT in the interval or box gives normalized error $O(q/D)$. There are at most $(r_0 + 1)(1 + \pi(e^{L^a}))^{r_0}$ terms and $q \leq e^{r_0 L^a}$, so the total error is

$$D^{-1} \exp(O(L^a \log L)). \quad (6.2)$$

It is smaller than every fixed power of L^{-1} because $a < 199/200$. The independent truncation remainder is at most $(\sum_{p \leq e^{L^a}} \nu(p))^{r_0+1} / (r_0 + 1)! \ll L^{-100}$, since that sum is at most $(4 + o(1)) \log L$. The exact products are $O(L^{-a})$ and $O(L^{-4a})$. Thus the two counts are $O(DL^{-a})$ and $O(D^3 L^{-4a})$, giving (6.1). $\square$

Proposition 6.2 (Density in the testing and overlap step). *Use the rough matrix (5.2), an origin interval of length at least $e^{L^{625}/2}$, and the vector $\sqrt{w} \lambda R_z$, truncated to its low- $\mathcal{Q}$ state and then projected by E . Its averaged squared norm is $O_h(MSW_1)$. For a bin the retained centred sum has bound*

$$|C_j^\circ| \ll_h \frac{SW_1 KW_1 (C\sqrt{W})^J}{L} + e^{-L^{0.9}}. \quad (6.3)$$

After summing bins and dividing by S_0 , the spectral contribution is

$$O_h \left(W_1^2 \frac{K}{\eta} (C/\sqrt{W})^J \right) + e^{-L^{0.8}}. \quad (6.4)$$

Proof. Repeat [20, Proposition 8.4, pp. 41–42] with all density factors retained. The squared vector is the truncated weight wR_z . Its product-law mean is at most SW_1 , since the small-prime coordinates are disjoint from $\mathcal{Q}$. Proposition 3.3 transfers this mean to each of the M block positions, giving the norm bound. Test (5.7) on the nonexceptional blocks. On every block the weighted absolute form is at most a fixed power of L times M : use $w \leq 17^{18 \log L}$ and $\sum_d a_d \leq (8W)^J$ on the exterior projection. Thus the fraction e^{-2k} of exceptional blocks costs exponentially little even after division by $SW_1^2 V_*$.

For completeness, the overlapping-block identity of [O, (8.12)–(8.13), p. 42] is unchanged. If an increasing edge has displacement $q' = hdu$, the number of blocks among $t = 0, \dots, N-1$, $N = \lfloor T_j \rfloor$, containing it is

$$m_N(n, q') = [\min(N-1, n-1) - \max(0, n+q' - M) + 1]_+. \quad (6.5)$$

For $M \leq n \leq N$ this is $M - q'$. The boundary involves at most $2M$ sites, and the remaining discrepancy is bounded by the deterministic row bound times

$$O_h \left(\frac{he^{100L+1}}{M} + \frac{M}{T_j} + \frac{1}{T_j} \right). \quad (6.6)$$

The row bound divided by M is a fixed power of L , $M = e^{103L} + O(1)$, and $T_j \geq e^{L^{625}}$. This includes floor errors and edges passing the prefix endpoint. Dividing the quadratic form by $2LM$ gives (6.3). There are $O(L/\eta)$ bins and $S_0 \geq SV_*/2$, $V_* \geq W^J$; these give (6.4). The argument did not use a signed-norm monotonicity principle. $\square$

Proof of Theorem 1.1. Choose a permissible absolute $C \geq 4$ in Proposition 5.2 before W , and set

$$\begin{aligned} A &= 625, \quad L = (\log X)^{1/A}, \quad \beta = 1/6, \\ W &= 4eC^2, \quad \alpha = 1/(1200W), \quad J = \lfloor \alpha \log L \rfloor, \\ \varepsilon &= \alpha/100, \quad \eta = L^{-\varepsilon}, \quad K = L^{\alpha \log 2 + \varepsilon}. \end{aligned} \quad (6.7)$$

All labels exceed z . The function $f_z = \lambda R_z$ is completely multiplicative and, for a graph label $q' = du$,

$$f_z(q'm)f_z(q'(m+h)) = f_z(m)f_z(m+h). \quad (6.8)$$

The square $f_z(q')^2 = 1$ proves this without a coprimality assumption between q' and m . The full-divisibility part of the centred sum is exactly

$$\sum_j \sum_{(d,u) \in \mathcal{S}_j} \frac{16^{\omega(u)}}{du} \frac{1}{T_j/(du)} \sum_{m \leq T_j/(du)} f_z(m)f_z(m+h). \quad (6.9)$$

Its cutoff is in $(Xe^{-\eta}, X]$. The positive upper sieve bounds the missing interval by $O_h(\eta X V_h(z))$ and the full rough count by $O_h(X V_h(z))$. If an actual tail is shorter, enlarge it to the complete interval $(Xe^{-\eta}, X]$. The required sieve moduli are $X^{o(1)}$, so this bound is uniform in z . Thus (6.9) differs from $S_0 X^{-1} \sum_{n \leq X} f_z(n)f_z(n+h)$ by $O_h(S_0 \eta V_h(z)) + e^{-L}$.

The padding deletion is Proposition 4.1. For the degree deletions repeat [20, Lemma 4.6, pp. 19–20], verifying the changed tilted law. Under w/S , the probabilities are $17/(p+16)$, and exponential Markov with parameter $1/20$ gives

$$\mathbb{P}^{\text{tilt}}(\omega_{\mathcal{Q}} > 18 \log L) \ll_h L^{-18/20+17(e^{1/20}-1)} \ll_h L^{-1/40}. \quad (6.10)$$

The centre tilt by a_d^+ forces at most J primes and leaves the others independent, of total mean at most $2WJ$. Its tail is at most $\exp(-6WJ + J + 2(e-1)WJ) \leq e^{-2WJ}$. In each fixed numerical label the small-prime pair has density $V_h(z)$; only these coordinates are independent of the centre and padding degree events. This retains the pair density in both deletions. For rare sites, the second moment of $w \prod_i (\omega_{\mathcal{P}_i} + V_i)$ is a fixed power of L : its $\mathcal{Q}$ factor is $\prod_{p \in \mathcal{Q}} (1 + 288/p)$, and the centre factors have moments bounded by fixed multiples of $V_i^2 + V_i$. Cauchy–Schwarz with (2.20), followed by the $O(L/\eta)$ bins, gives exponentially small error. All these are positive finite-residue expansions covered by Proposition 3.3; the padding threshold is encoded only after the low-degree cutoff. Together with Proposition 6.2, the non-centering errors, divided by $V_h(z)$, are

$$C_h \left\{ \eta + \frac{2^J(1 + \log L)^6}{K} + 2^J(L^{-1/40} + e^{-2WJ}) + \frac{K}{\eta}(C/\sqrt{W})^J \right\} + e^{-L^{0.8}}. \quad (6.11)$$

It remains to redo [20, Lemmas 4.1–4.2, 4.4, pp. 13–16] with f_z , the new range, and the densities. Use Proposition 6.1 with $a = 9949/10000$, and set $\Gamma = 89/100$, $s = 17/100$. In a partial centred term, a nonempty subset I of the J supplies is replaced by $-1/p$. Writing $b = \prod_{i \in I} p_i$, $d' = d/b$, and $n = ud'v$ extracts a reciprocal mass at most SV_* and leaves

$$\frac{1}{Y} \sum_{v \leq Y} \sum_{b \in \mathcal{Z}} \frac{c_b}{b} f_z(v) f_z(v + hb), \quad Y = T_j/(ud'). \quad (6.12)$$

The allowed b lie in at most two dyadic intervals, $H_0/2 \leq D \leq e^{2L}$, and $Y \geq e^{L^{625}/2}$. The coefficients are unimodular. This is [O, (4.8), p. 16] with the exact identity (6.8) in place of its original Liouville identity.

Replacing $\lambda(p)$ by zero at $p \leq z$ reduces (2.12) by at most $\sum_{p \leq z} 1/p \leq \beta \log L + O(1)$. Input 2.5 therefore gives $M(f_z; Y, Q) \geq (625/5) \log L$ eventually. Here $10 \leq D \leq Y$ and Q has exactly the prescribed cap; the three errors in (2.13) are bounded by

$$L^{-25/4}, \quad L^{-199/200} \log L, \quad L^{-25/28}, \quad \frac{1}{Y} \sum_{v \leq Y} \left| \sum_{m=1}^D f_z(v+m) e(\theta m) \right| \ll DL^{-\Gamma}. \quad (6.13)$$

The last assertion follows from unit intervals of origins, as in [O, (4.6), p. 15]; endpoint cost $O(D/Y)$ is exponentially small.

Average (6.12) over translations $1 \leq m \leq D$; the endpoint cost is $O_h(D/Y)$. Fourier orthogonality gives $(YD)^{-1} \sum_{v \leq Y} \int \mathcal{B} F_v G_v$, with $F_v = \sum_{m=1}^D f_z(v+m) e(\theta m)$ and $G_v = \sum_{a=1}^{(2h+1)D} f_z(v+a) e(-\theta a)$. For any such window, the elementary upper sieve gives rough count $O_h(DW_1)$. For clarity, even inclusion–exclusion of order $O(\log L)$ has total CRT endpoint error $D^{-1} \exp(O(L^{1/6} \log L))$, and its independent remainder can be any fixed negative power of L . This verifies the uniformity even for the smallest D . Thus $\|F_v\|_2 \|G_v\|_2 \ll_h DW_1$ and $|G_v| \ll_h DW_1$. On $|\mathcal{B}| \leq L^{-1-s}$ use Parseval; on the complement its measure is at most $L^{4+4s} \int |\mathcal{B}|^4 \ll D^{-1} L^{4+4s-4a}$. Use also $\|\mathcal{B}\|_\infty \ll L^{-a}$ and (6.13). The two partial contributions are

$$O_h(W_1 L^{-1-s}), \quad O_h(W_1 L^{4+4s-5a-\Gamma}). \quad (6.14)$$

After at most 2^J subsets, $O(L/\eta)$ bins and division by $V_h \asymp_h W_1^2$, their exponents are

$$\beta + \varepsilon + \alpha \log 2 - s, \quad \beta + \varepsilon + \alpha \log 2 + 5(1-a) + 4s - \Gamma. \quad (6.15)$$

Without the ε, α terms the margins are $-1/300$ and $-107/6000$. For $W \geq 10$ both are less than $-\varepsilon$.

In (6.11), the main deletion is $L^{-\varepsilon}(1 + \log L)^6$ and, since $C/\sqrt{W} = 1/(2\sqrt{e})$, the spectral exponent is $-\alpha/2 + 2\varepsilon$. The remaining degree terms have strict negative margins. Absorbing logarithmic factors gives

$$\left| \sum_{n \leq X} \lambda(n) \lambda(n+h) R_z(n) R_z(n+h) \right| \ll_h X V_h(z) L^{-\varepsilon/2}. \quad (6.16)$$

Lemma 2.4 supplies the unsigned and both single-sign terms with arbitrary-logarithmic absolute errors. Choose their saving larger than $2/3750 + \gamma$ and use the exact sign expansion

$$\mathbf{1}_{\lambda(n)=s, \lambda(n+h)=t} = \frac{1}{4} (1 + s\lambda(n))(1 + t\lambda(n+h)). \quad (6.17)$$

Equation (1.2) follows, for example with $\gamma = \varepsilon/(4 \cdot 625)$. The cutoffs satisfy $\log z \leq L^{1/6}$, exactly the stated range. All graph constants precede W and X ; choose the final threshold after h and those constants are fixed. Origin exceptions were internal averages, not exclusions of values of X . $\square$

7 Balanced soft small-prime weights

Lemma 7.1 (Prime-square masks and densities). *For $0 \leq r_1, r_2 \leq 1$ the function*

$$A_z(n) = S_z(n)S_z(n+h)r_1^{\omega_z(n)}r_2^{\omega_z(n+h)} \quad (7.1)$$

is an average of negative prime-divisibility CNF indicators, with total averaging mass one, together with the always retained negative prime-square tests. Its unsigned mean is compared by Lemma 3.1 in the prime-square model. When $r_1 = r_2 = r$, its pair mean is exactly (1.6). Its single and squared means are

$$W_r = \prod_{p \leq z} (1 - 1/p + r(p-1)/p^2), \quad W_{r^2} = \prod_{p \leq z} (1 - 1/p + r^2(p-1)/p^2). \quad (7.2)$$

Uniformly for $0 \leq r \leq 1$, $\delta = 1 - r$, and $z \geq 2$,

$$\log W_r = -\delta \log \log(2z) + O(1), \quad (7.3)$$

$$\log W_{r^2} = -(2\delta - \delta^2) \log \log(2z) + O(1), \quad (7.4)$$

$$\log V_{r,h}^{\text{sf}}(z) = -2\delta \log \log(2z) + O_h(1). \quad (7.5)$$

In particular

$$\begin{aligned} W_r &\asymp v^{-\delta}, \quad W_{r^2} \asymp v^{-(2\delta-\delta^2)}, \quad V_{r,h}^{\text{sf}} \asymp_h v^{-2\delta} \asymp_h W_r^2, \\ W_{r^2}/V_{r,h}^{\text{sf}} &\ll_h v^{\delta^2}, \quad W_r/V_{r,h}^{\text{sf}} \ll_h v^\delta, \quad v = \log(2z). \end{aligned} \quad (7.6)$$

Proof. Independently retain $p \nmid n$ with probability $1 - r_1$, and $p \nmid n + h$ with probability $1 - r_2$. Always retain $p^2 \nmid n, n+h$. On a coordinate modulo p^2 , $p \nmid n$ excludes the p residues $0, p, \dots, (p-1)p$, and a negative square test excludes one residue. Each realization is thus a conjunction of at most $2 \sum_{p \leq z} (p+1)$ negative equalities. For $z \leq e^{L^b}$, $b < 1/2$, the clause count is less than e^{L^3} and the moduli are at most e^L . Averaging Lemma 3.1 costs no scalar coefficient mass. The same argument is valid with $b < 1$ whenever the coordinate-size condition holds separately.

List the residues modulo p^2 . Outside the forbidden roots modulo p the pair weight is one. A root at one endpoint has $p-1$ admissible lifts of weight r . A common root has $p-1$ or $p-2$ admissible lifts of weight r^2 , according as $p^2 \mid h$ or not. This gives (1.6); at $p=2$, $h \equiv 2 \pmod{4}$, the factor is exactly $1/2$. The analogous single-site count gives (7.2).

For the full range $0 \leq r \leq 1$ the single local factors are

$$q_r(p) = 1 - \delta/p - r/p^2, \quad q_{r^2}(p) = 1 - (2\delta - \delta^2)/p - r^2/p^2, \quad v_{r,h}(p) = 1 - 2\delta/p - 2r/p^2 \quad (p \nmid h). \quad (7.7)$$

The single factors are at least $1 - 1/p$. For $p \geq 5$, the factors are at least $4/5, 4/5, 3/5$, respectively. Taylor expansion of their logarithms therefore has a uniform $O(p^{-2})$ error. The factors at $p=2, 3$ and at $p \mid h$ are finitely many positive exceptions with uniform positive lower bounds: the common-root factors are at least $1 - 1/p$, and the generic factor at 3 is at least $1/3$. Mertens' harmonic sum and $\sum_p p^{-2} < \infty$ give (7.3)–(7.5); exponentiation gives (7.6). Only these product comparisons are uniform in r . Theorem 1.3 still fixes r before X . $\square$

Proposition 7.2 (Classical single-sign orthogonality). *For each fixed $0 < \rho_* < 1$ and every $B > 0$, uniformly for $2 \leq z \leq \exp((\log X)^{\rho_*})$ and $r_1, r_2 \in [0, 1]$,*

$$\sum_{n \leq X} \lambda(n) A_z(n), \quad \sum_{n \leq X} \lambda(n+h) A_z(n) \ll_{h, \rho_*, B} X (\log X)^{-B}. \quad (7.8)$$

Proof. Average the negative prime tests in Lemma 7.1. It suffices to treat uniformly the deterministic mask $M(n) = \prod_{p \in \mathcal{T}_1} \mathbf{1}_{p \nmid n} \prod_{p \in \mathcal{T}_2} \mathbf{1}_{p \nmid n+h}$, $\mathcal{T}_i \subset \{p \leq z\}$, with the two squarefree masks still present. Put $T = X^{1/100}$ and

$$S_{z,T}(n) = \sum_{\substack{b \leq T, \\ b^2 \mid n}} \mu(b). \quad (7.9)$$

The full identity is the same sum without $b \leq T$. Thus $|S_z - S_{z,T}| \leq \sum_{b > T, b^2 \mid n} 1$ and $|S_{z,T}(n)| \leq \tau(n)$. Replacing both factors has L^1 error at most

$$\sum_{n \leq X} \sum_{b > T, b^2 \mid n} 1 + \sum_{n \leq X} \tau(n) \sum_{c > T, c^2 \mid n+h} 1 \ll_{\xi} X/T + X^{1+\xi}/T = O_h(X^{199/200}), \quad \xi = 1/200. \quad (7.10)$$

Use the complete shifted prefix through $X+h$ and $\tau(n) \ll_{\xi} n^{\xi}$; no additional floor sum is charged.

For fixed $b, c \leq T$, the two square divisibilities are inconsistent or give one class modulo $q_0 = [b^2, c^2] \leq T^4$. Tests at primes dividing q_0 are automatic or make the mask zero. Remove them. At each remaining prime at most two roots are excluded, and CRT works with d coprime to q_0 . Input 2.2, at level $D = X^{1/8}$, gives the L^1 upper-sieve approximation error on this progression

$$O\left(\frac{X}{q_0} E(s) + D \log(2D)\right), \quad s \geq \frac{1}{8}(\log X)^{1-\rho_*}. \quad (7.11)$$

Its local product is at most one; its dimension condition is uniformly bounded by the two-root sieve. The modulus and prime masks introduce no new constants. Summing this error costs $O(XE(s) + T^2 D \log(2D))$, since

$$\sum_{b, c \geq 1} [b^2, c^2]^{-1} \leq \zeta(2)^3, \quad T^2 D = X^{29/200}. \quad (7.12)$$

For the inequality write $b = gu, c = gv$, $(u, v) = 1$ and drop that coprimality. Inserting a Liouville sign cannot increase this L^1 error.

All remaining sums have modulus $q = q_0 d \leq X^{33/200}$. For fixed q there are at most $\tau(q)^3$ choices of b, c, d and at most $\tau(q)$ root classes per choice. The absolute remainder is bounded by

$$\sum_{q \leq X^{33/200}} \tau(q)^4 E_q, \quad E_q = \max_{a \bmod q} \max_{y \leq 2X} \left| \sum_{\substack{n \leq y \\ n \equiv a \pmod{q}}} \lambda(n) \right|. \quad (7.13)$$

Lemma 2.3 gives $\sum E_q \ll_A X \log^{-A} X$ and $E_q \leq 2X/q + 1$. Since $\tau(q)^8 \leq \tau_{256}(q)$, the divisor-variable bounds give

$$\sum_{q \leq X^{33/200}} \tau(q)^8 E_q \ll X \log^{256}(2X), \quad \sum \tau(q)^4 E_q \ll_A X \log^{128-A/2} X. \quad (7.14)$$

Choose A sufficiently large. The translated sign is a difference of two prefixes in another residue class. This proves (7.8) using only the classical FL and BV inputs. $\square$

Lemma 7.3 (Soft padding recursion). *Fix $0 < r < 1$, use padding 16, and let $b = b_z$ and b^+ be its cap in (7.19). In a fixed non- $\mathcal{P}$ residue environment put*

$$\rho_j^{\pm}(n, d) = \frac{1}{w(n)} \sum_{\substack{u \mid n, u \in \mathcal{Q}^{\text{sf}} \\ j\eta \leq \log(du) < (j+1)\eta}} a(u) b^{\pm}(n \pm hdu), \quad \rho_j = \max(\rho_j^+, \rho_j^-). \quad (7.15)$$

For fixed d_i and $\epsilon_i \in \{\pm 1\}$ define

$$c_i^0(n, u) = \frac{La(u)}{w(n)} \mathbf{1}_{u|n} \mathbf{1}_{\text{bin}} \mathbf{1}_{\rho_j(n, d_i) \leq KW_r/L}. \quad (7.16)$$

With $F_{m+1} = 1$, the recursion obeys

$$\begin{aligned} F_i(n) &= \sum_u c_i^0(n, u) b(n + \epsilon_i h d_i u) F_{i+1}(n + \epsilon_i h d_i u) \\ &\leq KW_r \|F_{i+1}\|_\infty, \quad F_1(n) \leq (KW_r)^m. \end{aligned} \quad (7.17)$$

On a closed word the balanced edge factors satisfy

$$\prod_{i=1}^m \sqrt{b(n_i) b(n_{i+1})} = \prod_{i=1}^m b(n_{i+1}) = \prod_{i=1}^m b(n_i), \quad n_{m+1} = n_1, \quad (7.18)$$

including repeated visits and zero weights.

Proof. Since $0 \leq b \leq b^+$, the row sum of $c_i^0(n, u) b(n + \epsilon_i h d_i u)$ is at most $L \rho_j^{\epsilon_i}(n, d_i) \mathbf{1}_{\rho_j \leq KW_r/L} \leq KW_r$. Here $\rho_j^{\epsilon_i}$ means ρ_j^+ for $\epsilon_i = 1$ and ρ_j^- for $\epsilon_i = -1$. This is the backward induction of [20, Lemma 7.3, p. 37] with a nonnegative row constant KW_r . It is uniform in the starting site and does not require $KW_r \geq 1$. For (7.18), every visited factor occurs in the two square roots at its incident edges, counted with visit multiplicity. Nonnegativity gives the identity, including when a factor is zero. Make it before dropping closure in the positive path enlargement; then (7.17) at $m = 2k$ gives $(KW_r)^{2k}$. Adding the eligible padding-degree condition only decreases these sums. $\square$

Proposition 7.4 (Balanced operator and capped cutoff). *Fix $0 < r < 1$ and retain (6.7). Let $b = b_z$. Use the symmetric edge matrix (7.20) below and test it with $\sqrt{w(n)b(n)}\lambda(n)$. Define*

$$M_r = \left\lceil \frac{100 \log L}{-\log r} \right\rceil, \quad b^+(n) = \max(b(n), r^{M_r}), \quad r^{M_r} \leq L^{-100}. \quad (7.19)$$

Use $b^+(n \pm hdu)$ in the target padding probabilities (4.1), take their maximum ρ_j , and cut at $\rho_j \leq KW_r/L$, after the low- Q cutoff. The actual matrices and vectors use b . More explicitly, for $m = n + hdu$ let

$$\begin{aligned} \tau_j^b(n, d) &= \mathbf{1}_{\omega_Q(n) \leq \lfloor 18 \log L \rfloor} \mathbf{1}_{\rho_j(n, d) \leq KW_r/L} \sigma_j(n), \\ A_d^b(n, m) &= \frac{L 16^{\omega(u)}}{\sqrt{w(n)w(m)}} \mathbf{1}_{u|n} c_d(n) \tau_j^b(n, d) \tau_j^b(m, d) \sqrt{b(n)b(m)}. \end{aligned} \quad (7.20)$$

Set $A_d^b(m, n) = A_d^b(n, m)$ and other entries to zero. In the following proof write $A_d = A_d^b$. The associated centred sums use $f = g = \lambda b$, $a = 16$, $\epsilon = 1$ and $\tau_j = \tau_j^b$. They satisfy the finite comparison of Proposition 3.3, the moment and spectral bounds of Proposition 5.2 with W_1 replaced by W_r , and averaged testing norm $O_{h,r}(MSW_r)$. Explicitly, with $H_{bd,d'} = \mathbf{1}_{d \neq d'} A_{d'}^b$,

$$\mathbb{E}_* \|H_b^k\|_{\text{HS}}^2 \leq \{KW_r (C_2 \sqrt{W})^J\}^{2k}, \quad \left\| E \left(\sum_d A_d^b \right) E \right\| \leq KW_r (C \sqrt{W})^J. \quad (7.21)$$

The spectral estimate excludes at most a fraction e^{-2k} of the block origins; the finite-origin and constant conventions are those of Proposition 5.2. Their padding deletion is

$$O_{h,r} \left(W_r^2 \frac{2^J (1 + \log L)^6}{K} \right). \quad (7.22)$$

Proof. The exact quadratic pairing contains $b(n)b(m)$, because each endpoint contributes a square root in the matrix and a square root in the vector. At zero-weight vertices all entries and the vector vanish. On the remaining space weighted Schur uses the positive weight $\sqrt{wb}$ and gives

$$|A_d(n, m)| \frac{\sqrt{w(m)b(m)}}{\sqrt{w(n)b(n)}} = \frac{L16^{\omega(u)}}{w(n)} a_d(n) \mathbf{1}_{u|n} b(m) \times \text{retained cuts.} \quad (7.23)$$

Thus the oriented padding row is at most KW_r . On a closed word,

$$\prod_{i=1}^{2k} \sqrt{b(n_i)b(n_{i+1})} = \prod_{i=1}^{2k} b(n_{i+1}), \quad n_{2k+1} = n_1. \quad (7.24)$$

This includes repeated visits and zeros. The denominators have the identical closed-word identity for w . Apply it before dropping closure; no square root remains to be encoded.

For finite comparison, fix an active state of at most $18 \log L$ padding primes, hence at most $N \leq L^{18 \log 2}$ candidates. A bad weighted sum is the OR over level vectors $(k_1, \dots, k_N) \in \{0, \dots, M_r\}^N$ whose scalar sum exceeds the threshold, of the events $b^+(n_i) \geq r^{k_i}$ for all i . If $k_i = M_r$ this event is automatic; otherwise it is exactly

$$S_z(n_i) = 1 \quad \text{and} \quad \omega_z(n_i) \leq k_i. \quad (7.25)$$

Negation yields an AND of ORs of square divisibilities $p^2 \mid n_i$ or conjunctions of $k_i + 1$ distinct prime divisibilities. The square-divisibility alternative is essential at $b = 0$. On a p^2 coordinate, $p \mid n_i$ is an OR of p equalities and $p^2 \mid n_i$ one equality. Substitute the interval DNFs of Proposition 3.3, distribute the bottom $O_r(\log L)$ conjunctions and flatten ORs. The depth is three, and the logarithmic size is at most

$$N \log(M_r + 1) + O_r(L^{1/6} \log L + \log^2 L) + O(L \log L) < L^{14}. \quad (7.26)$$

The coordinate moduli are at most e^L , pairwise coprime to the disjoint graph coordinates. Use $t = \lceil L^{589} \rceil$ and the identical CRT/Walsh correction, giving aggregate error e^{-L^9} . Every residual factor b is the average in Lemma 7.1; repeated occurrences use independent auxiliary marks, which correctly produce b to that multiplicity. Their probabilities sum to one. They only add CNF conditions and introduce no scalar-mass cost or physical-site independence.

The padding experiment is unchanged from Proposition 4.1. For different eligible paddings the generic three-site factor, at $p > 3$, $p \nmid h(u_1 - u_2)$, is exactly $1 - 3\delta/p - 3r/p^2$. Comparing it with three single-site factors changes a convergent $O(p^{-2})$ product. Lemma 2.1 bounds exceptional primes as before, and gives

$$\mathbb{E}_{\text{small}}[b(n)b(n \pm hdu_1)b(n \pm hdu_2)] \ll_{h,r} W_r^3(1 + \log L)^6. \quad (7.27)$$

At equal paddings the mean of $b(n)b(n + hdu)^2$ has generic logarithmic coefficient $-(3\delta - \delta^2)/p$ and is at most $C_{h,r} W_r^3 v^{\delta^2}$. If $\rho_j^{(b)}$ denotes the uncapped target probability, then

$$\mathbb{E}_*[wb \sum_j (\rho_j^{(b)})^2] \ll_{h,r} SW_r^3 \{(1 + \log L)^6/L + v^{\delta^2} (v/L)^{24/17}\} \ll_{h,r} SW_r^3(1 + \log L)^6/L. \quad (7.28)$$

The diagonal-to-main exponent is at most $(1/6)(24/17 + 1) + 1 - 24/17 = -1/102$. As before the different-padding estimate is used on eligible bin pairs before enlargement. Since $b^+ \leq b + r^{M_r}$, the cap's added squared collision is $O_r(L^{-200} SW_r/L)$, by the unmasked collision estimate and

independence of the small-prime coordinates. Lemma 7.1 gives $W_r \gg L^{-1/6}$ for this whole fixed- r range, so the extra term is negligible. The maximum of the two orientations only costs a constant. On a bad row $wb\rho_j^{(b)} \leq Lwb\rho_j^2/(KW_r)$. Multiply the independent centre majorant $2^J/d$, sum, and obtain (7.22).

We now check the changed graph proof in the same order as Proposition 5.2. The trace expansion [O, pp. 24–25] uses (7.24) and remains a signed finite expansion. In [O, Lemma 5.5, p. 26] the integer powers of b , the capped cutoff and the padding state belong to the non- $\mathcal{P}$ factor R ; singleton overwrites leave them unchanged, and G is still only the witness factor. The three negligible classes drop b using $0 \leq b \leq 1$, only in their nonnegative majorants: [O, Lemma 5.6, p. 26] retains the $U/2$ extra unlit reciprocals; [O, Lemma 6.2, pp. 28–29] retains the high-rank lit-consistency equations; [O, Lemma 6.5, pp. 31–33] retains the witness and padding divisibilities and the ordered unit-coefficient eliminations. In each class the pointwise comparison (5.11) gives the same $\exp(O(L \log L))$ change of padding coefficient, and no b is replaced by a positive lower bound. After these three classes are discarded, the forest hypotheses [O, Lemmas 7.1–7.2, pp. 34–37] hold on the smaller support. On surviving words keep the arrival b : Lemma 7.3, with the closed identity (7.18) made before closure is dropped, gives the padding bound $(KW_r)^{2k}$. The completion [O, pp. 37–38] retains the same C_2 , since $W_r \gg L^{-1/6}$ makes the main logarithm at least $-O_W(L \log L)$, even if $KW_r < 1$. Weighted Schur [O, Lemma 8.2, p. 40] uses (7.23) on $b > 0$, on each invariant level set of a_d . The outgoing row uses ρ_j^+ and the incoming row ρ_j^- , each capped at KW_r/L . Vertices with $b = 0$ have zero rows and columns, so no division by zero occurs. Input 5.1 applies with $a = 2KW_r$, $b = 2KW_r(8W)^{J/2}$. This gives the same permissible absolute C . Finally the truncated squared vector is wb with independent mean at most SW_r . The overlap identity (6.5) and deterministic exceptional-block argument apply with the same fixed polynomial bounds. This proves all asserted testing and spectral estimates. $\square$

Proof of Theorem 1.3. By Lemma 7.1, $V_{r,h}^{\text{sf}} \asymp_{h,r} W_r^2$ throughout the fixed- r range. For every label $q' = du$, whose primes exceed z ,

$$b(q'm) = b(m), \quad (\lambda b)(q'm) = \lambda(q')(\lambda b)(m), \quad (\lambda b)(q'm)(\lambda b)(q'(m+h)) = (\lambda b)(m)(\lambda b)(m+h). \quad (7.29)$$

The small prime-square valuations are preserved even when q' and m share large primes. Thus the raw identity and normalization are unchanged, with a pair density $V_{r,h}^{\text{sf}}$. Positive window bounds for the pair and single masks follow by averaging the elementary upper sieve over the marks of Lemma 7.1, dropping square restrictions for an upper bound. The local products change by bounded positive factors, since the difference is $O(p^{-2})$ outside fixed exceptional primes. An $O(\log L)$ truncation has normalized CRT error $H^{-1}e^{O(L^{1/6} \log L)}$ for $\log H \gg L^{199/200}$, and negligible independent remainder. This proves bounds $O_{h,r}(HV_{r,h}^{\text{sf}})$, $O_r(HW_r)$ and $O_r(HW_{r^2})$, including the raw tail of length ηX .

Proposition 7.4 gives the relative spectral term

$$O_{h,r}((K/\eta)(C/\sqrt{W})^J). \quad (7.30)$$

Its padding deletion is (7.22); degree and rare-site deletions preserve the pair density by the disjoint small prime-square coordinates. Their bounds are those in (6.11). For centering $f = \lambda b$ is multiplicative and 1-bounded; its prime values change from those of λ only at $p \leq z$ and by at most one. Thus (6.13) still holds with $\Gamma = .89$. The small-frequency piece uses the squared density W_{r^2} , and the large-frequency piece the pointwise mean W_r . At both uses Lemma 7.1 supplies the full- r comparisons. The two relative exponents, with $a = .9949$, $s = .17$, $\beta = 1/6$, are

$$\beta\delta^2 + \varepsilon + \alpha \log 2 - s, \quad \beta\delta + \varepsilon + \alpha \log 2 + 5(1-a) + 4s - \Gamma. \quad (7.31)$$

They do not exceed (6.15), because $\delta, \delta^2 \leq 1$. The raw tail has cost $O_{h,r}(\eta V_{r,h}^{\text{sf}})$. The full relative budget therefore gives

$$\left| \sum_{n \leq X} \lambda(n) \lambda(n+h) b(n) b(n+h) \right| \ll_{h,r} X V_{r,h}^{\text{sf}} L^{-\varepsilon/2}. \quad (7.32)$$

The unsigned mass is its independent product plus an exponentially small finite-comparison error by Lemma 7.1; Proposition 7.2 gives arbitrary logarithmic savings for both single signs. Expanding (6.17) proves (1.5), with the same $\gamma = \varepsilon/(4 \cdot 625)$. The constants in the graph moments are absolute; circuit thresholds and positive comparisons may depend on the fixed r . No choice $r = r(X)$ is made. $\square$

8 Bounds on the number of prime factors

Proposition 8.1 (Positive moments). *For fixed even h , $z \leq X^{o(1)}$ and $1 \leq u \leq 2$,*

$$\sum_{2 \leq n \leq X} R_z(n) R_z(n+h) u^{\omega(n)} \ll_h X (\log X)^{u-1} (\log z)^{-(u+1)} + O(\sqrt{X} \log^2 X), \quad (8.1)$$

$$\sum_{n \leq X} b_z(n) b_z(n+h) u^{\omega(n)} \ll_h X (\log X)^{u-1} (\log z)^{-(u+1)\delta} + O(\sqrt{X} \log^2 X), \quad (8.2)$$

$$\delta = 1 - r.$$

The same estimates hold at the other endpoint. The soft upper bound applies to every fixed $0 < r < 1$.

Proof. Apply the Nair–Tenenbaum theorem [12, Theorem 1, pp. 1–2] on each dyadic interval $[Y, 2Y]$ with $Y \geq \sqrt{X}$, to $Q_1(t) = t$, $Q_2(t) = t + h$ and the nonnegative weights

$$F(a, b) = u^{\omega(a)} R_z(a) R_z(b), \quad F(a, b) = u^{\omega(a)} r^{\omega_z(a) + \omega_z(b)}. \quad (8.3)$$

In the second bound the squarefree masks have been dropped. The polynomials are primitive, irreducible and coprime; the product has degree two, discriminant $h^2 \neq 0$, and no fixed prime divisor since h is even. For the function class in that theorem, coprime-product growth is bounded by $2^{\omega(a)} \leq \tau(a)$, hence both a fixed A^Ω bound and, for every fixed $\xi > 0$, a uniform $B_\xi n^\xi$ bound. Choose its interval exponent and coefficient exponent to be $1/2$, and then a permissible ξ . The intervals have length $Y > Y^{1/2}$, and the coefficient threshold holds for large Y with fixed h . These verify the hypotheses uniformly in z, u (and in the bounded prime values of r).

Both forms have one root modulo every integer. In the positive divisor sum on the right side of the cited theorem, drop the product cutoff and enlarge to Euler factors over primes at most $2Y$. For the hard weight, its local logarithms, including the theorem’s sieve product, are $(u-1)/p + O(p^{-2})$ for $p > z$, and $-2/p + O_h(p^{-2})$ for $p \leq z$. The fixed divisors of h give bounded factors. This yields $C_h Y (\log Y)^{u-1} (\log z)^{-(u+1)}$. For the soft weight the prime values are ur, r below z and $u, 1$ above z , giving logarithm

$$(u-1) \log \log X - (u+1) \delta \log \log z + O_h(1). \quad (8.4)$$

Higher prime-power errors are uniformly summable. Geometric dyadic summation proves the stated main terms. Below $\sqrt{X}$, $\sum 2^{\omega(n)} \leq \sum \tau(n) \ll \sqrt{X} \log X$, within the remainder. Swapping the two functions proves the other endpoint. No signed or lower estimate is taken from this theorem. $\square$

Lemma 8.2 (A useful proportional tail). *If a positive pair weight has mass of order $X \log^{-2\kappa} X$ and one-endpoint moments bounded by $CX \log^{-2\kappa+(1-\kappa)(u-1)} X + O(\sqrt{X} \log^2 X)$, $1 \leq u \leq 2$, then, for $0 < \kappa < 1/2$, the mass with $\omega(n) > (1 - \kappa/2) \log \log X$ is $O(X \log^{-2\kappa-\kappa^2/32} X) + O(\sqrt{X} \log^2 X)$.*

Proof. Take $\tau = 1 - \kappa/2$, $u = 1 + \kappa/(4(1 - \kappa))$, and $a = u - 1$. Markov's inequality gives relative exponent $E = (1 - \kappa)(u - 1) - \tau \log u$. Since $\log(1 + a) \geq a - a^2/2$,

$$E \leq -\frac{a\kappa}{2} + \frac{\tau a^2}{2} = \frac{\kappa^2(-3 + (7/2)\kappa)}{32(1 - \kappa)^2} \leq -\kappa^2/32. \quad (8.5)$$

The factor $u^{-\tau \log \log X}$ multiplying the remainder is at most one. $\square$

Proof of Theorem 1.2 and Remark 1.5. Put $\rho = 1/3750$, $z = \exp((\log X)^\rho)$. For $\kappa = \rho$ use the hard weight of Theorem 1.1 and (8.1). For $0 < \kappa < \rho$ fix $r = 1 - \kappa/\rho$ and use the weight of Theorem 1.3 and (8.2). Lemma 7.1 gives $V_{r,h}^{\text{sf}}(z) \asymp_h W_r^2 \asymp_h (\log X)^{-2\kappa}$ for this entire fixed- r range, also when $\kappa > \rho/2$ and hence $r < 1/2$. In either case each sign pattern has weighted mass comparable to $X \log^{-2\kappa} X$, and

$$\sum_{n \leq X} \text{pair weight } u^{\omega(n)} \leq CX(\log X)^{-2\kappa+(1-\kappa)(u-1)} + O(\sqrt{X} \log^2 X). \quad (8.6)$$

Here $\delta\rho = \kappa$ in the soft case. Write $m = 1 - \kappa$, $\tau = m + \epsilon < 1$, $a = \epsilon/\tau \in (0, 1)$ and $u = 1 + a$. The relative Markov exponent for $\omega(n) > \tau \log \log X$ is

$$ma - \tau \log(1 + a) \leq -\epsilon a + \tau a^2/2 = -\epsilon^2/(2\tau) < 0. \quad (8.7)$$

The other endpoint has the same bound. Both unsigned tails are negligible relative to each sign-pattern mass. The weight already excludes square factors at $p \leq z$. Failure of squarefreeness at larger primes costs at most

$$(2X + h) \sum_{m > z} m^{-2} = O_h(X/z), \quad (8.8)$$

using complete prefixes to count square multiples. Remove these exceptions as well. Then $\Omega = \omega$ at both endpoints, and every remaining weight is at most one, so its cardinality is at least its weighted mass. This proves (1.10). At the hard endpoint retain both roughness conditions and take

$$\kappa = 1/3750, \quad \epsilon = 1/60000, \quad \kappa - \epsilon = 1/4000, \quad 2\kappa = 1/1875. \quad (8.9)$$

This gives Theorem 1.2. $\square$

Corollary 8.3. *At the maximal z of Theorem 1.3, every sign pair occurs for at least $c_{h,r}X/(\log X)^{2\kappa}$ squarefree pairs with*

$$\Omega(n), \Omega(n + h) \leq (1 - \kappa/2) \log \log X, \quad \kappa = (1 - r)/3750. \quad (8.10)$$

At the hard endpoint the same argument gives $\Omega(n), \Omega(n + h) \leq (1 - 1/7500) \log \log X$; Theorem 1.2 gives the stated smaller upper bound.

Proof. Apply Lemma 8.2, (8.8) and the preceding pattern masses, or take $\epsilon = \kappa/2$ in Remark 1.5. $\square$

9 The full prime-factor weight: phases, norms and mass

In this and the remaining sections use padding $a(u) = 4^{\omega(u)}$, $w = 5^{\omega_Q}$, $\mathcal{Q} = \{p : y < p \leq e^L, p \not\equiv 1 \pmod{5}, p \nmid h\}$, and the original padding probability

$$\rho_j(n, d) = \frac{1}{w(n)} \sum_{\substack{u|n, u \in \mathcal{Q}^{\text{sf}} \\ j\eta \leq \log(du) < (j+1)\eta}} 4^{\omega(u)}. \quad (9.1)$$

The cutoff is $\rho_j \leq K/L$ and the low- $\mathcal{Q}$ cutoff is $400 \log L$. The original matrix has entry $L 4^{\omega(u)} c_d(n) \mathbf{1}_{u|n} \tau_j(n, d) \tau_j(m, d) / \sqrt{w(n)w(m)}$, where now τ_j contains these two cuts and σ_j , and $m = n + hdu$. No full Ω value enters this matrix or its finite-residue comparison.

Proposition 9.1 (Numerical signs and two layers). *Multiply the directed increasing edge with label $q' = du$ by any prescribed $\epsilon(q') \in \{\pm 1\}$, and form*

$$\mathcal{A}_d = \begin{pmatrix} 0 & A_d^+ \\ (A_d^+)^* & 0 \end{pmatrix}. \quad (9.2)$$

Duplicate E in both layers. The moment and spectral estimates of [20, Theorem 5.4 and Proposition 8.3, pp. 24, 41] have the same permissible absolute constants, with K as row constant. Their finite comparison holds with L^{18} by Proposition 3.2.

Proof. Check the original proof in order. The Hilbert–Schmidt expansion [O, pp. 24–25] projects each lifted closed word to a physical word already counted there; it retains unequal consecutive tuples within each half. The projected lifted words form a subset of the physical closed words, and each physical word has at most two lifts, according to its starting layer. For a fixed numerical word its product of edge signs is a scalar of modulus one, independent of residue draws. Remove it from the signed expectation before taking the absolute value. Writing X_w for the unphased contribution of a physical word, the positive word sums therefore obey

$$\sum_{\tilde{w} \in \mathcal{W}_{\text{lift}}} |\mathbb{E}_* X_{\tilde{w}}| \leq 2 \sum_{w \in \mathcal{W}_{\text{phys}}} |\mathbb{E}_* X_w|. \quad (9.3)$$

This compares absolute word sums, as used in [O, p. 24], rather than signed operator norms. The exact singleton identity [O, Lemma 5.5, p. 26] and the unlit, rank and singleton majorants [O, pp. 26–33] are unchanged. The eligible numerical ranges, reciprocal weights and rare witnesses are the ones verified in Proposition 2.9; removing primes up to y only prunes their positive sums. The surviving perfect blocks and forest universe [O, Lemmas 7.1–7.2, pp. 34–37] are unchanged subsets. The padding recursion [O, Lemma 7.3, p. 37] has oriented row bound K , with no numerical sign in the positive row. Completion [O, pp. 37–38] now has external factor $2|\mathcal{D}|^2 M \leq e^{108L}$, the envelope already used there, so the same absolute C_2 may be fixed. The signed finite expansion is covered by Proposition 3.2 before absolute values. The arithmetic and counting proofs [O, Lemmas 5.2–5.3, 5.5–5.6, 6.1–6.5 and 7.1–7.2, pp. 22–37] use no numerical value of K , η or A . They use the bin eligibility only through $du \leq e^{101L}$ and the degree bound on u ; these hold for $\eta \leq 1$. Within the padding-summation step, K enters through the row recursion of Lemma 7.3; its bound is then carried through the completion. The exponent A belongs to the finite-origin comparison, replaced here by Proposition 3.2.

For the one-orientation block A_d^+ , weighted absolute row and column sums, with $\phi = \sqrt{w}$, satisfy

$$\sum_m |A_d^+(n, m)| \frac{\phi(m)}{\phi(n)} \leq K a_d(n), \quad \sum_n |A_d^+(n, m)| \frac{\phi(n)}{\phi(m)} \leq K a_d(m). \quad (9.4)$$

Each is at most the corresponding two-orientation bound of [O, (8.4), p. 40]. Since a_d is constant along an edge, the weighted Cauchy–Schwarz calculation is

$$\begin{aligned} \|A_d^+ v\|^2 &\leq \sum_n K a_d(n) \sum_m |A_d^+(n, m)| \frac{\phi(n)}{\phi(m)} |v_m|^2 \\ &\leq K^2 \sum_m a_d(m)^2 |v_m|^2. \end{aligned} \quad (9.5)$$

Its transpose has the same bound. These are the Schur bounds on the invariant level sets of a_d ; the two-orientation proof in [O, Lemma 8.2, p. 40] uses their sum. For $v = (v_1, v_2)$, with the duplicated projection $v = Ev$, this gives

$$\|\mathcal{A}_d\| \leq K \leq 2K, \quad \sum_d \|\mathcal{A}_d v\|^2 \leq K^2 (8W)^J \|v\|^2 \leq 4K^2 (8W)^J \|v\|^2. \quad (9.6)$$

Thus the original permissible two-orientation constants still apply. If $\mathcal{A} = \sum_d \mathcal{A}_d$ and $A^+ = \sum_d A_d^+$, the exact testing identity for real vectors is

$$\langle (g_1, g_2), \mathcal{A}(g_1, g_2) \rangle = \langle g_1, A^+ g_2 \rangle + \langle g_2, (A^+)^* g_1 \rangle = 2\langle g_1, A^+ g_2 \rangle. \quad (9.7)$$

All edge signs and testing vectors here are real. In particular this recovers twice the ordered increasing-edge bilinear form for the three sign-exponent cases. Input 5.1 and the Markov proof of [O, Proposition 8.3, p. 41] give $K(C_3 \sqrt{W})^J$ outside e^{-2k} of origins, with $C_3 = 3 \max(2, 2\sqrt{8}, eC_2)$ as in [O, (8.7)]. This proves the stated constants without comparing signed submatrix norms by entrywise domination. $\square$

Proposition 9.2 (Positive actual-integer norms). *Uniformly for $99/100 \leq r \leq 1$, $2 \leq y \leq e^{L^{1/12}}$, $e^{L^{18}/2} \leq Y \leq e^{2L^{18}}$, put $B_r(n) = r^{\Omega(n)} S_y(n)$ and $S_r = \prod_{p \in \mathcal{Q}} (1 + 4r^2/p)$. Then*

$$\sum_{n \leq Y} B_r(n)^2 w(n) \ll Y S_r (\log Y)^{r^2-1}, \quad \sum_{n \leq Y} B_r(n)^2 \ll Y (\log Y)^{r^2-1}, \quad (9.8)$$

$$\sum_{n \leq Y} B_r(n)^2 e^{\omega_{\mathcal{T}}(n)} \ll Y (\log Y)^{r^2-1} \exp \left((e-1) r^2 \sum_{p \in \mathcal{T}} 1/p \right) \quad (9.9)$$

for any prime set $\mathcal{T} \subset [2, e^L]$.

Proof. Shiu’s nonnegative mean theorem in the form reproduced in [12, p. 1] bounds a mean by $CY \exp(\sum_{p \leq Y} (F(p) - 1)/p)$ for functions of fixed divisor growth. For the two functions in (9.8), $F \leq 5^{\omega(n)} \leq \tau_5(n)$; for (9.9), $F \leq 3^{\omega(n)} \leq \tau_3(n)$. Thus $F(p^j) \leq A^j$ and $F(n) \leq B_{\xi} n^{\xi}$ have uniform constants for every fixed $\xi > 0$. Apply the theorem with $q = 1$, interval and progression exponents $1/2$ on dyadic long blocks above $\sqrt{Y}$. The small prefix costs $O(\sqrt{Y} \log^4 Y)$. The prime values are $5r^2$ on $\mathcal{Q}$, r^2 elsewhere, or er^2 on $\mathcal{T}$ and r^2 elsewhere. The squarefree cutoff changes only higher powers. Uniformly summable $O(p^{-2})$ errors identify the extra $\mathcal{Q}$ product with S_r , and Mertens gives the displayed powers of $\log Y$. These are bounds on actual integers, rather than a residue model for full Ω . $\square$

Proposition 9.3 (Unsigned full-weight pair mass). *For fixed even h , uniformly for $99/100 \leq r \leq 1$ and $2 \leq y \leq \exp((\log X)^{1/216})$,*

$$c_h X (\log X)^{2r-2} \leq \sum_{n \leq X} r^{\Omega(n) + \Omega(n+h)} S_y(n) S_y(n+h) \leq C_h X (\log X)^{2r-2} \quad (9.10)$$

for sufficiently large X .

Proof. Choose a cutoff z_0 and independently for each $p \leq z_0$ choose two thresholds with $\mathbb{P}(T_i > k) = r^k$, $k \geq 0$; at $r = 1$ take $T_i = \infty$. For $p \leq y$ replace each threshold by its minimum with 2. The complement of the event $p^{T_1} \mid n$ or $p^{T_2} \mid n + h$ has expectation exactly the two local prime-power weights, including the squarefree cap. This is a nonnegative sieve measure on integers and threshold choices, within the probability setup of [6, pp. 1–3].

For squarefree d , expand each bad-prime union into at most three conjunction terms. Conditional on thresholds a consistent term is one prime-power CRT progression, counted with error $O(1)$ regardless of its modulus; inconsistent terms are zero. Average to obtain

$$A_d = Xg(d) + O(3^{\omega(d)}), \quad g(d) = \prod_{p \mid d} g(p). \quad (9.11)$$

Infinite thresholds may be truncated with an infinity option; bounded convergence then applies to each finite sieve sum. The divisor upper/lower identities of Input 2.2 hold for every sample of the bad-prime set and may be integrated.

For $p \nmid h$, $p > y$, a p -adic count gives

$$1 - g(p) = 1 - \frac{2(1 - r)}{p - r}. \quad (9.12)$$

Indeed at one of the two distinct roots the valuation- k density is $(1 - 1/p)p^{-k}$, and summing $1 - 2/p + 2(1 - 1/p) \sum_{k \geq 1} r^k p^{-k}$ gives the formula. For $p \leq y$ the factor is (1.6). The finitely many $p \mid h$ have positive local expectations. All bad events are subsets of the ordinary two-root event, so $g(p) \leq \nu_h(p)/p < 1$. In the uncapped case one may also use $1 - r^k \leq (1 - r)k$ and the mean valuation $2/(p - 1)$. Thus the sieve dimension condition is uniformly bounded by dimension two. Logarithms of generic factors have coefficient $-2(1 - r)/p$; the squarefree cap changes them by $O(p^{-2})$, and fixed exceptional factors stay uniformly positive for $r \geq .99$. Hence

$$\prod_{p \leq z_0} (1 - g(p)) \asymp_h (\log z_0)^{2r-2} \quad (y \leq z_0). \quad (9.13)$$

Take a fixed sufficiently large s_0 with lower-sieve relative error at most $1/4$, $D = X^{1/4}$ and $z_0 = X^{1/(4s_0)}$. Eventually $y \leq z_0 \leq D^{1/2}$. The total CRT error is $O(D \log^2 D)$, smaller than the main term. The finite prime-weighted mass is thus bounded below by a positive multiple of $X(\log X)^{2r-2}$. At each endpoint at most $4s_0 + 1$ prime factors counted with multiplicity exceed z_0 . Restoring them costs at most the fixed factor $r^{8s_0+2} \geq (.99)^{8s_0+2} > 0$, proving the lower bound.

For the upper bound drop S_y and apply [12, Theorem 1, pp. 1–2] to the same fixed polynomials $t, t + h$ and $F(a, b) = r^{\Omega(a) + \Omega(b)}$. All function-class bounds hold uniformly since $0 \leq F \leq 1$; the polynomial, interval and coefficient hypotheses were verified in Proposition 8.1. The positive Euler factors give coefficient $(2r - 2)/p$ with uniformly summable higher-power errors. Dyadic summation completes the bound. The restoring constant is not asserted uniform as r tends to zero. $\square$

10 External padding collision

Input 10.1 (Halász in the Montgomery–Tenenbaum form, [29, Chapter III.4]). *For $Y \geq 3$ and $1 \leq T \leq Y/2$, if F is multiplicative, $F(1) = 1$ and $|F(n)| \leq 1$, then*

$$\frac{1}{Y} \left| \sum_{m \leq Y} F(m) \right| \ll (1 + M_F) e^{-M_F} + T^{-1/2}, \quad M_F = \min_{|v| \leq 2T} \sum_{p \leq Y} \frac{1 - \Re(F(p)p^{-iv})}{p}. \quad (10.1)$$

A rescaling of the height parameter gives the convention $|v| \leq 2T$ in this display. Our use has $T = L^{1000}$ and $Y \geq e^{L^{18}/2}$, so the stated height and length requirements hold for large L .

Proposition 10.2 (Common-prefix collision). *Use the range and notation of Proposition 9.2. Conditional on an integer um , choose a squarefree $\mathcal{Q}$ divisor v with probability $4^{\omega(v)}/w(um)$. Then*

$$\sum_{u \in \mathcal{Q}^{\text{sf}}} \frac{4^{\omega(u)} r^{2\omega(u)}}{u} \sum_{m \leq Y} B_r(m)^2 \mathbb{P}_{v|um}(|\log(u/v)| \leq 1) \ll_h Y S_r(\log Y)^{r^2-1} \frac{\log(2L)}{L}. \quad (10.2)$$

The u -sum is finite and has the same physical prefix $m \leq Y$ for every u .

Proof. Replace the probability by $\mathbb{E}_v \exp(it \log(u/v))$ and call the resulting sum $H(t)$, $|t| \leq 2$. For a $\mathcal{Q}$ prime put $z_p = p^{it}$, $k_p = 4r^2/p$ and

$$C_{0,p} = 1 + \frac{k_p(4 + z_p)}{5}, \quad C_{1,p} = \frac{1 + 4z_p^{-1} + k_p(z_p + 4)}{5}. \quad (10.3)$$

If $p \nmid m$, not choosing it in u contributes 1; choosing it contributes $k_p(z_p/5 + 4/5)$, since the conditional v probabilities are $1/5, 4/5$. If $p \mid m$, these same v probabilities apply whether or not it is chosen in u , giving $C_{1,p}$. The finite Euler identity is therefore

$$H(t) = \left(\prod_{p \in \mathcal{Q}} C_{0,p} \right) \sum_{m \leq Y} F_t(m), \quad F_t(p^j) = \begin{cases} B_r(p^j)^2 C_{1,p}/C_{0,p}, & p \in \mathcal{Q}, \\ B_r(p^j)^2, & p \notin \mathcal{Q}. \end{cases} \quad (10.4)$$

Define F_t multiplicatively with $F_t(1) = 1$. For $c = \Re z_p$, direct algebra gives

$$|C_{0,p}|^2 - |C_{1,p}|^2 = \frac{8}{25}(1 - c)(1 + 5k_p + 2k_p c) \geq 0, \quad \Re C_{0,p} \geq 1. \quad (10.5)$$

Thus $|F_t| \leq 1$, as required in Input 10.1. Its prime values are

$$F_t(p) = \frac{r^2}{5}(1 + 4p^{-it}) + O(p^{-1}) \quad (p \in \mathcal{Q}), \quad F_t(p) = r^2 \quad (p \notin \mathcal{Q}). \quad (10.6)$$

The harmonic sum of these coefficient errors is $O(1)$.

Put $v_0 = \log(2y)$ and $\Psi(t) = \log((1 + L|t|)/(1 + v_0|t|))$. Fixed-modulus PNT and partial summation give, uniformly for $|t| \leq 3$,

$$\sum_{p \in \mathcal{Q}} \frac{1 - \cos(t \log p)}{p} = \frac{3}{4}\Psi(t) + O_h(1), \quad (10.7)$$

$$\left| \prod_{p \in \mathcal{Q}} C_{0,p} \right| \ll_h S_r \exp(-3r^2\Psi(t)/5). \quad (10.8)$$

For the latter, the leading logarithmic difference from $1 + 4r^2/p$ is $-(4r^2/5p)(1 - \cos(t \log p))$; all $O(p^{-2})$ errors are summable. This is the same partial summation as [20, (4.12), p. 18], now with the lower cutoff y .

Take $T = L^{1000}$. For any candidate twist v , (10.6) expands the distance as

$$(1 - r^2) \log \log Y + r^2 S_Y(v) - \frac{4}{5} r^2 S_Q(v) + \frac{4}{5} r^2 S_Q(v + t) + O_h(1), \quad (10.9)$$

where $S_Y(v) = \sum_{p \leq Y} (1 - \cos(v \log p)) / p$ and S_Q is the $\mathcal{Q}$ sum. If $|v| \leq 1$, partial summation of PNT gives $S_Y(v) = \log(1 + |v| \log Y) + O(1)$. The increasing concave function $x \mapsto \log((1 + Lx)/(1 + v_0x))$ vanishes at zero, and is subadditive. Thus $\Psi(t) \leq \Psi(v) + \Psi(v + t)$. Since $\log Y \geq L^{18}/2$, for sufficiently large L ,

$$\log(1 + |v| \log Y) \geq \frac{6}{5} \log(1 + L|v|) \geq \frac{6}{5} \Psi(v) \quad (|v| \leq 1). \quad (10.10)$$

One may verify the inequality separately for $L|v| \leq 1$ and $1 \leq L|v| \leq L$. Using (10.7) in (10.9) therefore gives

$$M_{F_t} \geq (1 - r^2) \log \log Y + \frac{3}{5} r^2 \Psi(t) - O_h(1) \quad (10.11)$$

for these twists. For $1 \leq |v| \leq 2T$, work on $\sigma = 1 + 1/\log Y$. Replacing the sharp prime sum by its $p^{-1/\log Y}$ smoothing costs $O(1)$, by partial summation; higher powers cost $O(1)$ too. The Euler product and Euler summation with cutoff comparable to $|v| + 3$ give $|\zeta(\sigma + iv)| \ll \log(|v| + 3)$, hence $S_Y(v) \geq \log \log Y - O(\log \log(3T))$. Also $S_Q(v) \leq 2 \sum_Q 1/p \leq (3/2) \log L + O_h(1)$, $S_Q(v + t) \geq 0$, and $\log \log Y \geq 18 \log L - O(1)$. These inequalities imply (10.11) in the large-twist range as well. This checks the full Halász minimum, not only a zero twist.

Apply (10.1), (10.8), and $M_{F_t} \leq 2 \log \log Y + O(1)$ to obtain

$$|H(t)| \ll_h Y S_r(\log Y)^{r^2-1} \log(2L) \left(\frac{1 + v_0|t|}{1 + L|t|} \right)^{6r^2/5}. \quad (10.12)$$

The $T^{-1/2} = L^{-500}$ term is absorbed uniformly; the other density and phase factors have magnitude at least L^{-2} in the stated compact r and Y ranges. Finally apply the positive kernel (4.6) to $\log(u/v)$, interchange only finite sums, and use

$$\int_0^2 \left(\frac{1 + v_0 t}{1 + L t} \right)^\nu dt \ll L^{-1} + (v_0/L)^\nu \ll L^{-1}, \quad \nu = 6r^2/5 > 1, \quad v_0 \leq 2L^{1/12}, \quad (11/12)\nu > 1. \quad (10.13)$$

This proves (10.2), retaining complex phases throughout the Euler identity and distance minimization. $\square$

Lemma 10.3 (Uniform upper sieve for marked primes). *There is an absolute $u_0 > 0$ such that, for an interval I of N consecutive positive integers, $z \geq 2$ and $\log N / \log z \geq u_0$, every set $\mathcal{R} \subset \{p \leq z\}$ satisfies*

$$\#\{n \in I : p \nmid n \text{ for every } p \in \mathcal{R}\} \leq 2N \prod_{p \in \mathcal{R}} (1 - 1/p) + O(N^{1/2} \log^2(2N)), \quad (10.14)$$

with an absolute error constant. Consequently, for $0 \leq \delta \leq 1$, $r = 1 - \delta$,

$$\sum_{n \in I} r^{\omega_z(n)} \ll N (\log(2z))^{-\delta} + N^{1/2} \log^2(2N). \quad (10.15)$$

Proof. Use Selberg's upper sieve with divisor level $U = N^{1/4}$. Here is its finite quadratic construction, which also verifies uniformity in the marked set. Let $\mathcal{D}_U = \{d \leq U : d \mid \prod_{p \in \mathcal{R}} p\}$ and

$$G_{\mathcal{R}}(U) = \sum_{d \in \mathcal{D}_U} \frac{1}{\varphi(d)}, \quad F_{\mathcal{R}} = \prod_{p \in \mathcal{R}} (1 - 1/p)^{-1}. \quad (10.16)$$

Put $y_q = \mu(q)/(\varphi(q)G_{\mathcal{R}}(U))$ for $q \in \mathcal{D}_U$, and define $\lambda_d/d = \sum_{q \in \mathcal{D}_U, d|q} \mu(q/d)y_q$. Finite divisor inversion gives $\lambda_1 = 1$ and $y_q = \sum_{d \in \mathcal{D}_U, q|d} \lambda_d/d$. Since $(d, e) = \sum_{q|(d,e)} \varphi(q)$, the quadratic main term is exactly

$$\sum_{d,e \in \mathcal{D}_U} \frac{\lambda_d \lambda_e}{[d, e]} = \sum_{q \in \mathcal{D}_U} \varphi(q) y_q^2 = G_{\mathcal{R}}(U)^{-1}. \quad (10.17)$$

Moreover

$$\lambda_d = \frac{\mu(d)d}{\varphi(d)G_{\mathcal{R}}(U)} \sum_{\substack{m \leq U/d \\ m \prod_{p \in \mathcal{R}} p, (m, d)=1}} \frac{1}{\varphi(m)}, \quad |\lambda_d| \leq d/\varphi(d) \leq \tau(d). \quad (10.18)$$

The sifted indicator is bounded by $(\sum_{d|n} \lambda_d)^2$. Counting each modulus $[d, e]$ in I as $N/[d, e] + O(1)$ therefore gives $N/G_{\mathcal{R}}(U) + O(U^2 \log^2(2U))$, independently of the interval position and marked set.

For the Rankin tail take $\sigma = 1/\log z$. The full squarefree divisor sum is $F_{\mathcal{R}}$, and

$$\begin{aligned} \frac{F_{\mathcal{R}} - G_{\mathcal{R}}(U)}{F_{\mathcal{R}}} &\leq U^{-\sigma} \prod_{p \in \mathcal{R}} \left(1 + \frac{p^{\sigma} - 1}{p}\right) \\ &\leq \exp\left(-\frac{\log U}{\log z} + C_0\right). \end{aligned} \quad (10.19)$$

The constant C_0 is absolute: $p^{\sigma} - 1 \leq (e - 1) \log p / \log z$ for $p \leq z$, and $\sum_{p \leq z} (\log p)/p \ll \log z$ by Mertens' theorem [19, Section 2.2]. Choose $u_0 \geq 4(C_0 + \log 2)$ once and for all. Then $G_{\mathcal{R}}(U) \geq F_{\mathcal{R}}/2$, proving (10.14). This proves the sieve bound for every marked set before any averaging.

Independently mark each prime up to z with probability δ . At a fixed n , the probability that no marked prime divides it is $r^{\omega_z(n)}$. Average (10.14); its error is uniform and

$$\mathbb{E}_{\mathcal{R}} \prod_{p \in \mathcal{R}} (1 - 1/p) = \prod_{p \leq z} (1 - \delta/p) \asymp (\log(2z))^{-\delta}. \quad (10.20)$$

The comparison follows from the logarithmic expansion and Mertens, uniformly in $0 \leq \delta \leq 1$. This gives (10.15). $\square$

11 Cancellation with the full weight

Proof of Theorem 1.4. In this proof abbreviate $C_{\Omega}, W_{\Omega}, \alpha_{\Omega}, \delta_{\Omega}$ by C, W, α, δ . Thus (1.7) fixes $W = 16eC^2$, $\alpha = 1/(1200W)$, $\delta = \sqrt{\alpha/216}$ and $r = 1 - \delta$. Set

$$\begin{aligned} L &= (\log X)^{1/18}, \quad J = \lfloor \alpha \log L \rfloor, \quad \varepsilon = \alpha/100, \\ b &= 1 + r^{-2}, \quad E_0 = 18\delta^2 = \alpha/12, \quad \eta = L^{-\varepsilon}, \quad K = L^{2\alpha \log b + 2E_0 + 4\varepsilon}. \end{aligned} \quad (11.1)$$

The exponent of K is less than 4α ; these are fixed powers of L , as required in Proposition 3.2. The centre supply is Definition 2.6, with the same H_0 and $W \leq V_i \leq 2W$. All label primes exceed y .

Put $f_i(n) = \lambda(n)^{e_i} B(n)$. For $q' = du$, complete additivity of Ω and complete multiplicativity of λ give the exact identity

$$f_1(q'm) f_2(q'(m+h)) = \lambda(q')^{e_1+e_2} r^{2\Omega(q')} f_1(m) f_2(m+h). \quad (11.2)$$

The small-prime-square mask is unchanged, with no coprimality condition on q' and m . Multiply the edge by $\lambda(q')^{e_1+e_2}$ and use Proposition 9.1. The raw term has positive harmonic mass

$$S_{0,r} = \sum_j \sum_{(d,u) \in \mathcal{S}_j} \frac{4^{\omega(u)} r^{2J+2\omega(u)}}{du}, \quad \frac{1}{2} r^{2J} S_r V_* \leq S_{0,r} \leq r^{2J} S_r V_*. \quad (11.3)$$

To prove the lower bound, its padding reciprocal law selects a prime with probability $4r^2/(p+4r^2)$, with expected logarithmic size at most $4L$ and degree at most $4\log L$. Use the two Markov exclusions in Proposition 2.7, now with this law. The raw cofactor cutoff remains in $(Xe^{-\eta}, X]$. The Nair–Tenenbaum upper theorem gives mass $O_h(\eta X(\log X)^{-2\delta})$ in the full missing interval of length comparable to ηX . Here the weights r^Ω are bounded by one, the polynomials are the fixed $t, t+h$, and $\eta X > X^{1/2}$ eventually, so the interval and function-class hypotheses checked in Proposition 8.1 hold. The normalized raw error is $O_h(\eta(\log X)^{-2\delta})$.

We next align all physical prefixes before enlarging padding sums. The positive centred-factor majorant is

$$a_d^+(n) = \prod_{p|d} (\mathbf{1}_{p|n} + 1/p) = \sum_{e|d} \frac{e}{d} \mathbf{1}_{e|n}. \quad (11.4)$$

In a term with $u \mid n$ and $e \mid n$, write $n = uem$; the prime supports of u, e are disjoint and exceed y . Then $B(uem)^2 = r^{2\omega(u)+2\omega(e)} B(m)^2$, and the bin gives

$$Xe^{-\eta} < T_j/(ud) \leq X, \quad T_j^{-1} \leq e^\eta/(Xud), \quad m \leq Xd/e. \quad (11.5)$$

Its coefficient in the normalized one-endpoint squared edge mass is bounded by $e^\eta e r^{2\omega(e)}/(Xd^2)$ times the external sum $\sum_u 4^{\omega(u)} r^{2\omega(u)}/u$ and the $B(m)^2$ sum through $Y = Xd/e$. Only now extend u to all of $\mathcal{Q}^{\text{sf}}$. Apply (9.8) without w , cancel the prefix Y against this coefficient and sum d, e . Since $\sum_{e|d} r^{2\omega(e)} = (1+r^2)^J$, the whole squared mass is

$$\ll_h S_r(\log X)^{r^2-1} (1+r^2)^J V_*. \quad (11.6)$$

Every eligible (d, u) belongs to one bin, so this alignment introduces no bin multiplicity. At the arrival, $n' = n + hdu = ue(m + hd/e)$ and $e \mid d$. Its cofactor $m' = m + hd/e$ is at most $(X+h)d/e$. Enlarge to this common prefix and repeat the same argument. This proves (11.6) at either endpoint, with logarithms comparable to $\log X$.

For a padding-cut failure, apply Cauchy–Schwarz in the positive edge measure. One squared factor has the bad indicator and $B(n)^2$, the other discards the indicator and uses (11.6). The original ρ_j reads only the active $\mathcal{Q}$ primes, so $\rho_j(uem, d) = \rho_j(um, d)$. Since the eligible u and every v in its bin have $|\log(u/v)| \leq \eta \leq 1$,

$$\mathbf{1}_{\rho_j(um, d) > K/L} \leq (L/K) \rho_j(um, d) \leq (L/K) \mathbb{P}_{v|um}(|\log(u/v)| \leq 1). \quad (11.7)$$

After (11.5) the prefix is the same $Y = Xd/e$ for all u . Proposition 10.2 makes the squared bad factor smaller than (11.6) by $O_h(\log(2L)/K)$. At the arrival use $m' = m + hd/e$ and the common prefix just proved; the same inequality applies. Dividing the resulting two-endpoint deletion by (11.3) gives

$$\ll_h (\log X)^{r^2-1} b^J \sqrt{\log(2L)/K}. \quad (11.8)$$

For a $\mathcal{Q}$ -degree failure, retain $\omega(u) \leq 100 \log L$ until $\omega_{\mathcal{Q}}(uem) > 400 \log L$ implies $\omega_{\mathcal{Q}}(m) > 300 \log L$. Markov and (9.9) give a squared-factor saving $O(L^{-200})$, since $\sum_{\mathcal{Q}} 1/p \leq \log L + O_h(1)$; only afterward enlarge the u -sum. Cauchy–Schwarz saves L^{-100} . For a $\mathcal{P}$ -degree failure, $\omega_{\mathcal{P}}(uem) \leq J + \omega_{\mathcal{P}}(m)$ and

$$e^{-(6W-1)J} \exp(2(e-1)WJ) \leq e^{-2WJ} \quad (W \geq 10). \quad (11.9)$$

This gives e^{-WJ} after Cauchy–Schwarz. For rare-site failures discard $B(n)B(n+hdu) \leq 1$ and use the positive deletion proof of [20, Lemma 4.6, pp. 19–20], with (2.20) and Proposition 3.2. The

ratio of SV_* , where $S = \prod_{p \in Q} (1 + 4/p)$, to $S_{0,r}$ and the inverse mass density are fixed powers of L , absorbed in its exponential error. The complete relative deletion error is

$$\ll_h L^{E_0} b^J \{ \sqrt{\log(2L)/K} + L^{-100} + e^{-WJ} \} + e^{-L^{0.8}}. \quad (11.10)$$

Here $(\log X)^{(r^2-1)+2\delta} = (\log X)^{\delta^2} = L^{E_0}$. The nonrare estimates were actual-integer positive norms, not a finite-residue model for B .

For the spectral term use Proposition 9.1 and the unweighted signed-trace comparison of Proposition 3.2. The actual squared testing norms of $\sqrt{w}f_i$ average to $O_h(MS_r(\log X)^{r^2-1})$, by Proposition 9.2; Cauchy–Schwarz handles the bilinear form. The overlap identity (6.5) and its deterministic exceptional-block bound apply to the two layers. All normalizations are fixed powers of L , so their exponential errors remain negligible. After the bins and (11.3), the relative spectral term is

$$\ll_h L^{E_0} \frac{K}{\eta} \left(\frac{C}{r^2 \sqrt{W}} \right)^J + e^{-L^{0.8}}. \quad (11.11)$$

The S_r in the actual norm cancels the S_r in the raw mass.

Finally repeat the centering proof of [20, Lemmas 4.1–4.2, 4.4, pp. 13–16] with the present f_i . At least one signed factor has prime value $-r$ at every prime, since S_y changes only higher powers. For that factor,

$$M(f_i; Y, Q) = rM(\lambda; Y, Q) + (1-r) \sum_{p \leq Y} 1/p \geq \frac{1}{4} \log \log Y - O(1). \quad (11.12)$$

Use precisely Input 2.5; $Y \geq e^{L^{18/2}}$, $H_0/2 \leq D \leq e^{2L} \ll Y$, and $Q = \min((\log Y)^{1/125}, (\log D)^5)$. The height is $|t| \leq Y$. Its errors give the short Fourier bound $DL^{-\Gamma}$ with

$$a = 994999/1000000, \quad \Gamma = 2571/100000, \quad s = 1/1000000, \quad a < 199/200, \quad \Gamma < 18/700. \quad (11.13)$$

In choosing the final threshold also require

$$\frac{1}{2} L^{199/200-a} \geq u_0, \quad z_m = e^{L^a}, \quad \frac{\log D}{\log z_m} \geq u_0 \quad (H_0/2 \leq D \leq e^{2L}), \quad (11.14)$$

where u_0 is fixed in Lemma 10.3. This is possible because $199/200 - a = 10^{-6} > 0$. The distance is eventually at least $(18/5) \log L$. The short error is $O(L^{-199/200} \log L)$, and the long error is $O(L^{-18/700})$. Proposition 6.1 has $\|\mathcal{B}\|_\infty \ll L^{-a}$ and fourth moment $O(D^{-1} L^{-4a})$ at this changed cutoff.

The averaged squared short-window norms are bounded by $D(\log Y)^{r^2-1}$: count each integer in at most D windows and apply (9.8) without w , followed by Cauchy–Schwarz in the origin average. For the pointwise other factor, use

$$B(m) \leq r^\omega_{p \leq e^{L^a}(m)}, \quad |G_v| \ll_h DL^{-a\delta}. \quad (11.15)$$

For the pointwise assertion, take $N = (2h+1)D$ and $z = z_m$ in Lemma 10.3. Condition (11.14) gives $\log N / \log z_m \geq u_0$. For every marked-prime set the Selberg sieve has divisor level $N^{1/4}$ and the Rankin estimate gives $G_{\mathcal{R}}(N^{1/4}) \geq F_{\mathcal{R}}/2$; average only after applying this uniform bound. The expected product is $\prod_{p \leq z_m} (1 - \delta/p) \asymp L^{-a\delta}$. The remainder $O_h(D^{1/2} \log^2(2D))$ is smaller than $DL^{-a\delta}$ throughout the stated D range, for sufficiently large L . Dropping the squarefree mask and taking absolute values of the endpoint function therefore proves $|G_v| \ll_h DL^{-a\delta}$ in every

positive window. The separately required CRT estimate in Proposition 6.1 remains $D^{-1}e^{O(L^a \log L)}$, negligible because $a < 199/200$.

For a nonempty partial-centred subset I , put $q_I = \prod_{i \in I} p_i$; the physical shift is hq_I . The subset extracts only $J - |I|$ centre primes. Relative to r^{2J} in the raw mass its missing factor is $r^{-2|I|}$, so summing subsets gives $(1 + r^{-2})^J = b^J$. All remaining numerical signs have modulus one and leave the shift-polynomial estimate unchanged. Padding extraction has harmonic mass S_r . The Fourier split $|\mathcal{B}| \leq L^{-1-s}$ now gives the two relative terms

$$L^{E_0 + \varepsilon + \alpha \log b - s}, \quad L^{5(1-a) + 4s - \Gamma + (36-a)\delta + \varepsilon + \alpha \log b}. \quad (11.16)$$

The squared-norm loss is L^{E_0} ; the pointwise loss is $L^{36\delta - a\delta}$. If only the second factor is signed, reindex by hq_I before Fourier averaging, put the signed factor in F , and allow negative offsets in G . Removing the initial $O_h(D)$ origins makes all arguments positive and costs $O_h(D/Y)$, exponentially little. Hence this centering bound covers all three nonzero exponent pairs. The input is the classical MRT distance bound; the half-plane $\Re s > 7/8$ is not used.

We record the numerical margins. Since $C_3 \geq 6\sqrt{8}$, $C^2 \geq 288$; using $e > 8/3$ gives $\alpha < 1/14700000$, $\delta < 1/56000$ and $r \geq .99$. Also $\log b < .71$ and

$$\log(4\sqrt{e}r^2) - 2\log b = \frac{1}{2} + 6\log r - 2\log((1+r^2)/2) > 2/5. \quad (11.17)$$

For example $\log r \geq -1/99$ and the last logarithm is nonpositive. The exponent in (11.11) is at most

$$3E_0 + 5\varepsilon - (2/5)\alpha = \alpha/4 + \alpha/20 - 2\alpha/5 = -\alpha/10. \quad (11.18)$$

The padding deletion exponent in (11.10) is -2ε before its square-root logarithm; the raw error has exponent $-\varepsilon$, and both degree terms are smaller. For the two centering terms, $E_0 + 2\varepsilon + \alpha \log b < s$, and

$$5(1-a) + 4s + 36/56000 + 1/10000000 < .02571 = \Gamma; \quad \text{the left side is less than } .025652. \quad (11.19)$$

Thus both exponents in (11.16) are less than $-\varepsilon$. Absorb fixed logarithmic factors into $\varepsilon/2$ to obtain

$$\left| \sum_{n \leq X} f_1(n)f_2(n+h) \right| \ll_h X(\log X)^{-2\delta} L^{-\varepsilon/2}, \quad L^{-\varepsilon/2} = (\log X)^{-\alpha/3600}. \quad (11.20)$$

This is (1.8); Proposition 9.3 proves (1.9). All constants were fixed before h, X ; only the final lower threshold depends on h . $\square$

Corollary 11.1. *With (1.7), let $M_0(X, y) = \sum_{n \leq X} B(n)B(n+h)$. For each sign pair the weighted mass is*

$$\sum_{\substack{n \leq X \\ (\lambda(n), \lambda(n+h)) = (s, t)}} B(n)B(n+h) = \frac{1}{4} M_0(X, y) (1 + O_h((\log X)^{-\gamma_\Omega})). \quad (11.21)$$

Every sign pattern also occurs for $\gg_h X/(\log X)^{2\delta_\Omega}$ integers $n \leq X$ with both endpoints squarefree and

$$\Omega(n), \Omega(n+h) \leq (1 - \delta_\Omega/2) \log \log X. \quad (11.22)$$

Proof. Expand (6.17), use all three signed estimates in Theorem 1.4, and divide by (9.10); no exact unsigned Euler leading constant is asserted. For the cardinality assertion take $y = \exp((\log X)^{1/216})$, $\tau = 1 - \delta/2$ and $u = 1 + \delta/(4r)$. Then $ru = 1 - 3\delta/4 < 1$. The Nair–Tenenbaum theorem for the nonnegative bounded function $(ru)^{\Omega(n)} r^{\Omega(n+h)}$ gives $O_h(X(\log X)^{ru+r-2})$; its fixed-polynomial hypotheses are those already verified, and squarefree masks may be dropped for this upper bound. Relative to the unsigned mass the Markov exponent is $r(u-1) - \tau \log u$. Equation (8.5) with $\kappa = \delta$ makes it at most $-\delta^2/32$. The second endpoint is identical. The squarefree support at small primes and (8.8) with $z = y$ remove the remaining $O_h(X/y)$ exceptions. Each retained weight is at most one, so positive weighted mass bounds the cardinality below. This proves (11.22). $\square$

References

- [1] Louay M. Bazzi. Polylogarithmic independence can fool DNF formulas. *SIAM Journal on Computing*, 38(6):2220–2272, 2009. doi: 10.1137/070691954.
- [2] Mark Braverman. Poly-logarithmic independence fools AC^0 circuits. In *IEEE Conference on Computational Complexity*, pages 3–8, 2009. URL <https://www.cs.cmu.edu/~odonnell/hits09/braverman-fooling-AC0.pdf>. Six-page author version, Main Theorem and Corollary 2, p. 1.
- [3] Julien Cassaigne, Sébastien Ferenczi, Christian Mauduit, Joël Rivat, and András Sárközy. On finite pseudorandom binary sequences. III. The Liouville function. I. *Acta Arithmetica*, 87(4): 367–390, 1999.
- [4] Dimitrios Charamaras and Florian K. Richter. Asymptotic independence of $\Omega(n)$ and $\Omega(n+1)$ along logarithmic averages, 2024. URL <https://arxiv.org/abs/2412.17583v3>. arXiv:2412.17583v3, 17 August 2026; version 1, 23 December 2024.
- [5] Hédi Daboussi and András Sárközy. On the correlation of the truncated Liouville function. *Acta Arithmetica*, 108(1):61–76, 2003. doi: 10.4064/aa108-1-6.
- [6] Kevin Ford. Sieve methods, 2023. URL <https://ford126.web.illinois.edu/sieve2023.pdf>. Lecture notes, Spring 2023, Theorem 3.6.
- [7] D. A. Goldston, S. W. Graham, J. Pintz, and C. Y. Yıldırım. Small gaps between almost primes, the parity problem, and some conjectures of Erdős on consecutive integers, 2008. URL <https://arxiv.org/abs/0803.2636v1>. Version 1, 18 March 2008.
- [8] Jizhou Guo. Goldbach-type representations with prescribed Liouville signs, October 2026. URL <https://github.com/aster2024/openai-math-followups/tree/main/liouville-goldbach>. Preprint, 8 October 2026.
- [9] Jizhou Guo. Sums of two rough numbers with prescribed Liouville signs, 2026. URL <https://github.com/aster2024/openai-math-followups/tree/main/rough-goldbach-parity>. Preprint, 9 October 2026.
- [10] Jizhou Guo. Shifted primes with a prescribed parity of the number of prime factors, 2026. URL <https://github.com/aster2024/openai-math-followups/tree/main/shifted-prime-parity>. Preprint, 9 October 2026.

- [11] Harald Andrés Helfgott and Maksym Radziwiłł. Expansion, divisibility and parity, 2021. URL <https://arxiv.org/abs/2103.06853v2>. Version 2, 13 April 2021.
- [12] Kevin Henriot. Nair–Tenenbaum bounds uniform with respect to the discriminant, 2011. URL <https://arxiv.org/abs/1102.1643v1>. Version 1, 8 February 2011.
- [13] Jason Hickey. salt: a formal mathematics project in Lean 4, 2026. URL <https://github.com/jyh/salt>. GitHub repository; pull requests 44 (merged 5 September 2026) and 126 (merged 14 September 2026); commit ffd2a9e2974fb5c79cd3cfedaa51ff26a5e685de, 9 October 2026.
- [14] Scott D. Hughes. Effective logarithmic two-point Chowla bounds in every window, 2026. URL <https://arxiv.org/abs/2609.28526v1>. arXiv:2609.28526v1, 22 September 2026.
- [15] Alexander P. Mangerel. On the bivariate Erdős–Kac theorem and correlations of the Möbius function. *Mathematical Proceedings of the Cambridge Philosophical Society*, 169(3):547–605, 2020. doi: 10.1017/S0305004119000288. Theorem numbering follows arXiv:1612.09544v2, 28 January 2017.
- [16] Kaisa Matomäki, Maksym Radziwiłł, and Terence Tao. An averaged form of Chowla’s conjecture. *Algebra & Number Theory*, 9:2167–2196, 2015. doi: 10.2140/ant.2015.9.2167. URL <https://arxiv.org/abs/1503.05121v3>. Corrected version arXiv:1503.05121v3, 1 March 2022; Theorem 1.7 and equation (1.12).
- [17] Kaisa Matomäki, Maksym Radziwiłł, and Terence Tao. Sign patterns of the Liouville and Möbius functions, 2015. URL <https://arxiv.org/abs/1509.01545v2>. Version 2, 23 September 2015.
- [18] Alastair McCaffer. Twin primes via Liouville parity detection and sieve-weighted Chowla estimates, 2026. URL <https://doi.org/10.5281/zenodo.20388364>. Zenodo, 26 May 2026.
- [19] Hugh L. Montgomery and Robert C. Vaughan. *Multiplicative Number Theory I: Classical Theory*, volume 97 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2007. doi: 10.1017/CBO9780511618314. URL <https://websites.umich.edu/~hlm/mt1.html>.
- [20] OpenAI. Ordinary two-point correlations of multiplicative functions, 2026. URL <https://github.com/openai/math/blob/adc7f1241b42e322a6451854ab7e4b4c146bf78a/preprints/Ordinary-two-point-correlations-of-multiplicative-functions-September-24-2026/final.pdf>. OpenAI Math Release preprint, 24 September 2026.
- [21] OpenAI. The joint Dickman law for consecutive integers, September 2026. URL <https://github.com/openai/math/blob/adc7f1241b42e322a6451854ab7e4b4c146bf78a/preprints/The-joint-Dickman-law-for-consecutive-integers-September-24-2026/paper.pdf>. 24 September 2026. Repository commit adc7f1241b42e322a6451854ab7e4b4c146bf78a.
- [22] Cédric Pilatte. Improved bounds for the two-point logarithmic Chowla conjecture, 2023. URL <https://arxiv.org/abs/2310.19357v3>. arXiv:2310.19357v3, 25 August 2026; version 1, 30 October 2023.
- [23] János Pintz. Are there arbitrarily long arithmetic progressions in the sequence of twin primes? II, 2010. URL <https://arxiv.org/abs/1004.1067>. 7 April 2010.

- [24] Sergei Preobrazhenskii and Tatyana Preobrazhenskaya. Correlations of the Möbius and Liouville functions and the twin prime conjecture, 2015. URL <https://arxiv.org/abs/1405.0682v7>. arXiv:1405.0682v7, 4 May 2022 (the text of version 6, 20 October 2015, is the same); listed on arXiv under the title “A note on correlations of arithmetic functions”.
- [25] Alexander A. Razborov. A simple proof of Bazzi’s theorem, 2008. URL <https://homepage.mi-ras.ru/~razborov/bazzi.pdf>. Author manuscript, 10 September 2008, Theorem 1.
- [26] Terence Tao. The logarithmically averaged Chowla and Elliott conjectures for two-point correlations. *Forum of Mathematics, Pi*, 4:e8, 2016. doi: 10.1017/fmp.2016.6. Theorem numbering follows arXiv:1509.05422v4, 29 July 2016.
- [27] Terence Tao and Joni Teräväinen. The structure of correlations of multiplicative functions at almost all scales, with applications to the Chowla and Elliott conjectures. *Algebra & Number Theory*, 13(9):2103–2150, 2019. doi: 10.2140/ant.2019.13.2103. Numbering follows arXiv:1809.02518v2.
- [28] Terence Tao and Joni Teräväinen. Quantitative correlations and some problems on prime factors of consecutive integers, 2025. URL <https://arxiv.org/abs/2512.01739v2>. arXiv:2512.01739v2, 25 April 2026; version 1, 1 December 2025.
- [29] Gérald Tenenbaum. *Introduction to Analytic and Probabilistic Number Theory*, volume 163 of *Graduate Studies in Mathematics*. American Mathematical Society, third edition, 2015. doi: 10.1090/gsm/163. URL <https://doi.org/10.1090/gsm/163>. Chapter III.4, Distribution of additive functions and mean values of multiplicative functions.

Use of AI

The proofs and the text of this paper were produced with AI systems (GPT-6 Astra, GPT-6.1 Sol and Claude Opus 5.5) under the author’s direction. They were checked by independent reviews with AI models, and the algebraic identities by computation on finite ranges (programs in the ancillary files). The author takes responsibility for the content.