

# Bootstrap and Bivariate Regression

Kirk Bansak

February 14, 2023

# Problem Sets

- Problem Set 2 is now due.
- Problem Set 3 is due next Thursday, February 23, by 12:29PM PT.
- Reminder: You should not use any R packages or functionalities beyond what is already loaded when you open up RStudio, unless a package is explicitly mentioned.
- The packages automatically loaded by default (and hence allowed) are: `base`, `datasets`, `grDevices`, `graphics`, `methods`, `stats`, `utils`
- You may always use `ggplot2`.

# First, revisiting the bootstrap...

When conducting statistical analyses, we will often want to not only produce estimates/predictions, but also quantify the **uncertainty** surrounding those estimates/predictions.

Two possible methods:

- 1 Classic statistical inference
- 2 **Bootstrap**

# Welcome to the Bootstrap World

## Real World

Unknown  
probability  
distribution

Observed random  
sample

$$P \longrightarrow Z = (Z_1, \dots, Z_n)$$

↓

$$\hat{\theta} = s(Z)$$

Statistic of interest

# Welcome to the Bootstrap World

## Real World

Unknown  
probability  
distribution

Observed random  
sample

$$P \longrightarrow Z = (Z_1, \dots, Z_n)$$


↓

$$\hat{\theta} = s(Z)$$

Statistic of interest


- Given no other information, the observed sample  $Z$  contains all the available information about the underlying population distribution  $P$

# Welcome to the Bootstrap World



- Given no other information, the observed sample  $Z$  contains all the available information about the underlying population distribution  $P$
- Thus, **resampling** from  $Z$  is the best guide to what can be expected from resampling from  $P$


# Welcome to the Bootstrap World



- Given no other information, the observed sample  $Z$  contains all the available information about the underlying population distribution  $P$
- Thus, **resampling** from  $Z$  is the best guide to what can be expected from resampling from  $P$
- Just like we drew  $Z$  from  $P$ , let's draw a **resample**  $Z^*$  from  $Z$



# Welcome to the Bootstrap World



- Given no other information, the observed sample  $Z$  contains all the available information about the underlying population distribution  $P$
- Thus, **resampling** from  $Z$  is the best guide to what can be expected from resampling from  $P$
- Just like we drew  $Z$  from  $P$ , let's draw a **resample**  $Z^*$  from  $Z$
- If  $n$  is sufficiently large, the observed sample  $Z$  should be a good approximation of  $P$  (i.e. treat  $Z$  as  $\hat{P}$ )

# (Nonparametric) Bootstrap

- 1 Decide on a summary statistic  $s$  of interest.
  - e.g. if  $Z$  contains only a single variable, perhaps the mean of that variable

# (Nonparametric) Bootstrap

- 1 Decide on a summary statistic  $s$  of interest.
  - e.g. if  $Z$  contains only a single variable, perhaps the mean of that variable
- 2 Draw  $B$  resamples of  $n$  observations from  $Z$  **with replacement**.
  - Note: If  $Z$  contains multiple variables, resampled observations should be kept intact (i.e. full rows should be drawn).

Call the  $b$ th resample  $Z_b^*$ .

# (Nonparametric) Bootstrap

- 1 Decide on a summary statistic  $s$  of interest.
  - e.g. if  $Z$  contains only a single variable, perhaps the mean of that variable
- 2 Draw  $B$  resamples of  $n$  observations from  $Z$  **with replacement**.
  - Note: If  $Z$  contains multiple variables, resampled observations should be kept intact (i.e. full rows should be drawn).

Call the  $b$ th resample  $Z_b^*$ .

- 3 For each  $Z_b^*$ , compute  $s(Z_b^*)$  and store it, creating the set  $\{s(Z_1^*), \dots, s(Z_B^*)\}$

# (Nonparametric) Bootstrap

- 1 Decide on a summary statistic  $s$  of interest.
  - e.g. if  $Z$  contains only a single variable, perhaps the mean of that variable
- 2 Draw  $B$  resamples of  $n$  observations from  $Z$  **with replacement**.
  - Note: If  $Z$  contains multiple variables, resampled observations should be kept intact (i.e. full rows should be drawn).

Call the  $b$ th resample  $Z_b^*$ .

- 3 For each  $Z_b^*$ , compute  $s(Z_b^*)$  and store it, creating the set  $\{s(Z_1^*), \dots, s(Z_B^*)\}$
- 4 You can assess the distribution of  $\{s(Z_1^*), \dots, s(Z_B^*)\}$ ; for instance, its standard deviation represents the **standard error** of your statistic.

# (Nonparametric) Bootstrap

- 1 Decide on a summary statistic  $s$  of interest.
  - e.g. if  $Z$  contains only a single variable, perhaps the mean of that variable
- 2 Draw  $B$  resamples of  $n$  observations from  $Z$  **with replacement**.
  - Note: If  $Z$  contains multiple variables, resampled observations should be kept intact (i.e. full rows should be drawn).

Call the  $b$ th resample  $Z_b^*$ .

- 3 For each  $Z_b^*$ , compute  $s(Z_b^*)$  and store it, creating the set  $\{s(Z_1^*), \dots, s(Z_B^*)\}$
- 4 You can assess the distribution of  $\{s(Z_1^*), \dots, s(Z_B^*)\}$ ; for instance, its standard deviation represents the **standard error** of your statistic.
- 5 Or to compute 95% CI, use 2.5/97.5 percentiles of  $\{s(Z_1^*), \dots, s(Z_B^*)\}$  as the lower/upper bounds (**bootstrap percentile CI**)

Does this look familiar?

Does this look familiar?

You just programmed a bootstrap on the problem set!



To *R*!

# Quick Review

# Supervised Learning

We assume some relationship between  $Y$  and  $X = (X_1, X_2, \dots, X_p)$ , such that:

$$Y = f(X) + \epsilon$$

- $f$  is some fixed but unknown function of  $X_1, X_2, \dots, X_p$
- $\epsilon$  is a random irreducible **error term**

## Supervised Learning

Using observed data ( $X$  and  $Y$ ) to estimate  $f$  with  $\hat{f}$

**Ultimate Goal:** Build a  $\hat{f}$  that is close to  $f$

# Using a Linear Model for $f$

For each election  $i$ , where  $i$  is used to index different observations,

Let:

$Vote_i =$  Incumbent Vote Share in election  $i$ .

$Approval_i =$  Incumbent Net Approval in election  $i$ .

We want to predict incumbent vote share using approval as our input.

# Using a Linear Model for $f$

For each election  $i$ , where  $i$  is used to index different observations,

Let:

$Vote_i$  = Incumbent Vote Share in election  $i$ .

$Approval_i$  = Incumbent Net Approval in election  $i$ .

We want to predict incumbent vote share using approval as our input.

Thus, we need a function  $f$  that relates  $Approval_i$  to  $Vote_i$ , and one fundamental approach is to employ a linear function:

# Using a Linear Model for $f$

For each election  $i$ , where  $i$  is used to index different observations,

Let:

$Vote_i$  = Incumbent Vote Share in election  $i$ .

$Approval_i$  = Incumbent Net Approval in election  $i$ .

We want to predict incumbent vote share using approval as our input.

Thus, we need a function  $f$  that relates  $Approval_i$  to  $Vote_i$ , and one fundamental approach is to employ a linear function:

$$Vote_i = f(Approval_i) + \epsilon_i$$

$$Vote_i = \beta_0 + \beta_1 Approval_i + \epsilon_i$$

# Bivariate Regression: Function Perspective

Employing linear function  $f$  to relate  $Approval_i$  to  $Vote_i$ :

$$Vote_i = \beta_0 + \beta_1 Approval_i + \epsilon_i$$

$\beta_0$  and  $\beta_1$  are two unknown quantities known as the model **coefficients** or **parameters**.

# Bivariate Regression: Function Perspective

Employing linear function  $f$  to relate  $Approval_i$  to  $Vote_i$ :

$$Vote_i = \beta_0 + \beta_1 Approval_i + \epsilon_i$$

$\beta_0$  and  $\beta_1$  are two unknown quantities known as the model **coefficients** or **parameters**.

Now we need to use our training data to estimate  $\beta_0$  and  $\beta_1$ .



# Bivariate Regression: Function Perspective

Employing linear function  $f$  to relate  $Approval_i$  to  $Vote_i$ :

$$Vote_i = \beta_0 + \beta_1 Approval_i + \epsilon_i$$


$\beta_0$  and  $\beta_1$  are two unknown quantities known as the model **coefficients** or **parameters**.

Now we need to use our training data to estimate  $\beta_0$  and  $\beta_1$ .


By using our estimates,  $\hat{\beta}_0$  and  $\hat{\beta}_1$ , we can then make predictions for  $Vote_i$ :

$$\widehat{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 Approval_i$$


# Bivariate Regression: Geometric Perspective



# Bivariate Regression: Geometric Perspective



# Test Your Knowledge



- 1 What corresponds to  $\hat{\beta}_0$ ?
- 2 What corresponds to  $\hat{\beta}_1$ ?
- 3 What corresponds to  $\hat{\epsilon}_i$ ?
- 4 What's the difference between  $Y$  and  $\hat{Y}$ ?
- 5 Bonus: where are we interpolating and where are we extrapolating?

# Fitting a Bivariate Regression

## Regression Model:

$$\text{Vote}_i = \beta_0 + \beta_1 \text{Approval}_i + \epsilon_i$$

# Fitting a Bivariate Regression

## Regression Model:

$$\text{Vote}_i = \beta_0 + \beta_1 \text{Approval}_i + \epsilon_i$$

## Fitted Regression:

$$\text{Vote}_i = \underbrace{\hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i}_{\text{Vote}_i} + \hat{\epsilon}_i$$

# Fitting a Bivariate Regression

## Regression Model:

$$\text{Vote}_i = \beta_0 + \beta_1 \text{Approval}_i + \epsilon_i$$

## Fitted Regression:

$$\text{Vote}_i = \underbrace{\hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i}_{\widehat{\text{Vote}_i}} + \hat{\epsilon}_i$$

## Residuals:

$$\hat{\epsilon}_i = \text{Vote}_i - \widehat{\text{Vote}_i}$$

# Fitting a Bivariate Regression

## Regression Model:

$$\text{Vote}_i = \beta_0 + \beta_1 \text{Approval}_i + \epsilon_i$$

## Fitted Regression:

$$\text{Vote}_i = \underbrace{\hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i}_{\widehat{\text{Vote}}_i} + \hat{\epsilon}_i$$

## Residuals:

$$\hat{\epsilon}_i = \text{Vote}_i - \widehat{\text{Vote}}_i$$


## Goal:

Choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to 'make residuals small' (i.e. get  $\widehat{\text{Vote}}_i$  'close to'  $\text{Vote}_i$ )



# Bivariate Regression: Geometric Perspective

Choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  such that the regression line is 'close to' the data points.



# How Should We Choose the Parameter Values?

## **Preliminary Ideas:**

Choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the following?

# How Should We Choose the Parameter Values?

## Preliminary Ideas:

Choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the following?

$$\sum_{i=1}^N |\hat{\epsilon}_i| = \sum_{i=1}^N \left| \text{Vote}_i - \widehat{\text{Vote}}_i \right|$$

# How Should We Choose the Parameter Values?

## Preliminary Ideas:

Choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the following?

$$\sum_{i=1}^N |\hat{\epsilon}_i| = \sum_{i=1}^N \left| \text{Vote}_i - \widehat{\text{Vote}}_i \right|$$

Good idea, but not the most common approach.

# Fitting a Bivariate Regression

The classic approach is to choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the **sum of squared residuals**:

$$\sum_{i=1}^N \hat{\epsilon}_i^2$$

# Fitting a Bivariate Regression

The classic approach is to choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the **sum of squared residuals**:

$$\sum_{i=1}^N \hat{\epsilon}_i^2$$

$$\sum_{i=1}^N \hat{\epsilon}_i^2 = \sum_{i=1}^N \left( \text{Vote}_i - \widehat{\text{Vote}_i} \right)^2$$

# Fitting a Bivariate Regression

The classic approach is to choose  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the **sum of squared residuals**:

$$\sum_{i=1}^N \hat{\epsilon}_i^2$$

$$\sum_{i=1}^N \hat{\epsilon}_i^2 = \sum_{i=1}^N \left( \text{Vote}_i - \widehat{\text{Vote}_i} \right)^2$$

$$\sum_{i=1}^N \hat{\epsilon}_i^2 = \sum_{i=1}^N \left( \text{Vote}_i - \hat{\beta}_0 - \hat{\beta}_1 \text{Approval}_i \right)^2$$

In statistics, the quantity we want to minimize is often called the:  
**Loss Function** (also sometimes called the **Objective Function**)



In statistics, the quantity we want to minimize is often called the: **Loss Function** (also sometimes called the **Objective Function**)

The following loss function

$$\sum_{i=1}^N \hat{\epsilon}_i^2$$

is the **Sum of Squared Residuals**.

In statistics, the quantity we want to minimize is often called the: **Loss Function** (also sometimes called the **Objective Function**)

The following loss function

$$\sum_{i=1}^N \hat{\epsilon}_i^2$$

is the **Sum of Squared Residuals**.

It is also often called the **Residual Sum of Squares**, **Sum of Squared Error**, or **Residual Squared Error**

In statistics, the quantity we want to minimize is often called the: **Loss Function** (also sometimes called the **Objective Function**)

The following loss function

$$\sum_{i=1}^N \hat{\epsilon}_i^2$$

is the **Sum of Squared Residuals**.

It is also often called the **Residual Sum of Squares**, **Sum of Squared Error**, or **Residual Squared Error**

**Ordinary Least Squares (OLS) Regression:** Choosing parameter values (in bivariate case,  $\hat{\beta}_0$  and  $\hat{\beta}_1$ ) to minimize the sum of squared residuals

# Bivariate Ordinary Least Squares Regression

**OLS Regression:** Choose the coefficients to **minimize the sum of squared residuals**.

In our bivariate case:

# Bivariate Ordinary Least Squares Regression

**OLS Regression:** Choose the coefficients to **minimize the sum of squared residuals**.

In our bivariate case:

$$\arg \min_{\hat{\beta}_0, \hat{\beta}_1} \sum_{i=1}^N \left( \text{Vote}_i - \hat{\beta}_0 - \hat{\beta}_1 \text{Approval}_i \right)^2$$

# Bivariate Ordinary Least Squares Regression

**OLS Regression:** Choose the coefficients to **minimize the sum of squared residuals**.

In our bivariate case:

$$\arg \min_{\hat{\beta}_0, \hat{\beta}_1} \sum_{i=1}^N \left( \text{Vote}_i - \hat{\beta}_0 - \hat{\beta}_1 \text{Approval}_i \right)^2$$

Minimization of loss function can be accomplished via:

- 1 Calculus
- 2 Computer optimization algorithms

# Bivariate Ordinary Least Squares Solution

$$\arg \min_{\hat{\beta}_0, \hat{\beta}_1} \sum_{i=1}^N \left( y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i \right)^2$$

# Bivariate Ordinary Least Squares Solution

$$\arg \min_{\hat{\beta}_0, \hat{\beta}_1} \sum_{i=1}^N \left( y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i \right)^2$$

Recall that finding an extreme (minimum or maximum) involves taking the derivative and setting to zero.



# Bivariate Ordinary Least Squares Solution

$$S(\hat{\beta}_0, \hat{\beta}_1) = \sum_{i=1}^N (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$$

# Bivariate Ordinary Least Squares Solution

$$\begin{aligned} S(\hat{\beta}_0, \hat{\beta}_1) &= \sum_{i=1}^N (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \\ &= \sum_{i=1}^N (y_i^2 - 2y_i \hat{\beta}_0 - 2y_i \hat{\beta}_1 x_i + \hat{\beta}_0^2 + 2\hat{\beta}_0 \hat{\beta}_1 x_i + \hat{\beta}_1^2 x_i^2) \end{aligned}$$

# Bivariate Ordinary Least Squares Solution

$$\begin{aligned} S(\hat{\beta}_0, \hat{\beta}_1) &= \sum_{i=1}^N (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \\ &= \sum_{i=1}^N (y_i^2 - 2y_i\hat{\beta}_0 - 2y_i\hat{\beta}_1 x_i + \hat{\beta}_0^2 + 2\hat{\beta}_0\hat{\beta}_1 x_i + \hat{\beta}_1^2 x_i^2) \end{aligned}$$

$$\frac{\partial S(\hat{\beta}_0, \hat{\beta}_1)}{\partial \hat{\beta}_0} =$$

# Bivariate Ordinary Least Squares Solution

$$\begin{aligned} S(\hat{\beta}_0, \hat{\beta}_1) &= \sum_{i=1}^N (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \\ &= \sum_{i=1}^N (y_i^2 - 2y_i \hat{\beta}_0 - 2y_i \hat{\beta}_1 x_i + \hat{\beta}_0^2 + 2\hat{\beta}_0 \hat{\beta}_1 x_i + \hat{\beta}_1^2 x_i^2) \end{aligned}$$

$$\frac{\partial S(\hat{\beta}_0, \hat{\beta}_1)}{\partial \hat{\beta}_0} = \sum_{i=1}^N (-2y_i + 2\hat{\beta}_0 + 2\hat{\beta}_1 x_i)$$

# Bivariate Ordinary Least Squares Solution

$$\begin{aligned} S(\hat{\beta}_0, \hat{\beta}_1) &= \sum_{i=1}^N (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \\ &= \sum_{i=1}^N (y_i^2 - 2y_i \hat{\beta}_0 - 2y_i \hat{\beta}_1 x_i + \hat{\beta}_0^2 + 2\hat{\beta}_0 \hat{\beta}_1 x_i + \hat{\beta}_1^2 x_i^2) \end{aligned}$$

$$\frac{\partial S(\hat{\beta}_0, \hat{\beta}_1)}{\partial \hat{\beta}_0} = \sum_{i=1}^N (-2y_i + 2\hat{\beta}_0 + 2\hat{\beta}_1 x_i)$$

$$\frac{\partial S(\hat{\beta}_0, \hat{\beta}_1)}{\partial \hat{\beta}_1} =$$

# Bivariate Ordinary Least Squares Solution

$$\begin{aligned} S(\hat{\beta}_0, \hat{\beta}_1) &= \sum_{i=1}^N (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \\ &= \sum_{i=1}^N (y_i^2 - 2y_i \hat{\beta}_0 - 2y_i \hat{\beta}_1 x_i + \hat{\beta}_0^2 + 2\hat{\beta}_0 \hat{\beta}_1 x_i + \hat{\beta}_1^2 x_i^2) \end{aligned}$$

$$\frac{\partial S(\hat{\beta}_0, \hat{\beta}_1)}{\partial \hat{\beta}_0} = \sum_{i=1}^N (-2y_i + 2\hat{\beta}_0 + 2\hat{\beta}_1 x_i)$$

$$\frac{\partial S(\hat{\beta}_0, \hat{\beta}_1)}{\partial \hat{\beta}_1} = \sum_{i=1}^N (-2y_i x_i + 2\hat{\beta}_0 x_i + 2\hat{\beta}_1 x_i^2)$$

# Bivariate Least Squares Solution

The **first order conditions** are

$$0 = \sum_{i=1}^N (-2y_i + 2\hat{\beta}_0 + 2\hat{\beta}_1 x_i)$$

and

$$0 = \sum_{i=1}^N (-2y_i x_i + 2\hat{\beta}_0 x_i + 2\hat{\beta}_1 x_i^2)$$

# Bivariate Least Squares Solution

The **first order conditions** are

$$0 = \sum_{i=1}^N (-2y_i + 2\hat{\beta}_0 + 2\hat{\beta}_1 x_i)$$

and

$$0 = \sum_{i=1}^N (-2y_i x_i + 2\hat{\beta}_0 x_i + 2\hat{\beta}_1 x_i^2)$$

Now solving for  $\hat{\beta}_0$  and  $\hat{\beta}_1$  yields the following equations:

$$\hat{\beta}_0 N = \left( \sum_{i=1}^N y_i \right) - \hat{\beta}_1 \left( \sum_{i=1}^N x_i \right)$$

$$\hat{\beta}_1 \sum_{i=1}^N x_i^2 = \left( \sum_{i=1}^N x_i y_i \right) - \hat{\beta}_0 \left( \sum_{i=1}^N x_i \right)$$



# Bivariate Least Squares Solution

And these equations

$$\hat{\beta}_0 N = \left( \sum_{i=1}^N y_i \right) - \hat{\beta}_1 \left( \sum_{i=1}^N x_i \right)$$

$$\hat{\beta}_1 \sum_{i=1}^N x_i^2 = \left( \sum_{i=1}^N x_i y_i \right) - \hat{\beta}_0 \left( \sum_{i=1}^N x_i \right)$$

# Bivariate Least Squares Solution

And these equations

$$\begin{aligned}\hat{\beta}_0 N &= \left( \sum_{i=1}^N y_i \right) - \hat{\beta}_1 \left( \sum_{i=1}^N x_i \right) \\ \hat{\beta}_1 \sum_{i=1}^N x_i^2 &= \left( \sum_{i=1}^N x_i y_i \right) - \hat{\beta}_0 \left( \sum_{i=1}^N x_i \right)\end{aligned}$$

rearranged yield the **bivariate ordinary least squares (OLS)** estimators:

$$\begin{aligned}\hat{\beta}_0 &= \bar{y} - \hat{\beta}_1 \bar{x} \\ \hat{\beta}_1 &= \frac{\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^N (x_i - \bar{x})^2} = \frac{\text{Cov}(x, y)}{\text{Var}(x)}\end{aligned}$$

# Our Estimated Prediction Function

$$\text{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i + \hat{\epsilon}_i$$

# Our Estimated Prediction Function

$$\text{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i + \hat{\epsilon}_i$$

$$\text{Vote}_i = 50.76 + 0.16 \times \text{Approval}_i + \hat{\epsilon}_i$$


# Our Estimated Prediction Function

$$\text{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i + \hat{\epsilon}_i$$

$$\text{Vote}_i = 50.76 + 0.16 \times \text{Approval}_i + \hat{\epsilon}_i$$

Our Estimated **Prediction Function**:

$$\widehat{\text{Vote}}_i = 50.76 + 0.16 \times \text{Approval}_i$$



$$\widehat{\text{Vote}}_i = 50.76 + 0.16 \times \text{Approval}_i$$

Interpret the parameter values:

$$\widehat{\text{Vote}}_i = 50.76 + 0.16 \times \text{Approval}_i$$

Interpret the parameter values:

$$\hat{\beta}_0 = 50.76$$

$$\widehat{\text{Vote}}_i = 50.76 + 0.16 \times \text{Approval}_i$$

Interpret the parameter values:

$$\hat{\beta}_0 = 50.76$$

$$\hat{\beta}_1 = 0.16$$



# Predicting Election Result

Consider the following Gallup polling numbers:

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times \text{Approval}_{2024}$$

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times -4$$

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times \text{Approval}_{2024}$$

$$\begin{aligned}\widehat{\text{Vote}}_{2024} &= 50.76 + 0.16 \times -4 \\ &= 50.76 - 0.64 = 50.12\end{aligned}$$

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times \text{Approval}_{2024}$$

$$\begin{aligned}\widehat{\text{Vote}}_{2024} &= 50.76 + 0.16 \times -4 \\ &= 50.76 - 0.64 = 50.12\end{aligned}$$

What about the following:

- Net approval: 105%

# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times \text{Approval}_{2024}$$

$$\begin{aligned}\widehat{\text{Vote}}_{2024} &= 50.76 + 0.16 \times -4 \\ &= 50.76 - 0.64 = 50.12\end{aligned}$$

What about the following:

- Net approval: 105%
- Net approval: -50%



# Predicting Election Result

Consider the following Gallup polling numbers:

- 46% Approve, 50% Disapprove
- -4% Net Approval

$$\widehat{\text{Vote}}_{2024} = \hat{\beta}_0 + \hat{\beta}_1 \times \text{Approval}_{2024}$$

$$\widehat{\text{Vote}}_{2024} = 50.76 + 0.16 \times \text{Approval}_{2024}$$

$$\begin{aligned}\widehat{\text{Vote}}_{2024} &= 50.76 + 0.16 \times -4 \\ &= 50.76 - 0.64 = 50.12\end{aligned}$$

What about the following:

- Net approval: 105%
- Net approval: -50%

**Caution:** Function is defined for all values!