

Introduction to Multivariate Analysis

Lecture 1

August 24, 2005
Multivariate Analysis

Today's Lecture

- Introductions.
- Syllabus and course overview.
- Chapter 1 (a brief review, really):
 - Data organization/notation.
 - Graphical techniques.
 - Distance measures.
- Introduction to SAS.

Overview

• Today's Lecture

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Who are you?

To help all of use get to know each other better, please tell us your:

- Name.
- Department and specialty.
- Where you are from originally.
- Where you did you undergraduate work.

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Syllabus

Syllabus discussion...

Overview

Introductions

Course Overview

- Multivariate

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Multivariate Statistics

A taxonomy of multivariate statistical analyses shows that most techniques fall into one of the following categories:

1. Data reduction or structural simplification.
2. Sorting and grouping.
3. Investigation of the dependence among variables.
4. Prediction.
5. Hypothesis construction and testing.

Overview

Introductions

Course Overview

● Multivariate

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Data Organization

Overview

Introductions

Course Overview

Data Organization

- Arrays

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

- As a precursor of things to come, here is a preview of the ways data are organized in this book/course.
- Multivariate data are a collection of observations (or measurements) of:
 - p variables ($k = 1, \dots, p$).
 - n “items” ($j = 1, \dots, n$).
 - “items” can also be thought of as subjects/examinees/individuals or entities (when people are not under study) .
 - In some disciplines (such as educational measurement), “items” are considered the variables collected per individual.

Data Organization

- x_{jk} = measurement of the k^{th} variable on the j^{th} entity.

	Variable 1	Variable 2	...	Variable k	...	Variable p
Item 1:	x_{11}	x_{12}	...	x_{1k}	...	x_{1p}
Item 2:	x_{21}	x_{22}	...	x_{2k}	...	x_{2p}
⋮	⋮	⋮		⋮		⋮
Item j :	x_{j1}	x_{j2}	...	x_{jk}	...	x_{jp}
⋮	⋮	⋮		⋮		⋮
Item n :	x_{n1}	x_{n2}	...	x_{nk}	...	x_{np}

Arrays

- To represent the entire collection of items and entities, a rectangular array can be constructed:

$$\mathbf{X} = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1k} & \dots & x_{1p} \\ x_{21} & x_{22} & \dots & x_{2k} & \dots & x_{2p} \\ \vdots & \vdots & & \vdots & & \vdots \\ x_{j1} & x_{j2} & \dots & x_{jk} & \dots & x_{jp} \\ \vdots & \vdots & & \vdots & & \vdots \\ x_{n1} & x_{n2} & \dots & x_{nk} & \dots & x_{np} \end{bmatrix}$$

- In the next class, we will learn about how arrays like this have an algebra that makes life somewhat easier.
- All arrays will be symbolized by boldfaced font.

Overview

Introductions

Course Overview

Data Organization

• Arrays

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Array Example

Overview

Introductions

Course Overview

Data Organization

• Arrays

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

- So, putting things all together, envision standing outside of the Kansas Union Bookstore, asking people for receipts.
- You are interested in looking at two variables:
 - Variable 1: the total amount of the purchase.
 - Variable 2: the number of books purchased.
- You find four people, and here is what you see observe (with notation:

$$x_{11} = 42 \quad x_{21} = 52 \quad x_{31} = 48 \quad x_{41} = 58$$

$$x_{12} = 4 \quad x_{22} = 5 \quad x_{32} = 4 \quad x_{42} = 3$$

Array Example (Continued)

- The data array would look like:

$$\mathbf{X} = \begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \\ x_{31} & x_{32} \\ x_{41} & x_{42} \end{bmatrix} = \begin{bmatrix} 42 & 4 \\ 52 & 5 \\ 48 & 4 \\ 58 & 3 \end{bmatrix}$$

- Notice for any variable, x_{jk} :
 - The first subscript (j) represents the ROW location in the data array.
 - The second subscript (k) represents the COLUMN location in the data array.

Overview

Introductions

Course Overview

Data Organization

• Arrays

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Descriptive Statistics Review

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

- Sample Mean
- Sample Variance
- Sample Correlation

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

- When we have a large amount of data, it is often hard to get a manageable description of the nature of the variables under study.
- For this reason (and as a way of introducing a review topics from previous courses), descriptive statistics are used.
- Such descriptive statistics include:
 - Means.
 - Variances.
 - Covariances.
 - Correlations.

Sample Mean

- For the k^{th} variable, the sample mean is:

$$\bar{x}_k = \frac{1}{n} \sum_{j=1}^n x_{jk}$$

- An array of the means for all p variables then looks like this (which we will come to know as the mean vector):

$$\bar{\mathbf{x}} = \begin{bmatrix} \bar{x}_1 \\ \bar{x}_2 \\ \bar{x}_3 \\ \bar{x}_4 \end{bmatrix}$$

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

● Sample Mean

● Sample Variance

● Sample Correlation

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Sample Variance

- For the k^{th} variable, the sample variance is:

$$s_k^2 = s_{kk} = \frac{1}{n} \sum_{j=1}^n (x_{jk} - \bar{x}_k)^2$$

- Note the “kk” subscript, this will be important because the equation that produces the variance for a single variable is a derivation of the equation of the covariance for a pair of variables.
- Also note the division by n . Reasons for this will become apparent in the near future.
- For a pair of variables, i and k , the sample covariance is:

$$s_{ik} = \frac{1}{n} \sum_{j=1}^n (x_{ji} - \bar{x}_i)(x_{jk} - \bar{x}_k)$$

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

- Sample Mean
- Sample Variance
- Sample Correlation

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Sample Covariance Matrix

- Making an array of all sample covariances give us:

$$\mathbf{S}_n = \begin{bmatrix} s_{11} & s_{12} & \dots & s_{1p} \\ s_{21} & s_{22} & \dots & s_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ s_{p1} & s_{p2} & \dots & s_{pp} \end{bmatrix}$$

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

- Sample Mean
- Sample Variance
- Sample Correlation

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Sample Correlation

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

- Sample Mean
- Sample Variance
- Sample Correlation

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

- Sample covariances are dependent upon the scale of the variables under study.
- For this reason, the correlation is often used to describe the association between two variables.
- For a pair of variables, i and k , the sample correlation is found by dividing the sample covariance by the product of the standard deviation of the variables:

$$r_{ik} = \frac{s_{ik}}{\sqrt{s_{ii}}\sqrt{s_{kk}}}$$

- The sample correlation:
 - Ranges from -1 to 1.
 - Measures linear association.
 - Is invariant under linear transformations of i and k .
 - Is a biased estimator.

Sample Correlation Matrix

- Making an array of all sample covariances give us:

$$\mathbf{R} = \begin{bmatrix} 1 & r_{12} & \dots & r_{1p} \\ r_{21} & 1 & \dots & r_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ r_{p1} & r_{p2} & \dots & 1 \end{bmatrix}$$

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

- Sample Mean
- Sample Variance
- Sample Correlation

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Graphical Techniques

- Displaying multivariate data can be difficult due to our natural limitations of 3-dimensions.
- Several simple ways of displaying data include:
 - Bivariate scatterplots.
 - Three-dimensional scatterplots.
- But you already know those, some plots that can be achieved by multivariate methods include:
 - “Stars.”
 - Chernoff faces.

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques


- Bivariate Scatterplots
- Trivariate Scatterplots
- Stars
- Chernoff Faces
- Dendrograms
- Variable Space
- Network Diagrams

Distance Measures


SAS Introduction

Wrapping Up

Bivariate Scatterplots



Trivariate Scatterplots



Graphical Techniques

- But you already know those plots.
- Some plots that can be achieved by multivariate methods include:
 - “Stars.”
 - Chernoff faces.
 - Dendrograms.
 - Bivariate plots, but of the variable space.
 - Network graphs.

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

- Bivariate Scatterplots

- **Trivariate Scatterplots**

- Stars

- Chernoff Faces

- Dendrograms

- Variable Space

- Network Diagrams


Distance Measures

SAS Introduction


Wrapping Up

Stars


Arizona Public Service (1)




Boston Edison Co. (2)




Central Louisiana Electric Co. (3)




Commonwealth Edison Co. (4)



Chernoff Faces



Dendrograms



Variable Space Plots





Figure 12.18 A two-dimensional representation of universities produced by metric multidimensional scaling.

Network Diagrams



Distance Measures

- A great number of multivariate techniques revolve around the computation of distances:
 - Distances between variables.
 - Distances between entities.
- The formula for the Euclidean distance formula between the coordinate pair $P = (x_1, x_2)$ and the origin $P = (0, 0)$:

$$d(O, P) = \sqrt{x_1^2 + x_2^2}$$

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Distance Measures

- Elaborate discussions of distance measures will be found later in the class
- Just keep in mind that there are statistical analogs to distance measures, taking the variability of variables into account.
- Also be aware that there are literally an infinite number of distance measures!
- A distance measure must satisfy the following:
 - $d(P, Q) = d(Q, P)$
 - $d(P, Q) > 0$ if $P \neq Q$
 - $d(P, Q) = 0$ if $P = Q$
 - $d(P, Q) \leq d(P, R) + d(R, Q)$ (known as the triangle inequality)

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

Introduction to SAS

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

- SAS has a reputation for being...well...unliked by many in the social sciences.
- Why bother teaching it in this course?
 - New focus on SAS in our department.
 - Adding value to your degree (check out amstat.org or dice.com for details).
- For some good things about SAS, check out <http://www.pbs.org/cringely/pulpit/pulpit20020411.html>

Final Thought

- We just introduced what this course will be about.
- Things will become increasingly relevant as time progresses.
- Please be patient with SAS.
- We will now head down to the lab for a SAS introduction session.



Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

● Final Thought

● Next Class

Next Time

- Matrix algebra (Chapter 2, Supplement 2A)
- *SAS proc iml*

Overview

Introductions

Course Overview

Data Organization

Descriptive Statistics

Graphical Techniques

Distance Measures

SAS Introduction

Wrapping Up

- Final Thought
- Next Class