

The empirical recurrence for OEIS A222094, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

3 September 2026

Abstract

OEIS A222094 counts the distinct OCCUPANCY arrays obtained when every cell of an $n \times 2$ grid holds one token, each token moves to a neighbouring cell, and the direction a token leaves a cell in is restricted by the direction in which another token arrives there. The entry carries an empirical recurrence of order 7 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. What is counted is the size of an IMAGE — two different assignments of moves with the same occupancy count once — so it is not a walk count in the obvious graph; determinising the machine that emits the occupancy row by row turns it into one, on $S = 190$ states, after which the conjectured recurrence is settled by a finite exact computation.

2020 Mathematics Subject Classification. 05A15, 68Q45, 60C05.

1 The conjecture

OEIS A222094 is “Number of $2 \times n$ arrays of occupancy after each element stays put or moves to some horizontal, vertical or antidiagonal neighbor, without move-in move-out left turns.”. It has offset 1 and begins

3, 31, 306, 3014, 29671, 292017, 2873693, 28278630, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 17*a(n-1) - 90*a(n-2) + 217*a(n-3) - 258*a(n-4) + 156*a(n-5) - 44*a(n-6) + 4*a(n-7)$.

As of the “Last modified” line on the live entry (Aug 15 2018 12:37:46, revision 8) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

Every cell of an $n \times 2$ grid holds one token. Each token moves by one of the offsets

$(-1, 0), (-1, 1), (0, -1), (0, 0), (0, 1), (1, -1), (1, 0),$

a move being available only when it lands inside the grid. The offset $(0, 0)$ is among them: the entry lets a token stay put. The entry writes the array with n as the number of COLUMNS. Transposing it exchanges the two coordinates of every offset, which is done once so that the walk always runs down the rows; a transposition is a reflection and reverses the sense of a turn, so the left-hand test is negated in the transposed frame rather than being applied in the wrong one. The OCCUPANCY array records, for each cell, how many tokens land on it, and the entry

counts how many different occupancy arrays occur as the assignment of moves runs over all its possibilities.

The moves are not free of one another. Suppose the token at a cell A moves by d to the cell B , and the token standing at B moves by d' . The entry forbids the ordered pair (d, d') when

- $d \wedge d' > 0$, that is, the outgoing direction lies to the LEFT of the incoming one, the wedge being the cross product of the two offsets read as plane vectors;

and imposes nothing else. A token that stays put neither arrives anywhere nor leaves, so a zero offset on either side of the pair carries no condition at all.

The sense of “left” is fixed once and for all. Rows are numbered downwards, so the offset (d_1, d_2) is the plane vector $(d_2, -d_1)$, and $d \wedge d'$ denotes $d_2(-d'_1) - (-d_1)d'_2$. A positive value says d' is obtained from d by a rotation through an angle strictly between 0 and π counterclockwise, which is what a turn to the left is. The condition is not a 90° test: for a neighbour set containing diagonal offsets the turns of 45° and 135° are left turns too, and the entry’s data distinguishes the two readings.

That the count is the size of an IMAGE is the whole difficulty: two assignments with the same occupancy must be counted once, and a walk count over the assignments would count them twice.

3 One window decides everything

Write $c_i \in \Delta^2$ for the row of offsets chosen by the cells of row i , where Δ is the offset set above.

Lemma 1. *Both the occupancy of row i and every instance of the forbidden pair whose FIRST member is a move made in row i are determined by c_{i-1}, c_i, c_{i+1} , with the convention that an absent row is the symbol $\perp$.*

Proof. Every offset changes the row index by at most one, so a token landing on a cell of row i started in row $i - 1$, i or $i + 1$; that is the occupancy. For the pairs, a move made by the cell (i, j) with offset d lands on the cell $(i + d_1, j + d_2)$, whose row is $i - 1$, i or $i + 1$, so the offset d' of that cell is a coordinate of c_{i-1} , c_i or c_{i+1} . Reading a row of the array as $\perp$ makes “up” or “down” unavailable, which is exactly the boundary rule. $\square$

Each ordered pair is therefore tested exactly once, from the row of the token that makes the first move of the pair; every row plays that part exactly once along the array.

Let $\beta(p, c, x)$ be the occupancy row of c in the context (p, x) , undefined when some cell of c has no room to move or when some forbidden pair occurs. The occupancy array of an assignment $c_1, \dots, c_n$ is

$$(\beta(\perp, c_1, c_2), \beta(c_1, c_2, c_3), \dots, \beta(c_{n-1}, c_n, \perp)).$$

4 The image is a walk count after determinisation

Determinise. For a set D of pairs (p, c) and an occupancy row b put

$$\delta(D, b) = \{(c, x) : (p, c) \in D, \beta(p, c, x) = b\}, \quad f(D) = |\{\beta(p, c, \perp) : (p, c) \in D, \beta(p, c, \perp) \text{ defined}\}|,$$

and let $D_0 = \{(\perp, c) : c\}$. An induction on the number of rows shows D_t is exactly the set of pairs of consecutive move-rows completing a prefix with the emitted occupancy rows, so that with M the adjacency matrix of the digraph on the reachable nonempty sets, one edge per occupancy row it admits,

$$a(n) = \iota^\top M^{n-1} \tau, \quad \iota = e_{D_0}, \quad \tau = f.$$

The last row is counted by f and not by an edge: it is emitted with nothing below it, so it is a property of the state the walk stops in and τ is not the all-ones vector. There are $S = 190$ reachable sets. In particular a is C -finite.

5 The proof

Lemma 2. *Let $q(t) = t^7 - \sum_i c_i t^{7-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+7} - \sum_i c_i M^{j+7-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 3.

$$a(n) = 17a(n-1) - 90a(n-2) + 217a(n-3) - 258a(n-4) + 156a(n-5) - 44a(n-6) + 4a(n-7)$$

for every $n > 7$.

Proof. The vector $w = q(M)\tau$ was formed by 7 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 7 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

6 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 18 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest grids straight from the definition — enumerate every assignment of offsets to cells, discard the assignments containing a forbidden pair, form the occupancy array, and count the distinct arrays — with the forbidden-pair test written out from the entry’s wording rather than taken from the model. The published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A222094.
- [2] J. E. Hopcroft, R. Motwani and J. D. Ullman, *Introduction to Automata Theory, Languages, and Computation*, 3rd ed., Addison–Wesley, 2006. (The subset construction, Section 2.3.)
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.