

The empirical recurrence for OEIS A208793, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

4 September 2026

Abstract

OEIS A208793 counts arrays of width 5 over $\{0, 1, 2\}$ in which no cell is the average of its neighbours. The entry carries an empirical recurrence of order 87 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical: what a cell asks reaches one row above it and none below, so the condition on a whole row is settled by 2 consecutive rows and the count is a walk count on $S = 244$ states.

2020 Mathematics Subject Classification. 05A15, 05B20, 15A18.

1 The conjecture

OEIS A208793 is “Number of $n \times 5$ 0..2 arrays with no element equal the average of immediate neighbors vertically above, diagonally above and left, and horizontally left of it.”. It has offset 1 and begins

48, 4540, 461280, 46570746, 4706484836, 475618885520, 48061584255222, 4856847815865546, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 16*a(n-1) + 7184*a(n-2) + 184746*a(n-3) - 3131578*a(n-4) - 124401715*a(n-5) + 661252162*a(n-6) + 39523120524*a(n-7) - 118123039489*a(n-8) - 7850624206951*a(n-9) + 24857888681498*a(n-10) + 1055617081944672*a(n-11) - 4691635269200727*a(n-12) - 96201396540801379*a(n-13) + 618949424617596821*a(n-14) + 5582923300046393379*a(n-15) - 53485789201753572606*a(n-16) - 165295272048479010215*a(n-17) + 2957684753177551063480*a(n-18) - 1035336009064408740791*a(n-19) - 100981469184706605343247*a(n-20) + 28203159273917921315243321) + 1958218307267469546441483*a(n-22) - 10678991274236989231944843*a(n-23) - 15860504816102282024317857*a(n-24) + 208865772617363888022357100*a(n-25) - 12284463747752760853026) - 2498446803339436742224709601*a(n-27) + 4817916430390672239468689392*a(n-28) + 19335812204686064346788956452*a(n-29) - 64463757148412256911382507214*a(n-30) - 93954995867823933907083630896*a(n-31) + 549002790292685373898543363410*a(n-32) + 202312015860701016713941912111*a(n-33) - 3396740596860712442654476807076*a(n-34) + 884233527910524516578172350880*a(n-35) + 16219183768214788807417673238174*a(n-36) - 11086014255501737520512507408131*a(n-37) - 62028379885044189377037124735216*a(n-38) + 61435147009787366601296072459496*a(n-39) + 195164965958106958898874832790569*a(n-40) - 236747877463014164442162449333683*a(n-41) - 516048516087782719243755535520553*a(n-42) + 700316402736382947161324170310418*a(n-43) + 1166403451993100889419007657179300*a(n-44) - 1652016995665260476779805143119334*a(n-45) - 2281327061080054215718284161268520*a(n-46) + 3159515554290664910912403728855416*a(n-47) + 3883952157477963462721390622498708*a(n-48) - 4924285588757374188674992976589784*a(n-49) - 5746894817139837106268488654646280*a(n-50) + 6235373182476064497712484653451616*a(n-51) + 7326697529032251168271964204861200*a(n-$

52) -6344457848686341171551718625838112*a(n-53) -7937101028359967608486697497609920*a(n-54) +5075435069558950270245702300523776*a(n-55) +7183664823911109778092932080588992*a(n-56) -3059533897723250631258209991118016*a(n-57) -5332726309088620293109052366162560*a(n-58) +1255231074613317850053561921003264*a(n-59) +3183699846934102577570431862068736*a(n-60) -222438432500475722247461261259776*a(n-61) -1495801024256419383271165917998592*a(n-62) -110752957332074998118797184564224*a(n-63) +539067790490642302335605832910848*a(n-64) +109793470758977626308367851996160*a(n-65) -144151927853335865479105751150592*a(n-66) -47812960589426192530384101679104*a(n-67) +27226000357581931927082416361472*a(n-68) +13205108522611283355670735339520*a(n-69) -3296640606416885550525013131264*a(n-70) -2473383278429250490240357761024*a(n-71) +176384444786124457663332450304*a(n-72) +317827468317127571220149829632*a(n-73) +15523760639929701210054590464*a(n-74) -27099078076889424377305628672*a(n-75) -4061987217720465695027232768*a(n-76) +1323746119754742367299567616*a(n-77) +354440705832095156191363072*a(n-78) -20303736265320268772147200*a(n-79) -13732121335757562345160704*a(n-80) -721232319483660065570816*a(n-81) +186197813024275682033664*a(n-82) +19638823895629408763904*a(n-83) -687698477433218924544*a(n-84) -130521928539301740544*a(n-85) -1943832082828492800*a(n-86) +92288882866913280*a(n-87)

As of the “Last modified” line on the live entry (Oct 03 2025 15:01:30, revision 9) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

The arrays have 5 columns and a growing number of rows, with entries in $\{0, 1, 2\}$. Fix the list of offsets the entry names,

$$\Delta = \{(-1, -1), (-1, 0), (0, -1)\},$$

and for a cell (i, j) write

$$S(i, j) = \{(i + \delta_1, j + \delta_2) : (\delta_1, \delta_2) \in \Delta\} \cap (\text{the cells of the array})$$

for the neighbours it actually has. Only offsets landing inside the array count, so a cell on the border simply has fewer neighbours.

The condition is that no cell equals the average of the neighbours it has:

$$x(i, j) \neq \frac{1}{|S(i, j)|} \sum_{(a, b) \in S(i, j)} x(a, b)$$

whenever $S(i, j)$ is not empty. A cell with no neighbour at all is under no condition, and that is what the entry’s first published term says.

3 A window of 2 rows

The offsets reach one row up and none down, so everything the condition asks of one row is decided by that row together with one row above it and none below — a window of 2 consecutive rows, and nothing wider. Take the array one row at a time and let the state be the last row, with a distinguished start standing for the array before its first row, so that a walk of n steps is an array of n rows.

Appending a row completes exactly one window, and the row it settles is the bottom row of that window; a step is allowed when that row passes. Each row of the finished array is therefore tested exactly once.

Every state may end an array, so $\tau = \mathbf{1}$. Thus

$$a(n) = \iota^\top M^j \tau, \quad S = 244,$$

for the transfer matrix M on those S states. The walk index j and the entry's own n differ by a constant, read off by matching the published terms rather than assumed. In particular a is C -finite.

4 The proof

Lemma 1. *Let $q(t) = t^{87} - \sum_i c_i t^{87-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+87} - \sum_i c_i M^{j+87-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 2.

$$a(n) = 16a(n-1) + 7184a(n-2) + 184746a(n-3) - 3131578a(n-4) - 124401715a(n-5) + 661252162a(n-6) + 39$$

for every $n > 87$.

Proof. The vector $w = q(M)\tau$ was formed by 87 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 87 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 12 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition: every array of width 5 over $\{0, 1, 2\}$ with a few rows was written out cell by cell and each cell tested against the neighbours it has. No window, no state and no end vector are involved in that computation, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 1 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A208793.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.