

The empirical recurrence for OEIS A240413, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

3 September 2026

Abstract

OEIS A240413 counts arrays over $\{0, 1, 2, 3\}$ in which no entry may equal, modulo 4, a given constant plus the SUM of all the entries lying before it along one of several rays. The entry carries an empirical recurrence of order 15 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. None of those sums is bounded and none is local, but each is built one term at a time, so its residue modulo 4 is carried in the state; the count is then a walk count on $S = 24$ states.

2020 Mathematics Subject Classification. 05A15, 11A07, 15A18.

1 The conjecture

OEIS A240413 is “Number of $2 \times n$ 0..3 arrays with no element equal to zero plus the sum of elements to its left or one plus the sum of the elements above it or zero plus the sum of the elements diagonally to its northwest, modulo 4.”. It has offset 1 and begins

5, 10, 32, 80, 162, 493, 1109, 2656, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 3*a(n-2) + 10*a(n-3) - 3*a(n-4) - 6*a(n-5) - 3*a(n-6) + 8*a(n-7) - 7*a(n-8) - 9*a(n-9) - 4*a(n-10) + 13*a(n-11) - 14*a(n-12) + 17*a(n-13) - a(n-15)$ for $n > 16$

As of the “Last modified” line on the live entry (Jun 02 2025 09:56:02, revision 6) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

The array has 2 cells across one way, entries $x(i, j) \in \{0, 1, 2, 3\}$, and the entry names these rays out of a cell (i, j) , with the constant it attaches to each:

ray	the cells on it	constant
west	$(i, j - 1), (i, j - 2), \dots$	1
north	$(i - 1, j), (i - 2, j), \dots$	0
northwest	$(i - 1, j - 1), (i - 2, j - 2), \dots$	0

Writing $\Sigma_r(i, j)$ for the sum of the entries on ray r out of (i, j) , all of them inside the array, the condition is

$$x(i, j) \not\equiv c_r + \Sigma_r(i, j) \pmod{4}$$

for every ray r named. A ray that leaves the array at once contributes the empty sum, zero. The entry writes the array with n as the number of COLUMNS, so the slice added at each step is a column and the roles of the west and north rays are exchanged. The northwest ray is carried to itself by that exchange.

3 The rays as a bounded state

Take the array one *column* at a time, in the direction in which it grows, and index the positions of a *column* by p . Let x be the *column* just written.

Lemma 1. *Along the columns, the 3 ray sums named by this entry satisfy*

$$\begin{aligned} \Sigma_{\text{north}} & \text{ is the running total inside the current column,} \\ U'[p] &= U[p] + x[p], & \text{the west ray, per position,} \\ NW'[p] &= NW[p-1] + x[p-1], & NW'[0] = 0. \end{aligned}$$

Proof. Each ray is a set of cells at a fixed step from (i, j) repeated. Advancing one *column* moves a cell one step along the west ray, one step along the northwest ray together with one position to the right, and one step along the northeast ray together with one position to the left; in each case exactly the cell just left behind is the new term. At the border of the array the diagonal rays are empty, which is the stated boundary value. $\square$

Reduced modulo 4 these vectors take finitely many values, so they are a state, and one step of the walk writes a whole *column*: the position p is filled left to right and the running total of the north ray is carried along as it is filled, so every cell is tested against every ray at the moment it is written. The walk starts at the all-zero state, the empty array, and every state may end one, so $\tau = \mathbf{1}$ and

$$a(n) = \iota^\top M^j \tau, \quad S = 24.$$

The walk index j and the entry's own n differ by a constant, read off by matching the published terms rather than assumed. In particular a is C -finite.

4 The proof

Lemma 2. *Let $q(t) = t^{15} - \sum_i c_i t^{15-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+15} - \sum_i c_i M^{j+15-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 3.

$$a(n) = 3a(n-2) + 10a(n-3) - 3a(n-4) - 6a(n-5) - 3a(n-6) + 8a(n-7) - 7a(n-8) - 9a(n-9) - 4a(n-10) + 13a(n-11)$$

for every $n > 16$.

Proof. The vector $w = q(M) \tau$ was formed by 15 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 16 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 28 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition: fill the cells one at a time in the orientation the entry's own name writes and, for each candidate value, walk each ray to the edge of the array and add the entries up. No residue, no state and no recurrence of Lemma 1 is involved, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A240413.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.