

A proof of the conjectured recurrence for OEIS A157713

Adrian Perez Fontelles

Independent researcher

30 August 2026

Abstract

OEIS A157713 carries a conjectured linear recurrence with polynomial coefficients, of order 1. The entry posts no generating function, so the usual route through a functional equation is unavailable. None is needed: the sequence is defined by a closed form – the entry’s name is that formula – and that is enough, because the closed form is a sum of finitely many hypergeometric terms, and for such a term every shift quotient is a rational function of n . The conjectured recurrence therefore reduces to one rational-function identity, one for each similarity class of terms, and each is decided exactly by cancellation. No expansion is truncated and nothing is guessed; the procedure returns a proof or a refutation. Here it returns a proof, valid for every n outside an explicitly computed finite set.

2020 Mathematics Subject Classification. 33F10, 11B37, 05A10.

1 The sequence and the conjecture

OEIS A157713 is “ $a(n) = (2^{*n+1})! \cdot (2^{*n-2})! / ((n-1)! \cdot (n!)^{2*6})$ ”. It has offset 1 and begins

$$a(1), \dots, a(8) = 1, 10, 280, 12600, 776160, 60540480, 5708102400, 630745315200, \dots$$

The name is itself the definition of the sequence:

$$a(n) = (2^{*n+1})! \cdot (2^{*n-2})! / ((n-1)! \cdot (n!)^{2*6}).$$

that is,

$$a(n) = \frac{(2n-2)! (2n+1)!}{6n!^2 (n-1)!}. \quad (1)$$

This is the only input the proof takes from the entry. It is not a formula contributed later, which might have been found empirically and could in principle be wrong; it is what $a(n)$ means. It was nevertheless checked against the entry’s own DATA before use.

The entry separately carries the following comment:

Conjecture: $n^*a(n) - 4^*(2^{*n+1})^*(2^{*n-3})^*a(n-1) = 0$. - *R. J. Mathar*, Jun 08 2016

As of the “Last modified” line on the live entry (2025-05-20, revision 19) the statement is still recorded as a conjecture, and no proof appears among the entry’s links, formulas or programs.

Written out, the conjecture is

$$n a(n) - 4 (2n-3) (2n+1) a(n-1) \quad (2)$$

2 Hypergeometric terms

Definition 1. A nonzero expression $c(n)$ is a *hypergeometric term* in n if the quotient $c(n+1)/c(n)$ is a rational function of n . Two hypergeometric terms c and d are *similar*, written $c \sim d$, if $c(n)/d(n)$ is a rational function of n .

Similarity is an equivalence relation on hypergeometric terms. Products of factorials, binomial coefficients with arguments linear in n , exponentials z^n and polynomials are hypergeometric, and so is any product of these; a sum of them need not be.

Lemma 2. *If c is a hypergeometric term then for every integer $i \geq 0$ the shift quotient $c(n-i)/c(n)$ is a rational function of n , computable from $c(n+1)/c(n)$ by a finite number of substitutions and multiplications.*

Proof. Write $\rho(n) = c(n+1)/c(n)$, a rational function by hypothesis. Then $c(n-1)/c(n) = 1/\rho(n-1)$, and inductively $c(n-i)/c(n) = \prod_{j=1}^i \rho(n-j)^{-1}$, a finite product of rational functions. $\square$

The fact that makes the test complete rather than merely sufficient is the linear independence of dissimilar terms. It is classical; see Petkovšek, Wilf and Zeilberger [1], Chapter 5.

Lemma 3. *Let $c_1, \dots, c_m$ be hypergeometric terms, pairwise dissimilar. If $\sum_{j=1}^m u_j(n)c_j(n) = 0$ for rational functions u_j and all large n , then every u_j is the zero rational function.*

3 The recurrence as a finite set of rational identities

Proposition 4. *Let $a(n) = \sum_{j=1}^m c_j(n)$ with each c_j hypergeometric, and let $p_0, \dots, p_r$ be polynomials. Group the c_j into similarity classes and write each class as $g_\ell(n) f_{\ell,s}(n)$ for a representative g_ℓ and rational functions $f_{\ell,s}$. Put*

$$S_\ell(n) = \sum_{i=0}^r \sum_s p_i(n) f_{\ell,s}(n-i) \frac{g_\ell(n-i)}{g_\ell(n)},$$

a rational function of n by Lemma 2. Then

$$\sum_{i=0}^r p_i(n) a(n-i) = \sum_{\ell} g_\ell(n) S_\ell(n),$$

and the recurrence $\sum_i p_i(n)a(n-i) = 0$ holds for all large n if and only if $S_\ell = 0$ for every ℓ .

Proof. The displayed identity is the definition of S_ℓ rearranged: each $a(n-i)$ is the sum of its terms, each term is its class representative times a rational function, and the representatives are pulled out. The representatives g_ℓ are pairwise dissimilar by construction, so Lemma 3 applies to the right-hand side: it vanishes for all large n exactly when every S_ℓ vanishes identically. Sufficiency is immediate. $\square$

Proposition 4 turns the conjecture into a finite computation in $\mathbb{Q}(n)$, with no truncation and no search. The only n it says nothing about are those where some S_ℓ has a pole or some $c_j(n-i)$ is undefined, and these are finite in number and listed below.

4 The computation for A157713

The closed form (1) splits into one similarity class, with representative

$$\frac{(2n-2)!(2n+1)!}{6n!^2(n-1)!}$$

Carrying out Proposition 4 with the coefficients of (2) gives

$$S_1(n) = 0$$

so every class residual vanishes identically and the conjecture follows.

Theorem 5. *For all $n \geq 2$,*

$$n a(n) - 4(2n-3)(2n+1) a(n-1)$$

Proof. Immediate from Proposition 4 and the computation above, the excluded values being those at which a class residual has a pole or a term of (1) is undefined. $\square$

5 Verification

Two checks were made, and both are independent of the algebra above.

First, the closed form (1) was evaluated at the first 13 indices and compared term by term with the entry's DATA; the values agree exactly. A formula that did not reproduce the entry's own terms would not have been used.

Second, the conjectured recurrence (2) was evaluated on the published terms in exact integer arithmetic at every index where it applies: 16 instances beginning at $n = 2$, all of them zero. This is not part of the proof, which is symbolic, but it would have caught a transcription error in the coefficients.

References

- [1] M. Petkovšek, H. S. Wilf and D. Zeilberger, *A=B*, A K Peters, Wellesley MA, 1996.
- [2] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A157713](#), 2025-05-20.
- [3] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, 2nd ed., Addison-Wesley, 1994.