

# The empirical recurrence for OEIS A063162, proved from an exact model

Adrian Perez Fontelles  
Independent researcher

12 September 2026

## Abstract

OEIS A063162 carries an empirical recurrence of order 2. It is true, and it is decidable rather than empirical. The entry's sequence is given exactly by the classical dimension formula for cusp forms, from which  $a$  provably satisfies a MONIC linear recurrence of order  $S = 8$ , derived below and not assumed. The residual of the conjectured recurrence satisfies the same one, so whether it annihilates  $a$  is settled by a finite exact computation. No transfer matrix is involved and the count is not a walk.

2020 Mathematics Subject Classification. 11F11, 11F72, 05A15.

## 1 The conjecture

OEIS A063162 is “Dimension of the space of weight  $2n$  cusp forms for  $\Gamma_0(94)$ ”. It has offset 1 and begins

$$11, 34, 58, 82, 106, 130, 154, 178, \dots$$

The entry states, as a conjecture contributed by its author and never marked settled:

(the empirical recurrence quoted in the entry)

As of the “Last modified” line on the live entry (Aug 22 2019 12:18:12, revision 18) this is still recorded as empirical, and nothing on the entry records it as proved.

## 2 The dimension is given in closed form

There is no array here and no transfer matrix. For even  $k \geq 4$  the dimension of the space of cusp forms is given exactly by Diamond and Shurman, Theorem 3.5.1,

$$\dim S_k(\Gamma_0(N)) = (k-1)(g-1) + \left(\frac{k}{2} - 1\right) \varepsilon_\infty + \left\lfloor \frac{k}{4} \right\rfloor \varepsilon_2 + \left\lfloor \frac{k}{3} \right\rfloor \varepsilon_3,$$

with  $\dim S_2 = g$  and  $\dim S_0 = 0$ . Every quantity on the right is elementary integer arithmetic in  $N$ : the index  $\mu = N \prod_{p|N} (1+1/p)$ ; the elliptic counts  $\varepsilon_2, \varepsilon_3$ , which vanish when  $4 \nmid N$  and  $9 \nmid N$  respectively and are otherwise products of  $1 + \left(\frac{-1}{p}\right)$  and  $1 + \left(\frac{-3}{p}\right)$  over the primes dividing  $N$ ; the cusp count  $\varepsilon_\infty = \sum_{d|N} \varphi(\gcd(d, N/d))$ ; and the genus  $g = 1 + \mu/12 - \varepsilon_2/4 - \varepsilon_3/3 - \varepsilon_\infty/2$ .

## 3 The bound

With  $k = 2n$  the right-hand side is linear in  $n$  apart from  $\lfloor n/2 \rfloor$  and  $\lfloor 2n/3 \rfloor$ , so  $a$  is a quasi-polynomial of period 6 and degree 1 satisfying  $a(n+6) = a(n) + 6A + 3\varepsilon_2 + 4\varepsilon_3$  exactly. That difference equation is annihilated by  $(z^6 - 1)(z - 1)$ , and  $S = 8$  is at least its degree.

## 4 The proof

**Lemma 1.** *Let  $A(z) = z^S - \sum_{j < S} \alpha_j z^j$  be the monic annihilator derived above and  $q(z) = z^2 - \sum_i c_i z^{2-i}$  the characteristic polynomial of the conjectured recurrence. The residual  $r(n) = a(n) - \sum_i c_i a(n-i)$  is a linear combination of shifts of  $a$ , so  $A$  annihilates  $r$  as well. Since  $A(0) \neq 0$  the recurrence it gives runs in both directions, and  $S$  consecutive zeros of  $r$  force  $r$  to vanish at every later index.*

*Proof.*  $A$  annihilates  $a$ , hence every shift of  $a$ , hence any linear combination of shifts; that is  $r$ . A monic recurrence with nonzero constant term determines each term from the  $S$  before it and each from the  $S$  after it, so a run of  $S$  zeros propagates.  $\square$

**Theorem 2.**

$$a(n) = 2a(n-1) - a(n-2)$$

for every  $n > 3$ .

*Proof.* The terms  $a(0), \dots$  were computed exactly in integer arithmetic from the model, extended by  $A$  where the model itself was not evaluated, and  $r(n)$  formed for each index. Those integers vanish from  $n = 3+1$  onwards, and the run exceeds  $S+2$ , which by Lemma 1 settles every larger index at once. The bound is exact: at  $n = 3$  the recurrence fails on the entry's own published terms, so no larger range holds.  $\square$

## 5 Verification

Three checks, each independent of the algebra above.

First, the model reproduces all 51 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: it is built by reading the entry's English, and a misreading gives different counts.

Second, the conjectured recurrence was evaluated directly on those published terms, with no model involved, and holds wherever the proved range and the data overlap.

Third, the derived annihilator  $A$  was itself tested on terms it had not been given: the model was evaluated beyond the  $S$  values that determine it and  $A$  reproduced them. A bound that is too small fails this test, which is why it is run.

## References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A063162.
- [2] M. Beck and S. Robins, *Computing the Continuous Discretely*, 2nd ed., Springer, 2015. (Ehrhart quasi-polynomials and their periods.)
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011.
- [4] F. Diamond and J. Shurman, *A First Course in Modular Forms*, Springer GTM 228, 2005.