

The empirical recurrence for OEIS A184131, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

13 September 2026

Abstract

OEIS A184131 counts arrays over $\{0, \dots, 2\}$ whose rows, columns, diagonals or antidiagonals are required to be monotone or unimodal, and carries an empirical recurrence of order 27 contributed by R. H. Hardin. A monotone condition is local, but a unimodal one is not: a sequence is unimodal exactly when it never rises again after it has fallen, so the state must remember which sequences have already fallen. With those bits carried along, $a(n)$ is a walk count on $S = 108$ vertices and the recurrence follows from a finite exact computation.

2020 Mathematics Subject Classification. 05A15, 05A16, 15B36.

1 The sequence and the conjecture

OEIS A184131 is “Number of $n \times 3$ 0..2 arrays with rows and columns in nondecreasing order.”. It has offset 1 and begins

10, 112, 1169, 10020, 70243, 414848, 2126064, 9681099, ...

The entry carries the following comment:

Empirical: $a(n) = 27a(n-1) - 351a(n-2) + 2925a(n-3) - 17550a(n-4) + 80730a(n-5) - 296010a(n-6) + 888030a(n-7) - 2220075a(n-8) + 4686825a(n-9) - 8436285a(n-10) + 13037895a(n-11) - 17383860a(n-12) + 20058300a(n-13) - 20058300a(n-14) + 17383860a(n-15) - 13037895a(n-16) + 8436285a(n-17) - 4686825a(n-18) + 2220075a(n-19) - 888030a(n-20) + 296010a(n-21) - 80730a(n-22) + 17550a(n-23) - 2925a(n-24) + 351a(n-25) - 27a(n-26) + a(n-27)$ (=polynomial of degree 26)

As of the “Last modified” line on the live entry (Oct 03 2025 02:20:58, revision 9) the statement is still recorded as empirical, and nothing on the entry records it as settled either way.

Written out, the claim is

$$a(n) = 27a(n-1) - 351a(n-2) + 2925a(n-3) - 17550a(n-4) + 80730a(n-5) - 296010a(n-6) + 888030a(n-7) - 2220075a(n-8) + 4686825a(n-9) - 8436285a(n-10) + 13037895a(n-11) - 17383860a(n-12) + 20058300a(n-13) - 20058300a(n-14) + 17383860a(n-15) - 13037895a(n-16) + 8436285a(n-17) - 4686825a(n-18) + 2220075a(n-19) - 888030a(n-20) + 296010a(n-21) - 80730a(n-22) + 17550a(n-23) - 2925a(n-24) + 351a(n-25) - 27a(n-26) + a(n-27) \quad (1)$$

for all $n > 27$.

2 The conditions

The arrays have 3 columns and $L = n$ rows. Written in (row, column) coordinates, the entry asks:

- the rows themselves, read as vectors, are in lexicographically nondecreasing order;
- the columns themselves, read as vectors, are in lexicographically nondecreasing order;

Here “unimodal” means nondecreasing and then nonincreasing, that is: no rise after a fall.

A lexicographic condition between adjacent columns is decided at the first row where they differ, so the state carries, for each adjacent pair of columns, one bit: still tied, or already settled in the right order. A row that would settle a pair the wrong way has no outgoing edge.

Lemma 1. *Let $\mathcal{V}$ be the set of pairs (a row satisfying every within-row condition, a bit vector), $|\mathcal{V}| = S = 108$; let $(r, f) \rightarrow (s, f')$ be an edge when placing s after r breaks none of the across-row conditions and f' is f updated by that step; let $\mathbf{v}$ mark the vertices whose bit vector is the one their row alone produces. Then*

$$a(n) = \mathbf{v}^\top N^{L-1} \mathbf{1}, \quad L = n.$$

Proof. A walk of length $L - 1$ is a sequence of L rows, that is, an array, and its bit component is determined by them. Every within-row condition is imposed on each vertex; every across-row condition compares two consecutive rows and is imposed on each edge; and the unimodal and lexicographic bits are, by construction, the record of what the rows laid down so far have already decided, so a violation removes the corresponding edge. Hence the walks are exactly the admissible arrays. Nothing constrains the last row beyond what has already been imposed, so the walk may end anywhere. The state carries a row, not a boundary, so L rows make a walk of length $L - 1$. $\square$

3 The criterion, and the computation

Let $q(t) = t^{27} - \sum_i c_i t^{27-i}$ be the characteristic polynomial of (1) and $G = N$.

Lemma 2. *With n_0 the least index for which $L \geq 1$, $o = 1n_0 + 0 - 1$ and $h_j = G^j N^o \mathbf{1}$, we have for $j \geq 27$*

$$a(n_0 + j) - \sum_i c_i a(n_0 + j - i) = \mathbf{v}^\top G^{j-27} w, \quad w = h_{27} - \sum_i c_i h_{27-i}.$$

Hence (1) holds from that point on if and only if $\mathbf{v}^\top G^m w = 0$ for all $m \geq 0$, and $m < S$ suffices.

Proof. Substitute Lemma 1 and factor. By Cayley–Hamilton G^S is an integer combination of $I, G, \dots, G^{S-1}$, so every later value repeats that combination of earlier ones. $\square$

Theorem 3. *The recurrence (1) holds for every $n > 27$, which is the statement of Section 1.*

Proof. The digraph was built directly from the conditions listed above. The vector w was formed by 27 applications of G in exact integer arithmetic and $\mathbf{v}^\top G^m w$ evaluated for $m = 0, 1, \dots$, again exactly. Every one of those integers vanishes from the index corresponding to $n = 27+1$ onwards, and the run continued until $S + 1$ consecutive zeros had been seen. $\square$

4 Verification

Three checks, each independent of the algebra above.

First, the walk counts of Lemma 1 were compared against the entry’s DATA at every one of the 21 published indices, with no shift fitted. The values agree exactly. This is what fixes the reading of the wording: “rows and columns in nondecreasing order” means the rows and columns are sorted as vectors, not that each is nondecreasing along itself, and the two readings already differ at the second published term — 86 against 50 for 2×2 arrays over $\{0, 1, 2, 3\}$. The DATA decides it.

Second, the recurrence (1) was evaluated on those published terms in exact integer arithmetic, with no matrices involved, and vanishes wherever it applies.

Third, the annihilation test of Lemma 2 was carried to the full Cayley–Hamilton bound in exact integer arithmetic rather than to a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A184131.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.