

The empirical recurrence for OEIS A250837: recovering a conjecture that is not written down

Adrian Perez Fontelles
Independent researcher

2 September 2026

Abstract

OEIS A250837 records “Empirical recurrence of order 70”, with the recurrence itself in a linked file rather than in the entry. The conjecture can nevertheless be settled, and the recurrence recovered, without reading that file. The entry counts arrays under a condition local to a bounded window of lines, so the count is a walk count on $S = 243$ vertices and satisfies some monic recurrence of order at most S ; Berlekamp–Massey applied to $2S$ exact terms therefore returns the MINIMAL such recurrence. Its order is exactly 70, the order the entry states, and the same is true of every tail of the sequence. Any recurrence of order 70 that the sequence satisfies from any point on must then have a characteristic polynomial that is a multiple of the minimal one and of the same degree, hence equal to it. So the entry’s recurrence is the one displayed here, whatever its file contains, and it is proved.

2020 Mathematics Subject Classification. 05A15, 11B37, 15A18.

1 A conjecture one cannot read

OEIS A250837 is “Number of $(n+1) \times (4+1)$ 0..2 arrays with no 2×2 subblock having a diagonal absolute difference less than its antidiagonal absolute difference.”. It has offset 1 and begins

12145, 633207, 33244665, 1751284191, 92293013405, 4864186536955, 256366497513917, 13511888452284935, ...

The entry states:

Empirical recurrence of order 70 (see link above)

As of the “Last modified” line on the live entry (Jul 23 2025 12:48:03, revision 6) this is still recorded as empirical. The recurrence’s coefficients are not in the entry text.

2 The count is a walk count

The entry’s condition is local: it constrains a bounded window of consecutive lines of the array and nothing further. The admissible configurations of one such window are the vertices of a finite digraph, an edge joining u to v when v may follow u , and an admissible array is exactly a walk. Hence, with M the adjacency matrix on $S = 243$ vertices, $a(n) = \iota^\top M^{j(n)} \tau$ for a linear reindexing j fixed by the entry’s shape and checked against its published terms. By the Cayley–Hamilton theorem M^S is an integer combination of $I, M, \dots, M^{S-1}$, so:

Lemma 1. *a satisfies a monic linear recurrence with integer coefficients of order at most $S = 243$.*

3 Recovering the recurrence

Lemma 2. *A sequence known to satisfy some linear recurrence of order at most S is determined, together with its minimal recurrence, by its first $2S$ terms: Berlekamp–Massey applied to them returns that minimal recurrence exactly.*

Running it on $2S = 486$ exact terms of the model returns order 70 — the order the entry states — with the integer coefficients

$$a(n) = \sum_{i=1}^{70} c_i a(n-i),$$

c_1	171	c_{19}	−239606971695451761982	c_{37}	−334136544097810133497854736	c_{55}
c_2	−12770	c_{20}	494039923130973772692	c_{38}	−166998889442252844095305674	c_{56}
c_3	568260	c_{21}	2367823118049680664991	c_{39}	616077273221350233016092302	c_{57}
c_4	−17082911	c_{22}	−25102021030858871279629	c_{40}	−706884859189108568747968587	c_{58}
c_5	368361864	c_{23}	113075671927705249692803	c_{41}	413771062437248778627412006	c_{59}
c_6	−5814102825	c_{24}	−302376181632887946390043	c_{42}	34496881560279732282365842	c_{60}
c_7	65545032107	c_{25}	366372316711723696419225	c_{43}	−313330965283660359131013818	c_{61}
c_8	−458238722195	c_{26}	743980516614800245885814	c_{44}	231560376614896038396042427	c_{62}
c_9	288747481331	c_{27}	−5059602345998141677350606	c_{45}	45217989193469656409609892	c_{63}
c_{10}	40750135491485	c_{28}	13232681300340199373145943	c_{46}	−131511761664588312747296688	c_{64}
c_{11}	−616010519308184	c_{29}	−18398473987441755679670457	c_{47}	−94092336900333796541772203	c_{65}
c_{12}	5440977968672018	c_{30}	2587634587721003905485739	c_{48}	349799193819697408566025100	c_{66}
c_{13}	−32727115501445323	c_{31}	48391593244802113896144833	c_{49}	−383584161462376515863094612	c_{67}
c_{14}	133085727795328278	c_{32}	−107001161456218046459751479	c_{50}	264550480319063536716900346	c_{68}
c_{15}	−331695264981231982	c_{33}	84351841040496801987677871	c_{51}	−157791603108748091862845130	c_{69}
c_{16}	544606099298459810	c_{34}	102360566649242533629148309	c_{52}	90431337338644542603147424	c_{70}
c_{17}	−4136879332717280759	c_{35}	−392228493141829085492855848	c_{53}	−30758541788177974280610418	
c_{18}	44117144656891988959	c_{36}	543264190214521731899644472	c_{54}	−7922706059003486572462956	

Theorem 3. *The recurrence above holds for every n , and it is the recurrence the entry records.*

Proof. It holds by Lemma 2, which returns a recurrence the sequence satisfies from its first term. For the identification: the entry asserts that the sequence satisfies SOME recurrence of order 70, possibly only from some index on. Let q be that recurrence’s characteristic polynomial and $q_{\min}$ the minimal polynomial of the tail of the sequence. Then $q_{\min} \mid q$. Berlekamp–Massey run on the sequence with its first $70 + 4$ terms removed returns order 70 as well, so $\deg q_{\min} = 70 = \deg q$; both are monic, so $q = q_{\min}$, which is the polynomial computed above. $\square$

4 Verification

Three checks.

First, the walk model reproduces all 12 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry, and it is the only place where reading the entry’s English enters.

Second, the recovered recurrence was evaluated directly on those published terms, with no matrices and no Berlekamp–Massey involved, and holds at every index where the entry gives enough earlier terms to test it.

Third, the order was computed twice by independent routes: once modulo a 61-bit prime, where every intermediate is a machine word, and once in exact rational arithmetic; and once more on the sequence with its first $70 + 4$ terms removed, which is what the identification above requires. All three agree.

Remark 4. What is unusual here is that the conjecture’s statement was never read. The entry pins it down to a one-parameter family — the recurrences of order 70 that this sequence satisfies — and that family turns out to have exactly one member.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A250837.
- [2] J. L. Massey, Shift-register synthesis and BCH decoding, *IEEE Trans. Inform. Theory* **15** (1969), 122–127.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.