

A parity term from the level-4 cusp: a proof of Luschny's conjecture on OEIS A063321

Adrian Perez Fontelles

Independent researcher

24 August 2026

Abstract

OEIS A063321 tabulates the dimension of the space of weight- n cuspidal newforms for $\Gamma_1(48)$. In March 2012 P. Luschny conjectured the closed form $a(n) = 13(n + ((n-1) \bmod 2)) + 14(n + (n \bmod 2)) - 72$ for $n > 2$. We prove it. The route is the classical dimension formula for $S_k(\Gamma_1(M))$ together with the Atkin–Lehner–Li newform sieve; for $N = 48$ the sieve leaves only a short explicit combination, and the result is $\dim S_k(\Gamma_1(48))^{\text{new}} = \begin{cases} 27k - 31, & k \text{ even,} \\ 27k - 32, & k \text{ odd,} \end{cases}$ for $k \geq 2$. The parity term is not an accident, and its source can be named exactly: among all levels surviving the sieve, $M = 4$ is the only one for which Proposition 1 carries the correction $+\frac{1}{2}$ at odd k , and it enters with weight $\bar{\mu}(12) = -2$, contributing -1 for odd k and nothing for even k . We also record that the entry's data is offset by one against its own name: the tabulated row begins at weight 1, whose value is the sentinel -1 because weight 1 admits no dimension formula, so that $a(n)$ is the weight- $(n-1)$ dimension. Under that reading — the only one consistent with the data — the conjecture is exactly right.

1 The sequence and the conjecture

OEIS A063321 (N. J. A. Sloane, 14 July 2001) has offset 2 and begins

$-1, 23, 49, 77, 103, 131, 157, 185, 211, \dots$

The entry carries

Conjecture: $a(n) = 13(n + ((n-1) \bmod 2)) + 14(n + (n \bmod 2)) - 72$ for $n > 2$. — Peter Luschny, Mar 2012

As of the “Last modified” line on the live entry (23 August 2026) this is still recorded as an unproven conjecture, and no proof appears in the entry's links.

2 The dimension formulas

All of the following is classical; we quote it in the form given by Stein [1], Chapter 6, where it is assembled from Shimura [3, §2.6], Diamond–Shurman [2, Ch. 3] and Miyake [4, §2.5], and derived by Riemann–Roch applied to $X_1(N) \rightarrow X_1(1)$.

For a positive integer N put

$$\mu_0(N) = \prod_{p|N} (p^{v_p(N)} + p^{v_p(N)-1}), \quad \mu_1(N) = \begin{cases} \mu_0(N) & N = 1, 2, \\ \frac{1}{2}\varphi(N)\mu_0(N) & N \geq 3, \end{cases}$$

$$c_1(N) = \begin{cases} c_0(N) & N = 1, 2, \\ 3 & N = 4, \\ \sum_{d|N} \frac{1}{2}\varphi(d)\varphi(N/d) & \text{otherwise,} \end{cases} \quad g_1(N) = 1 + \frac{1}{12}\mu_1(N) - \frac{1}{4}\mu_{1,2}(N) - \frac{1}{3}\mu_{1,3}(N) - \frac{1}{2}c_1(N),$$

where $\mu_{1,2}(N) = \mu_{1,3}(N) = 0$ for $N \geq 4$. Here $g_1(N)$ is the genus of $X_1(N)$ and $c_1(N)$ is its number of cusps.

Proposition 1 ([1, Prop. 6.6]). $\dim S_2(\Gamma_1(N)) = g_1(N)$. For $k \geq 3$ and $N \geq 3$, writing $a(N, k) = (k-1)(g_1(N) - 1) + \left(\frac{k}{2} - 1\right)c_1(N)$,

$$\dim S_k(\Gamma_1(N)) = \begin{cases} a(N, k) + \frac{1}{2} & N = 4 \text{ and } k \text{ odd,} \\ a(N, k) + \lfloor k/3 \rfloor & N = 3, \\ a(N, k) & \text{otherwise.} \end{cases}$$

Proposition 2 ([1, Prop. 6.6], Atkin–Lehner–Li). $\dim S_k(\Gamma_1(N))^{\text{new}} = \sum_{M|N} \bar{\mu}(N/M) \dim S_k(\Gamma_1(M))$,

where $\bar{\mu}(R) = 0$ if $p^3 \mid R$ for some prime p , and otherwise $\bar{\mu}(R) = \prod_{p \parallel R} (-2)$, the product being over primes exactly dividing R .

Note that $\bar{\mu}(1) = 1$, $\bar{\mu}(p) = -2$, $\bar{\mu}(p^2) = 1$ and $\bar{\mu}(R) = 0$ once any prime occurs to the third power.

3 The indexing of the entry

The data of this entry comes from Stein’s table *Dimensions of the spaces* $S_k^{\text{new}}(\Gamma_1(N))$, linked from the entry. Two features of the data fix how it is to be read.

First, the initial term is -1 . A dimension cannot be negative, so this is not a dimension; it is the sentinel for an uncomputed value. Weight 1 is exactly the case for which no dimension formula is available — Stein opens Chapter 6 by saying that he gives *no simple formulas* for weight 1, and that it may not be possible to give any, since the methods behind all the formulas above fail at $k = 1$.

Second, comparing the remaining terms against Proposition 2 shows that they are the dimensions in weights $2, 3, 4, \dots$. Hence the tabulated row runs from weight 1, with weight 1 recorded as -1 , while the entry’s offset is 2. The consequence is

$$a(n) = \dim S_{n-1}(\Gamma_1(N))^{\text{new}} \quad (n \geq 3), \quad (1)$$

that is, the entry’s name is off by one against its own data. This is verified below against every published term, and against the two companion entries treated in the companion notes, which exhibit the same shift. We take (1) as the reading of the entry; it is the only one consistent with the data, and it is the reading under which the conjecture is a true statement.

4 The computation at level 48

Lemma 3. *Since $48 = 2^4 \cdot 3$, the divisors $M \mid 48$ with $\bar{\mu}(48/M) \neq 0$ are $M = 48, 24, 16, 12, 8, 4$ with weights $\bar{\mu}(1) = 1, \bar{\mu}(2) = -2, \bar{\mu}(3) = -2, \bar{\mu}(4) = 1, \bar{\mu}(6) = 4, \bar{\mu}(12) = -2$. Hence*

$$\begin{aligned} \dim S_k(\Gamma_1(48))^{\text{new}} = & \dim S_k(\Gamma_1(48)) - 2 \dim S_k(\Gamma_1(24)) - 2 \dim S_k(\Gamma_1(16)) \\ & + \dim S_k(\Gamma_1(12)) + 4 \dim S_k(\Gamma_1(8)) - 2 \dim S_k(\Gamma_1(4)). \end{aligned}$$

The invariants are $g_1 = 37, 5, 2, 0, 0, 0$ and $c_1 = 56, 24, 14, 10, 6, 3$ at levels 48, 24, 16, 12, 8, 4 respectively. Exactly one of these levels is 4, and it is the sole source of an exceptional term: by Proposition 1, $\dim S_k(\Gamma_1(4))$ carries an extra $+\frac{1}{2}$ when k is odd, which enters the combination with coefficient $\bar{\mu}(12) = -2$ and so contributes -1 at odd k .

Theorem 4. *For every $k \geq 2$,*

$$\dim S_k(\Gamma_1(48))^{\text{new}} = \begin{cases} 27k - 31, & k \text{ even}, \\ 27k - 32, & k \text{ odd}, \end{cases}$$

Proof. Substitute the values of g_1 and c_1 from Lemma 3 into Proposition 1 and form the combination of Proposition 2. Each $\dim S_k(\Gamma_1(M))$ occurring is, for $k \geq 3$, the affine function $(k-1)(g_1(M)-1) + (\frac{k}{2}-1)c_1(M)$ of k , with the single exception noted in Lemma 3; the combination of affine functions is affine, and the exceptional term contributes the stated correction. The case $k = 2$ is checked directly from $\dim S_2(\Gamma_1(M)) = g_1(M)$. $\square$

Corollary 5 (Luschny's conjecture). *For every $n > 2$,*

$$a(n) = 13(n + ((n-1) \bmod 2)) + 14(n + (n \bmod 2)) - 72 = \begin{cases} 27n - 59, & n \text{ even}, \\ 27n - 58, & n \text{ odd}, \end{cases}$$

which is Luschny's conjecture. Indeed $a(n) = \dim S_{n-1}(\Gamma_1(48))^{\text{new}}$, and n even makes $n-1$ odd, giving $27(n-1) - 32 = 27n - 59$, while n odd makes $n-1$ even, giving $27(n-1) - 31 = 27n - 58$.

Proof. Combine Theorem 4 with the reading (1) of Section 3. $\square$

Remark (why $n = 2$ is excluded, and correctly so). Under (1), $a(2)$ is the weight-1 value, which is the sentinel -1 rather than a dimension. Luschny's restriction to $n > 2$ is therefore exactly right, and could not be removed: the conjectured formula gives a different value at $n = 2$.

Remark (honest assessment). No new mathematics is involved: the dimension formulas are classical and the newform sieve is Atkin–Lehner–Li. What was missing was that nobody had carried out the substitution. The one genuinely delicate point is the indexing of Section 3; a proof addressed to the entry's name rather than its data would conclude that the conjecture is false, which is why that section is stated before any computation. The natural destination is a comment on the entry, together with a correction of the name or the offset.

5 Verification

A verification script, written separately from this note, checks every claim and prints every number quoted. It (i) implements Propositions 1 and 2 and validates them against Stein's own published tables — eight tabulated values of $\dim S_k(\Gamma_0(N))$, ten of $\dim S_k(\Gamma_1(N))$, and the two worked newform dimensions $\dim S_{12}(\Gamma_0(11))^{\text{new}} = 8$ and $\dim S_{12}(\Gamma_0(2007))^{\text{new}} = 1017$, all matching; (ii) confirms the sieve of Lemma 3; (iii) confirms Theorem 4 for $k = 2, \dots, 199$; (iv) confirms the shift (1) against all 51 published terms $n = 3, \dots, 53$, and independently against the two companion entries; and (v) confirms Luschny's formula against all 51 published terms, and against $\dim S_{n-1}(\Gamma_1(48))^{\text{new}}$ for $n = 3, \dots, 399$, with no discrepancy; and that $M = 4$ survives the sieve at $N = 48$ but not at $N = 32$ or $N = 64$, which is why those two companion entries have no parity term.

References

- [1] W. A. Stein, *Modular Forms: A Computational Approach*, Graduate Studies in Mathematics 79, AMS, 2007; Chapter 6 (Dimension Formulas).
- [2] F. Diamond and J. Shurman, *A First Course in Modular Forms*, Springer GTM 228, 2005; Chapter 3.
- [3] G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton University Press, 1994; §2.6.
- [4] T. Miyake, *Modular Forms*, Springer, 1989; §2.5.
- [5] N. J. A. Sloane et al., *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A063321>, consulted 24 August 2026.

A Suggested OEIS comment (plain text, house style)

Luschny's conjecture is true, and the parity term comes from the level-4 cusp. Since $48 = 2^4 \cdot 3$, the divisors M of 48 with $\text{mubar}(48/M)$ nonzero are $M = 48, 24, 16, 12, 8, 4$ with weights 1, -2, -2, 1, 4, -2, so

$$\begin{aligned} \dim S_k(\Gamma_1(48))^{\text{new}} &= \dim S_k(\Gamma_1(48)) - 2 \dim S_k(\Gamma_1(24)) \\ &\quad - 2 \dim S_k(\Gamma_1(16)) + \dim S_k(\Gamma_1(12)) + 4 \dim S_k(\Gamma_1(8)) \\ &\quad - 2 \dim S_k(\Gamma_1(4)). \end{aligned}$$

With $g_1 = 37, 5, 2, 0, 0, 0$ and $c_1 = 56, 24, 14, 10, 6, 3$ at levels 48, 24, 16, 12, 8, 4, the standard formula $\dim S_k(\Gamma_1(M)) = (k-1)(g_1(M)-1) + (k/2-1)c_1(M)$ applies at every level except $M = 4$, where for odd k there is an extra $+1/2$. That single term, entering with coefficient -2, is the whole source of the parity dependence:

$$\dim S_k(\Gamma_1(48))^{\text{new}} = 27k - 31 \text{ (k even), } 27k - 32 \text{ (k odd), } k \geq 2.$$

The data of this entry is the tabulated row starting at weight 1, whose value

is the sentinel -1 (weight 1 has no dimension formula), while the offset is 2. So $a(n)$ is the weight- $(n-1)$ dimension, giving $a(n) = 27n - 59$ for n even and $27n - 58$ for n odd, which is exactly the conjectured expression. The name of the entry is off by one against its data. Note that the companion entries A063305 (level 32) and A063337 (level 64) have no parity term, because $M = 4$ does not survive the sieve at those levels.