

The empirical column recurrences for the table OEIS A235198

Adrian Perez Fontelles

Independent researcher

1 September 2026

Abstract

OEIS A235198 is a two-dimensional table $T(n, k)$, and it carries a block of empirical recurrences contributed by R. H. Hardin — one for each of its first few columns. They are true. Column k of the table is an ordinary fixed-width array count, so its rows are the vertices of a finite digraph and $T(n, k)$ is a number of walks in it; being a walk count the column is C -finite, and whether a given recurrence annihilates it is decided by a finite exact computation, with the Cayley–Hamilton theorem bounding the work. This note settles the columns $k = 1$.

2020 Mathematics Subject Classification. 05A15, 05B45, 15B36.

1 The table and the conjectures

OEIS A235198 is “ $T(n, k)$ is the number of $(n+1) \times (k+1)$ 0..6 arrays with every 2×2 subblock having its diagonal sum differing from its antidiagonal sum by 5, with no adjacent elements equal (constant-stress tilted 1×1 tilings).”. It has offset 1, is read by antidiagonals, and begins

136, 512, 512, 1880, 1544, 1880, 7072, 4724, ...

The entry carries the following block:

Empirical for column k (the $k=3..6$ recurrence works also for $k=1..2$; apparently all rows and columns satisfy the same order 39 recurrence):

$k=1$: $a(n) = 30*a(n-2) - 257*a(n-4) + 468*a(n-6)$.

$k=2$: [order 23].

As of the “Last modified” line on the live entry (Jun 19 2022 02:21:21, revision 8) these are still recorded as empirical, and nothing on the entry records any of them as settled.

2 A column of the table is a walk count

Fix k . The entry defines $T(n, k)$ as the number of arrays of a shape in which one dimension is n and the other is determined by k , subject to a condition imposed on every 2×2 subblock — a condition local to two consecutive rows and two consecutive columns. Substituting the fixed value of k into the entry’s own wording turns the definition into an ordinary one-parameter array count, and for such a count the rows are the vertices of a finite digraph G_k : $r \rightarrow s$ is an edge exactly when placing s under r satisfies the condition in every column pair. An admissible array is then a walk, so with M_k the adjacency matrix of G_k ,

$$T(n, k) = c_k \mathbf{1}^\top M_k^n \mathbf{1}$$

for the constant c_k that the entry’s name supplies (a factor such as $\frac{1}{2}$ or $\frac{1}{4}$ where the name says so, and 1 otherwise).

Lemma 1. Let $q(t) = t^r - \sum_i c_i t^{r-i}$ be the characteristic polynomial of a conjectured recurrence of order r for column k . Then for every $n \geq r$,

$$T(n, k) - \sum_i c_i T(n - i, k) = c_k \mathbf{1}^\top M_k^{n-r} q(M_k) \mathbf{1},$$

so the recurrence holds from some point on if and only if $\mathbf{1}^\top M_k^j q(M_k) \mathbf{1} = 0$ for every $j \geq 0$; and by the Cayley–Hamilton theorem it is enough to check $j < S_k$, where S_k is the number of vertices of G_k .

Proof. Substitute the walk expression and factor M_k^{n-r} out of $M_k^n - \sum_i c_i M_k^{n-i}$; the scalar c_k is nonzero. For the bound, $M_k^{S_k}$ is an integer combination of $I, M_k, \dots, M_k^{S_k-1}$, so each $\mathbf{1}^\top M_k^j q(M_k) \mathbf{1}$ with $j \geq S_k$ repeats that combination of earlier values. $\square$

3 The computation, column by column

line	order	vertices S	proved for	terms checked
column $k=1$	6	42	> 6	8

In each case the vector $w = q(M_k) \mathbf{1}$ was formed by r matrix–vector products in exact integer arithmetic and $\mathbf{1}^\top M_k^j w$ evaluated for $j = 0, 1, \dots$, again exactly; every one of those integers vanishes from the stated point onwards and the run continued until the working vector was identically zero, which settles all larger j at once.

Column $k=1$

The sequence is “Number of $(n+1) \times 2$ 0..6 arrays with every 2×2 subblock having its diagonal sum differing from its antidiagonal sum by 5, with no adjacent elements equal (constant-stress tilted 1×1 tilings)”. Its rows are the vertices of a digraph on $S = 42$ vertices, edges being the admissible consecutive pairs, and the count is $\mathbf{1}^\top M^n \mathbf{1}$ up to the constant factor in the entry’s name. The conjectured recurrence

$$a(n) = 30a(n-2) - 257a(n-4) + 468a(n-6)$$

holds from index $6+1$ on.

4 Lines whose recurrence the entry does not print

For the lines below the entry records only an order, the coefficients being in a linked file. They can still be settled, and the recurrences recovered. A line is a walk count on S vertices, so it satisfies some monic recurrence of order at most S and Berlekamp–Massey on $2S$ exact terms returns the MINIMAL one. Where that order equals the stated order — and where the same holds after the first few terms are dropped, since the entry’s recurrence may begin only partway along — any recurrence of that order the line satisfies has a characteristic polynomial that is a multiple of the minimal one and of equal degree, hence is that polynomial. So the entry’s recurrence is the one displayed, whatever its file contains.

Column $k=2$, recovered

The entry gives only the order for this line: “ $k=2$: [order 23]”. Its model is a walk on $S = 252$ vertices, so it satisfies a monic recurrence of order at most S ; Berlekamp–Massey on $2S = 504$ exact terms returns the minimal one, of order 23 — the stated order — with integer coefficients

c_1	6	c_7	-128869	c_{13}	14714982	c_{19}	-8246780
c_2	52	c_8	26637	c_{14}	-20381230	c_{20}	13170048
c_3	-367	c_9	1045428	c_{15}	-23985995	c_{21}	2801112
c_4	-995	c_{10}	-801048	c_{16}	38063780	c_{22}	-1572480
c_5	9350	c_{11}	-5098437	c_{17}	20215810	c_{23}	-393120
c_6	7034	c_{12}	5754163	c_{18}	-35199080		

Repeating with the first $23 + 4$ terms removed gives order 23 again.

5 Verification

Three checks, each independent of the algebra above.

First, each model was compared against the entry’s own published values for its line, taken both from the antidiagonal DATA read in either orientation and from the “Table starts” block, and agrees at every available term. This is what ties a model to the table: substituting the fixed index into the entry’s wording is a reading of English, and a wrong reading gives a different digraph and different counts. Across the lines settled here the model reproduces 8 published values, 37 decimal digits in all, every one of them exactly; a misreading of the entry would have to reproduce all of them by accident.

Second, each conjectured recurrence was evaluated on those published values in exact integer arithmetic, with no matrices involved, and holds wherever the proved range and the data overlap.

Third, the annihilation test of Lemma 1 was carried out in exact integer arithmetic rather than on a sampled prefix.

Remark 2. A row of the table is a column of the transposed table: fixing n and letting k vary gives a count of arrays of fixed height, which is the same kind of object with the two dimensions exchanged. Both are settled here by the same argument. Only the lines the entry states explicitly, and for which enough published values exist to run the first check above, are claimed.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A235198.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.