

The empirical recurrence for OEIS A250467: recovering a conjecture that is not written down

Adrian Perez Fontelles
Independent researcher

7 September 2026

Abstract

OEIS A250467 records “Empirical recurrence of order 67”, with the recurrence itself in a linked file rather than in the entry. The conjecture can nevertheless be settled, and the recurrence recovered, without reading that file. The entry counts arrays under a condition local to a bounded window of lines, so the count is a walk count on a finite digraph and satisfies some monic recurrence of order at most the number of its states with distinct futures, here $S' = 68$; Berlekamp–Massey applied to $2S'$ exact terms therefore returns the MINIMAL such recurrence. Its order is exactly 67, the order the entry states, and the same is true of every tail of the sequence. Any recurrence of order 67 that the sequence satisfies from any point on must then have a characteristic polynomial that is a multiple of the minimal one and of the same degree, hence equal to it. So the entry’s recurrence is the one displayed here, whatever its file contains, and it is proved.

2020 Mathematics Subject Classification. 05A15, 11B37, 15A18.

1 A conjecture one cannot read

OEIS A250467 is “Number of $(n+1) \times (7+1)$ 0..1 arrays with nondecreasing $\min(x(i,j), x(i,j-1))$ in the i direction and nondecreasing $\min(x(i,j), x(i-1,j))$ in the j direction.”. It has offset 1 and begins

4757, 166765, 5286723, 169000988, 5383012776, 171499189113, 5462124226322, 173973296060000, ...

The entry states:

Empirical recurrence of order 67 (see link above)

As of the “Last modified” line on the live entry (Jul 23 2025 12:27:07, revision 6) this is still recorded as empirical. The recurrence’s coefficients are not in the entry text.

2 The count is a walk count

The entry’s condition is local: it constrains a bounded window of consecutive lines of the array and nothing further. The admissible configurations of one such window are the vertices of a finite digraph, an edge joining u to v when v may follow u , and an admissible array is exactly a walk. Hence, with M the adjacency matrix on $S = 256$ vertices, $a(n) = \iota^\top M^{j(n)} \tau$ for a linear reindexing j fixed by the entry’s shape and checked against its published terms. States with the same future contribute identically to $\iota^\top M^n \tau$, so they may be merged without changing any count; merging leaves $S' = 68$ of the 256, and it is that smaller bound the computation below uses. By the Cayley–Hamilton theorem M^S is an integer combination of $I, M, \dots, M^{S-1}$, so:

Lemma 1. *a satisfies a monic linear recurrence with integer coefficients of order at most $S' = 68$.*

3 Recovering the recurrence

Lemma 2. *A sequence known to satisfy some linear recurrence of order at most S is determined, together with its minimal recurrence, by its first $2S$ terms: Berlekamp–Massey applied to them returns that minimal recurrence exactly.*

Running it on $2S = 512$ exact terms of the model returns order 67 — the order the entry states — with the integer coefficients

$$a(n) = \sum_{i=1}^{67} c_i a(n-i),$$

c_1	9	c_{18}	2040297501360566	c_{35}	−24660411122591766774	c_{52}	13392305483595776
c_2	860	c_{19}	−2219809790388852	c_{36}	−1054840566249855608	c_{53}	13239630419865600
c_3	1128	c_{20}	−14198330546191550	c_{37}	25962978933617988912	c_{54}	−2029351351181312
c_4	−173938	c_{21}	19823133216123176	c_{38}	−1256785936152716256	c_{55}	−1729123595010048
c_5	−414920	c_{22}	73833004219003941	c_{39}	−21947916215512494896	c_{56}	225483459051520
c_6	16993857	c_{23}	−124596042232667133	c_{40}	2467492198255511712	c_{57}	168308947255296
c_7	35725323	c_{24}	−289138580911622922	c_{41}	14925463907601576256	c_{58}	−17866546544640
c_8	−982860326	c_{25}	575773677886994038	c_{42}	−2318742370830525824	c_{59}	−11800236654592
c_9	−1437896242	c_{26}	856661449759644905	c_{43}	−8165360893984628544	c_{60}	969870278656
c_{10}	36820348969	c_{27}	−2009659095857457031	c_{44}	1481749423691503872	c_{61}	567322607616
c_{11}	28999201369	c_{28}	−1919577049188649356	c_{45}	3587023991087163776	c_{62}	−33970192384
c_{12}	−939149924812	c_{29}	5399152531281015454	c_{46}	−695630317870508032	c_{63}	−17352359936
c_{13}	−174167911854	c_{30}	3222481493258002722	c_{47}	−1260394536910368768	c_{64}	692060160
c_{14}	16836140031976	c_{31}	−11323330878356223130	c_{48}	246098485535575040	c_{65}	296747008
c_{15}	−5381768492500	c_{32}	−3934369659780475081	c_{49}	352010252958166016	c_{66}	−6291456
c_{16}	−216870858071462	c_{33}	18737265000056552775	c_{50}	−66038849216618496	c_{67}	−2097152
c_{17}	159877914124844	c_{34}	3187594738909438924	c_{51}	−77424747844276224		

Theorem 3. *The recurrence above holds for every n , and it is the recurrence the entry records.*

Proof. It holds by Lemma 2, which returns a recurrence the sequence satisfies from its first term. For the identification: the entry asserts that the sequence satisfies SOME recurrence of order 67, possibly only from some index on. Let q be that recurrence’s characteristic polynomial and $q_{\min}$ the minimal polynomial of the tail of the sequence. Then $q_{\min} \mid q$. Berlekamp–Massey run on the sequence with its first $67 + 4$ terms removed returns order 67 as well, so $\deg q_{\min} = 67 = \deg q$; both are monic, so $q = q_{\min}$, which is the polynomial computed above. $\square$

4 Verification

Three checks.

First, the walk model reproduces all 14 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry, and it is the only place where reading the entry’s English enters.

Second, the recovered recurrence was evaluated directly on those published terms, with no matrices and no Berlekamp–Massey involved, and holds at every index where the entry gives enough earlier terms to test it.

Third, the order was computed twice by independent routes: once modulo a 61-bit prime, where every intermediate is a machine word, and once in exact rational arithmetic; and once more on the sequence with its first $67 + 4$ terms removed, which is what the identification above requires. All three agree.

Remark 4. What is unusual here is that the conjecture’s statement was never read. The entry pins it down to a one-parameter family — the recurrences of order 67 that this sequence satisfies — and that family turns out to have exactly one member.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A250467.
- [2] J. L. Massey, Shift-register synthesis and BCH decoding, *IEEE Trans. Inform. Theory* **15** (1969), 122–127.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.