

A condition on every subblock: the empirical recurrence for OEIS A183806

Adrian Perez Fontelles
Independent researcher

6 September 2026

Abstract

OEIS A183806 carries an empirical recurrence of order 17 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. The entry's condition is a condition on every subblock of a fixed size, so it is decided by a bounded window of consecutive lines; the admissible windows are the vertices of a finite digraph and the arrays counted are exactly the walks in it. Hence $a(n)$ is a walk count on $S = 24$ vertices and is C -finite, and whether the conjectured recurrence annihilates it is settled by a finite exact computation, with the Cayley–Hamilton theorem bounding the work. The entry does not count the arrays themselves but $1/16$ of them, so every count below is divided by 16; the division is exact at every term, which is itself a check on the model.

2020 Mathematics Subject Classification. 05A15, 05B45, 15B36.

1 The conjecture

OEIS A183806 is “ $1/16$ the number of $(n+1) \times 5$ binary arrays with no 2×2 subblock being a reflection across the shared element pair of any horizontal or vertical neighbor.”. It has offset 1 and begins

27, 237, 1953, 16953, 143191, 1232595, 10494847, 89998565, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 6a(n-1) + 81a(n-2) - 425a(n-3) - 1968a(n-4) + 10277a(n-5) + 16576a(n-6) - 92864a(n-7) - 64803a(n-8) + 320262a(n-9) + 224743a(n-10) - 99955a(n-11) - 959780a(n-12) - 1547921a(n-13) + 2262032a(n-14) + 2523060a(n-15) - 1692000a(n-16) - 1053000a(n-17)$

As of the “Last modified” line on the live entry (Oct 03 2025 02:02:52, revision 9) this is still recorded as empirical, and nothing on the entry records it as proved.

2 The count is a walk count

The entry counts arrays with 5 columns over the alphabet $\{0, 1\}$, growing one row at a time, and the condition it imposes is that no 2×2 subblock may be the mirror image of the subblock beside it or the subblock below it, reflected in the line the two share.

That condition is decided by a bounded window. A 2×2 subblock occupies 2 consecutive rows, and a condition relating one subblock to the next needs one more, so a window of 3 consecutive rows decides everything that the arrival of a new row settles; nothing outside the window is consulted. Take those windows as the vertices of a digraph, with an edge from u to v when v 's new row may follow u , and merge vertices with identical futures; that leaves $S = 24$.

An admissible array is exactly a walk, so with M the adjacency matrix and ι, τ the vectors recording which windows may begin and end an array,

$$a(n) = \frac{1}{16} \iota^\top M^n \tau,$$

the exponent fixed by the entry's own shape and checked against its published terms. In particular a is C -finite.

3 The proof

Lemma 1. *Let $q(t) = t^{17} - \sum_i c_i t^{17-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+17} - \sum_i c_i M^{j+17-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. A constant factor in front of a divides out of the residual and changes nothing here. $\square$

Theorem 2.

$$a(n) = 6a(n-1) + 81a(n-2) - 425a(n-3) - 1968a(n-4) + 10277a(n-5) + 16576a(n-6) - 92864a(n-7) - 64803a(n-8)$$

for every $n > 17$.

Proof. The vector $w = q(M)\tau$ was formed by 17 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 17 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

4 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 18 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: the digraph is built by reading the entry's English, and a misreading gives different counts.

Second, the reading was pinned before the digraph was written, by a program that enumerates every array of the given shape one by one and tests the entry's condition on each subblock directly. That program shares no code with the transfer matrix, so an error in one does not hide an error in the other.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 1 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A183806.

- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.