

The empirical recurrence for OEIS A209523, proved by counting patterns

Adrian Perez Fontelles
Independent researcher

31 August 2026

Abstract

OEIS A209523 counts arrays subject to a local condition *and* to the clause “new values 0..7 introduced in row major order”, which says the array is written in canonical form. That clause is not local, so a transfer matrix over the alphabet does not see it. It does not have to: what is being counted is not arrays but equality patterns, and the number of patterns is recovered from the numbers L_i of arrays over an i -letter alphabet by inverting a falling factorial. The result is the clean identity $8!a(n) = \sum_i \binom{8}{i} D_{8-i} L_i(n)$ with D_m the derangement numbers, and each L_i is an ordinary walk count. A second reduction — the chain is strongly lumpable over the patterns, because the condition is relabelling-invariant — brings the state space down from 8772 to $S = 98$. The entry’s empirical recurrence of order 51, contributed by R. H. Hardin, then follows from a finite exact computation.

2020 Mathematics Subject Classification. 05A15, 05A18, 15B36.

1 The sequence and the conjecture

OEIS A209523 is “Number of $n \times 4$ 0..7 arrays with every 2×2 subblock containing exactly one value repeat, and new values 0..7 introduced in row major order.”. It has offset 1 and begins

15, 700, 107693, 27323948, 9700576474, 4194606038254, 2002617518101082, 999998646599976052, ...

The entry carries the following comment:

Empirical: $a(n) = 741*a(n-1) - 80239*a(n-2) - 27206821*a(n-3) + 4305035452*a(n-4) + 223990610212*a(n-5) - 54620780269672*a(n-6) + 80102456008496*a(n-7) + 269012192646842580*a(n-8) - 5707166224378420708*a(n-9) - 614361714034268393448*a(n-10) + 20349127506784034500304*a(n-11) + 678382194937846639652480*a(n-12) - 31846630475071645146740528*a(n-13) - 302516138167984951194348896*a(n-14) + 260959278564146617831741115 - 42282326119900772602090488064*a(n-16) - 11829904605301568248908018568000*a(n-17) + 95434060058346872198828125007744*a(n-18) + 3049598030451163236051411209616896*a(n-19) - 40328192983623784101590586927442944*a(n-20) - 439535105889627937805745674742651136*a(n-21) + 8800995724627097037828678826851591680*a(n-22) + 29877440033653195430840885415475181568*a(n-23) - 1153471451160086939353282980879345710080*a(n-24) + 361417465904520256348084988960166447104*a(n-25) + 96388134984267610666941969352759061807104*a(n-26) - 237433518884853914547549341361474933817344*a(n-27) - 5316540495638084469441833849843792059400192*a(n-28) + 20393494451129478050239694172720827330625530299 - 199181629390484462577756831982634501905514496*a(n-30) - 94360773020839745817849465346613660761666331 - 5231256737767831422266603443721419521122631680*a(n-32) + 2739452469890391546206554193812788770738333 - 99803868178309860951994852593262049142125887488*a(n-34) - 52411300611543585452695952649081547330335 - 1425032472189917422814910850409521554940678373376*a(n-36) + 668627275143026130522176344822656532537 - 15273084369730432523206297036506635896548728242176*a(n-38) - 56091093092375144843369119047605323539 - 118040785197556405677917396892327617921876131577856*a(n-40) + 2986309209185552111684094780617604341 - 608824256895510126291165994808739474478152800010240*a(n-42) - 9625344197113551125978846576116137143 - 1905011080779867767627757598039656592066040429543424*a(n-44) + 1823172121497981180819938155626702343 - 3277374088192893278994603970667689592637714126077952*a(n-46) - 20927532337120974925554462377924295$

47) -2790425203670748510069753611132555534785277570580480*a(n-48) +1448099586316124905615364077335799
49) +912421745734027603434708150458948967598456307712000*a(n-50) -46246614632108395541136217892017375
51) for n>52

As of the “Last modified” line on the live entry (Oct 03 2025 15:18:30, revision 9) the statement is still recorded as empirical, and nothing on the entry records it as settled either way.

Written out, the claim is

$$a(n) = 741 a(n-1) - 80239 a(n-2) - 27206821 a(n-3) + 4305035452 a(n-4) + 223990610212 a(n-5) - 546207802 \quad (1)$$

for all $n > 52$.

2 What the canonical-form clause counts

Fix the shape: 4 columns and $L = n + 0$ rows, so $L \cdot 4$ cells. Call a partition of the cells into nonempty classes a *pattern*; an array over an alphabet determines a pattern, two cells lying in the same class exactly when they carry the same value.

Every 2×2 subblock of the array occupies two consecutive rows and two consecutive columns; writing those two rows as r and s the subblock is

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} r_j & r_{j+1} \\ s_j & s_{j+1} \end{pmatrix},$$

and the entry’s requirement on it is

$$\#\{\alpha, \beta, \gamma, \delta\} = 3. \quad (2)$$

The condition is stated purely in terms of equalities between entries, so it is invariant under any relabelling of the values and is therefore a property of the pattern alone. Call a pattern *admissible* when it has the property.

The clause “new values 0..7 introduced in row major order” selects, from each class of arrays that differ only by a relabelling, exactly one representative: reading the cells in row major order, the first occurrence of the value v must precede the first occurrence of $v + 1$. Every pattern with at most $K = 8$ classes has exactly one such representative, and every array in canonical form is the representative of its own pattern. Hence, writing $N_j(n)$ for the number of admissible patterns with exactly j classes,

$$a(n) = \sum_{j=0}^8 N_j(n). \quad (3)$$

Lemma 1. *Let $L_i(n)$ be the number of arrays over an alphabet of i letters satisfying the local condition, with no canonical-form clause. Then*

$$L_i(n) = \sum_j N_j(n) i(i-1) \cdots (i-j+1), \quad 8! \sum_{j=0}^8 N_j(n) = \sum_{i=0}^8 \binom{8}{i} D_{8-i} L_i(n),$$

where $D_m = m! \sum_{t=0}^m (-1)^t / t!$ is the number of derangements of m letters.

Proof. An array over i letters satisfying the condition determines an admissible pattern together with an injection of its classes into the alphabet, and conversely; a pattern with j classes admits $i(i-1) \cdots (i-j+1)$ injections. That is the first identity. Writing $i(i-1) \cdots (i-j+1) = j! \binom{i}{j}$ it reads $L_i = \sum_j (j! N_j) \binom{i}{j}$, so by binomial inversion $j! N_j = \sum_i (-1)^{j-i} \binom{j}{i} L_i$. Summing over $j \leq 8$ and exchanging the order of summation,

$$\sum_{j=0}^8 N_j = \sum_{i=0}^8 L_i \sum_{j=i}^8 \frac{(-1)^{j-i}}{i! (j-i)!} = \sum_{i=0}^8 \frac{L_i}{i!} \sum_{t=0}^{8-i} \frac{(-1)^t}{t!},$$

and multiplying by $8!$ turns the coefficient of L_i into $\frac{8!}{i!} \sum_{t \leq 8-i} (-1)^t / t! = \binom{8}{i} (8-i)! \sum_{t \leq 8-i} (-1)^t / t! = \binom{8}{i} D_{8-i}$, an integer. $\square$

3 Each L_i is a walk count, and the chain lumps

Fix i and let $V_i = \{0, \dots, i-1\}^4$ be the possible rows. For $r, s \in V_i$ declare $r \rightarrow s$ an edge when placing s directly after r satisfies (2) at every column, and let M_i be the adjacency matrix. An array is exactly a sequence of rows, and its condition is exactly the conjunction of the edge conditions along that sequence, so

$$L_i(n) = \mathbf{s}_i^\top M_i^{n+0-1} \mathbf{1}.$$

Because the condition involves no row before the first, every row may start a walk, and $\mathbf{s}_i = \mathbf{1}$.

Lemma 2. *Let $\pi(r)$ be the equality pattern of the row r . For all r, r' with $\pi(r) = \pi(r')$ and every pattern Q ,*

$$\#\{s : r \rightarrow s, \pi(s) = Q\} = \#\{s : r' \rightarrow s, \pi(s) = Q\}.$$

Consequently the numbers $\widehat{M}_i[P][Q]$ obtained by taking any representative of P are well defined, and with $e_P = i(i-1) \cdots (i-|P|+1)$ the number of rows of pattern P ,

$$\mathbf{s}_i^\top M_i^m \mathbf{1} = \sum_P \varepsilon_P f_m(P), \quad f_0 \equiv 1, \quad f_{m+1}(P) = \sum_Q \widehat{M}_i[P][Q] f_m(Q),$$

where $\varepsilon_P = e_P$ if rows of pattern P may start a walk and 0 otherwise.

Proof. If $\pi(r) = \pi(r')$ then the map sending the value of r at a column to the value of r' at that column is a well-defined injection between the letters used, and extends to a permutation σ of the alphabet with $r' = \sigma(r)$. The edge condition is a statement about equalities among entries, so $r \rightarrow s$ if and only if $\sigma(r) \rightarrow \sigma(s)$; and $\pi(\sigma(s)) = \pi(s)$. Hence $s \mapsto \sigma(s)$ is a bijection between the two successor sets preserving patterns, which is the first claim. It says the chain is strongly lumpable over π , so $(M_i^m \mathbf{1})(r)$ depends on r only through $\pi(r)$; calling that value $f_m(\pi(r))$ gives the recursion, and summing over r in each class gives the displayed identity, the number of rows of pattern P being the number of injections of its $|P|$ classes into the alphabet. $\square$

Assembling the 8 lumped chains into one block-diagonal matrix N of size $S = 98$ and putting the weights of Lemma 1 into the start vector, $\mathbf{v} = ((\binom{8}{i} D_{8-i} \varepsilon_P)_{i,P})$, gives

$$8! a(n) = \mathbf{v}^\top N^{n+0-1} \mathbf{1}. \quad (4)$$

The unlumped alphabet had 8772 rows in all; the lumped chain has $S = 98$ states.

4 The criterion, and the computation

Let $q(t) = t^{51} - \sum_i c_i t^{51-i}$ be the characteristic polynomial of (1).

Lemma 3. *For every n with $m = n + 0 - 1 \geq 51$,*

$$8! \left(a(n) - \sum_i c_i a(n-i) \right) = \mathbf{v}^\top N^{m-51} q(N) \mathbf{1}.$$

Hence (1) holds from that point on if and only if $\mathbf{v}^\top N^j q(N) \mathbf{1} = 0$ for every $j \geq 0$, and it is enough to check $j = 0, 1, \dots, S-1$.

Proof. Substitute (4) into the left side and factor N^{m-51} out of $N^m - \sum_i c_i N^{m-i}$. The scalar $8!$ is nonzero, so it does not affect whether the left side vanishes. For the last clause put $w = q(N)\mathbf{1}$: by Cayley–Hamilton N^S is an integer combination of $I, N, \dots, N^{S-1}$, so each $\mathbf{v}^\top N^j w$ with $j \geq S$ is the corresponding combination of earlier values. $\square$

Theorem 4. *The recurrence (1) holds for every $n > 52$, which is the statement of Section 1.*

Proof. The lumped transition counts were obtained by enumerating, for one representative row of each pattern, all admissible successors and sorting them by pattern — the successors being generated column by column, since the edge condition is a conjunction of one constraint per column. The vector $w = q(N)\mathbf{1}$ was then formed by 51 matrix–vector products in exact integer arithmetic and $\mathbf{v}^\top N^j w$ evaluated for $j = 0, 1, \dots$, again exactly. Every one of those integers vanishes from the index corresponding to $n = 52 + 1$ onwards, and the run continued until w was the zero vector, which settles all larger j at once. Lemma 3 gives the theorem. $\square$

5 Verification

Three checks, each independent of the algebra above.

First, the whole construction was compared against the entry’s DATA: the right-hand side of (4), divided by $8!$, was evaluated at every one of the 12 published indices and agrees exactly. This is a strong test of the modelling and not only of the arithmetic — the inversion of Lemma 1, the reading of the local condition, and the alignment of the entry’s index with the number of rows would each show up as a mismatch, and the quotient by $8!$ came out an integer at every index.

Second, the lumped chain was checked against the unlumped one: for the small shapes the walk counts $\mathbf{s}_i^\top M_i^m \mathbf{1}$ were computed directly over the full alphabet as well and agree with $\sum_P \varepsilon_P f_m(P)$ term by term.

Third, the recurrence (1) was evaluated on the published terms in exact integer arithmetic with no matrices involved, and the annihilation test of Lemma 3 was run until the working vector was identically zero rather than to a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A209523.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7; Stirling and falling-factorial inversion, Section 1.9.)
- [3] J. G. Kemeny and J. L. Snell, *Finite Markov Chains*, Springer, 1976. (Strong lumpability, Section 6.3.)
- [4] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.