

A disproof of the conjectured recurrence for OEIS A129366, and the recurrence that holds instead

Adrian Perez Fontelles
Independent researcher

30 August 2026

Abstract

OEIS A129366 carries a conjectured linear recurrence with polynomial coefficients, of order 6. It is false. The entry's own generating function gives, exactly, the quantity the recurrence claims to annihilate, and that quantity is not eventually zero: the smallest failure is at $n = 6$, where the left-hand side equals -73968 rather than 0, and the failure recurs for infinitely many n . We then give a recurrence of order 6 that does hold, derived from the same generating function and proved by the same criterion that refutes the conjectured one.

2020 Mathematics Subject Classification. 05A15, 11B37, 33F10.

1 The sequence and the conjecture

OEIS A129366 is “ $a(n) = \text{Sum}_{k=0..floor(n/2)} A000108(n-k)$ ”. It has offset 0 and begins

$$a(0), \dots, a(7) = 1, 1, 3, 7, 21, 61, 193, 617, \dots$$

The entry states, not as a conjecture,

G.f.: $(1/(1-x)) * (c(x) - x * c(x^2))$, where $c(x)$ is the g.f. of A000108(n).

that is,

$$A(x) = \sum_{n \geq 0} a(n)x^n = \frac{\sqrt{1-4x} - \sqrt{1-4x^2}}{2x(x-1)}. \quad (1)$$

This is the only input taken from the entry, and it was checked against every term of the entry's DATA section before use — all 25 of them — not against a sample.

The entry separately carries the following comment:

Conjecture: $n*(12*n+35)*(n-1)*a(n) + (n-1)*(12*n^2-701*n+1236)*a(n-1) + 2*(6*n^3-385*n^2+2285*n-3432)*a(n-2) + 4*(-405*n^3+5313*n^2-19970*n+23175)*a(n-3) + 8*(156*n^3-1724*n^2+5498*n-5175)*a(n-4) + 16*(393*n^3-4981*n^2+20393*n-26820)*a(n-5) - 32*(n-5)*(93*n-268)*(2*n-9)*a(n-6) = 0$. - R. J. Mathar, Feb 05 2015

As of the “Last modified” line on the live entry (2024-02-01, revision 13) the statement is still recorded as a conjecture, and nothing on the entry records it as settled either way.

Written out, the conjecture asserts

$$n(n-1)(12n+35)a(n) + (n-1)(12n^2-701n+1236)a(n-1) + 2(6n^3-385n^2+2285n-3432)a(n-2) - 4(2) \quad (2)$$

2 The residual, and what it decides

Write $\theta = x \frac{d}{dx}$, so that $\theta x^n = n x^n$, and set

$$B(x) = \sum_{i=0}^6 x^i (p_i(\theta + i)A)(x). \quad (3)$$

Lemma 1. $[x^n]B(x) = \sum_i p_i(n) a(n-i)$ for every n . Consequently the recurrence (2) holds for all $n > d$ if and only if B is a polynomial of degree at most d ; and if B is not a polynomial at all, (2) fails for infinitely many n .

Proof. θ acts diagonally on powers, so $(p_i(\theta + i)A)(x) = \sum_m p_i(m+i)a(m)x^m$, and multiplying by x^i shifts the exponent, giving $\sum_m p_i(m+i)a(m)x^{m+i} = \sum_n p_i(n)a(n-i)x^n$. Summing over i gives the first claim. The rest is immediate: a power series is a polynomial of degree at most d exactly when all its coefficients beyond x^d vanish, and a series with infinitely many nonzero coefficients is not a polynomial. $\square$

Lemma 1 is the same criterion used to *prove* such conjectures. Its force here is that it is an equivalence, so a negative answer is a disproof rather than a failure to find a proof — and the answer is computed exactly, in the field $\mathbb{Q}(x)[y]/(P(x,y))$ where A lives, with no series truncated.

3 The conjecture is false

Carrying out (3) for A129366 and reducing in $\mathbb{Q}(x)[y]/(P(x,y))$, the residual B is *not* a polynomial. By Lemma 1 the conjectured recurrence therefore fails for infinitely many n .

The failure is not confined to small indices, and it is not an artefact of the first few terms: expanding (1) and evaluating the left side of (2) directly gives

n	$\sum_i p_i(n)a(n-i)$
6	−73968
7	−271952
8	−1554560
9	−6925344
10	−33471568
11	−150229936
12	−678365952
13	−2982213920

Theorem 2. The recurrence (2) conjectured on OEIS A129366 is false. Its smallest failure is at $n = 6$, where the left-hand side equals −73968, and it fails for infinitely many n .

Proof. The tabulated values are computed from (1), whose coefficients agree with the entry's DATA at every published index, so each is a value of $\sum_i p_i(n)a(n-i)$ for the sequence A129366 itself; the entry at $n = 6$ is nonzero. That B is not a polynomial, established exactly above, gives infinitely many such n by Lemma 1. $\square$

Remark 3. The failure is not a matter of how the recurrence is indexed. Shifting the argument of the coefficients and the indices of the terms together, by any amount between −5 and 5, does not make it hold on the published data either.

4 A recurrence that does hold

The generating function (1) is algebraic over $\mathbb{Q}(x)$, so the sequence is P -recursive and a correct recurrence exists; it is obtained from (1) rather than guessed. It satisfies a linear differential equation with polynomial coefficients, and reading off the coefficient of each power of x turns that equation into the recurrence below.

Theorem 4. *For all $n \geq 6$,*

$$(-n^2 - n) a(n) + (7n^2 - 5n) a(n-1) + (-14n^2 + 46n - 36) a(n-2) + (108 - 48n) a(n-3) + (56n^2 - 328n + 504) a(n-4) = 0$$

Proof. The residual (3) formed with these coefficients reduces, in $\mathbb{Q}(x)[y]/(P(x, y))$, to 0 identically. By Lemma 1 the recurrence therefore holds at every n for which it is stated, that is for $n \geq 6$. $\square$

5 Verification

Three checks, each independent of the algebra above.

First, the generating function (1) was expanded and compared against the entry's DATA at every one of the 25 published indices; the values agree exactly. A generating function that reproduced only some of them would not have been used.

Second, the conjectured recurrence was evaluated on those published terms in exact integer arithmetic, with no generating function involved: it is nonzero at 8 of the 8 indices tabulated above.

Third, the recurrence of the previous section was evaluated the same way, on the published terms, and vanishes at all 19 indices where it applies.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A129366.
- [2] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999.
- [4] M. Petkovšek, H. S. Wilf and D. Zeilberger, *A=B*, A K Peters, 1996.