

The empirical recurrence for OEIS A223505, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

3 September 2026

Abstract

OEIS A223505 counts the ways of writing the vertices of a fixed graph G into an array of width 2 so that any two cells adjacent in one of the named directions carry vertices joined by an edge of G ; here G is graph named in the entry. The entry carries an empirical recurrence of order 3 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical: every condition joins cells inside one slice of the array or inside two consecutive slices, so a single slice is a state and the count is a walk count on a finite digraph with $S = 32$ vertices.

2020 Mathematics Subject Classification. 05C15, 05A15, 15A18.

1 The conjecture

OEIS A223505 is “Petersen graph (3,1) coloring a rectangular array: number of $2 \times n$ arrays where 0..5 label nodes of a graph with edges 0,1 0,3 3,5 3,4 1,2 1,4 4,5 2,0 2,5 and every array movement to a horizontal, diagonal or antidiagonal neighbor moves along an edge of this graph, with the array starting at 0.”. It has offset 1 and begins

6, 19, 115, 631, 3539, 19759, 110427, 617015, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 5 \cdot a(n-1) + 4 \cdot a(n-2) - 4 \cdot a(n-3)$ for $n > 4$.

As of the “Last modified” line on the live entry (Aug 21 2018 05:54:43, revision 8) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

Let G be the graph on the label set $\{0, 1, \dots, 5\}$ whose vertices are the 6 nodes $0, \dots, 5$ with the edges the entry lists, namely $\{0,1\}$, $\{0,2\}$, $\{0,3\}$, $\{1,2\}$, $\{1,4\}$, $\{2,5\}$, $\{3,4\}$, $\{3,5\}$, $\{4,5\}$.

An array x is a map from the cells of a rectangle to the vertices of G . The entry names the directions antidiagonal, diagonal and horizontal, that is the cell offsets

$$(1, -1), (1, 1), (0, 1),$$

and asks that whenever two cells of the array differ by one of those offsets, the two vertices they carry are joined by an edge of G :

$x(i, j) x(i + \delta_1, j + \delta_2) \in E(G)$ for every named offset (δ_1, δ_2) and every pair of cells in range.

The entry also fixes the top-left cell: $x(0, 0) = 0$.

Lemma 1. *The count does not depend on how the vertices are labelled.*

Proof. A relabelling is a bijection σ of the label set; if it is an isomorphism from one labelled copy of G to another then $x \mapsto \sigma \circ x$ is a bijection between the arrays counted for the two copies, because the condition mentions only the edge relation. Here the labelling is written out in the entry itself, so nothing is left to choose. $\square$

3 One slice is a state

Take the array one *column* at a time, in the direction in which it grows. The entry writes the array with n as the number of COLUMNS, so the slice added at each step is a column. Transposing the array exchanges the horizontal and vertical directions and carries each diagonal direction to itself, and that is how the directions are read below. Every offset the entry names has its two cells either in the same *column* or in two consecutive ones. So a *column*, a word of length 2 over the vertices of G , carries everything the next *column* needs: the conditions inside a *column* constrain that *column* alone, and the conditions across *columns* are a relation between the *column* just written and the one about to be written.

Let M be the matrix indexed by the legal *columns*, with $M_{uv} = 1$ when v may follow u . Add one further state for the empty array, whose successors are the legal first *columns*—those beginning with the vertex 0; then a walk of n steps from that state is exactly an array of n *columns*. Every state may end an array, so $\tau = \mathbf{1}$ and

$$a(n) = \iota^\top M^j \tau, \quad S = 32.$$

The *columns* themselves are built one cell at a time with the conditions applied as soon as they can be, which for the sparser graphs is what makes the state list short: a *column* of length 2 over 6 vertices is one of 6^2 words, but only $S - 1$ of them are legal. The walk index j and the entry's own n differ by a constant, read off by matching the published terms rather than assumed. In particular a is C -finite.

4 The proof

Lemma 2. *Let $q(t) = t^3 - \sum_i c_i t^{3-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+3} - \sum_i c_i M^{j+3-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 3.

$$a(n) = 5a(n-1) + 4a(n-2) - 4a(n-3)$$

for every $n > 4$.

Proof. The vector $w = q(M)\tau$ was formed by 3 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 4 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 20 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition, with the graph rebuilt by a different route—the cube from its corner points and face normals in space, the icosahedron as a pentagonal antiprism capped at both poles—and the arrays written out cell by cell, each cell tested against the cells it must be joined to. No slice, no state and no recurrence is involved in that computation, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A223505.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] H. S. M. Coxeter, *Regular Polytopes*, 3rd ed., Dover, 1973.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.