

The empirical product formula for OEIS A140915, proved

Adrian Perez Fontelles

Independent researcher

6 September 2026

Abstract

OEIS A140915 counts 6×8 matrices over $\{0, \dots, n\}$ whose rows and columns are non-decreasing, and carries a product formula for $a(n)$ marked empirical. Reversing rows and columns turns such a matrix into a plane partition inside a $6 \times 8 \times n$ box, so MacMahon's box theorem applies; the triple product telescopes in its third index and leaves a polynomial in n of degree 48. The entry's own expression, cleared of its denominator, becomes an identity between two polynomials of bounded degree, and is therefore settled by a finite exact computation. The box count itself is classical and is already recorded on this entry; what is proved here is the entry's own formula.

2020 Mathematics Subject Classification. 05A15, 05A17, 05A19.

1 The conjecture

OEIS A140915 is “Number of 6×8 matrices with elements in $0..n$ with each row and each column in nondecreasing order. $6,8,n$ can be permuted, see formula.”. It has offset 0 and begins

1, 3003, 2147145, 614083470, 90474964580, 7997986868872, ...

The entry states, as conjectures contributed by its author and never marked settled:

(empirical) set p,q,r to $n,8,6$ (in any order) in $s=p+q+r-1$; $a(n) = \text{product } i \text{ in } 0..r-1$
 $(\text{binomial}(s,p+i)*i!/(s-i)^{(r-i-1)})$

As of the “Last modified” line on the live entry (Jan 19 2025 11:09:13, revision 8) these are still recorded as empirical, and nothing on the entry records them as proved.

2 What is being counted

A matrix x of shape 6×8 with entries in $\{0, \dots, n\}$, nondecreasing along every row and down every column, becomes, on reversing the order of the rows and of the columns, an array that is NONINCREASING along rows and down columns with the same entries. That is exactly a plane partition whose parts are at most n and whose shape fits inside a 6×8 rectangle — a plane partition in a $6 \times 8 \times n$ box. The correspondence is a bijection, being an involution on arrays, so

$$a(n) = \text{PP}(6, 8, n).$$

Note which parameter grows: the shape is FIXED and it is the alphabet $0..n$ that n counts, so nothing about transfer matrices applies here.

3 The box count is a polynomial

Theorem 1 (MacMahon).

$$\text{PP}(p, q, r) = \prod_{i=1}^p \prod_{j=1}^q \prod_{k=1}^r \frac{i+j+k-1}{i+j+k-2}.$$

This is classical; see Stanley [2], and it is already recorded on this entry.

Theorem 2.

$$a(n) = \prod_{i=1}^6 \prod_{j=1}^8 \frac{n+i+j-1}{i+j-1},$$

a polynomial in n of degree 48 with rational coefficients.

Proof. In Theorem 1 the inner product over k telescopes: for fixed i and j ,

$$\prod_{k=1}^n \frac{i+j+k-1}{i+j+k-2} = \frac{n+i+j-1}{i+j-1},$$

every other factor cancelling against its neighbour. The remaining double product has $6 \cdot 8 = 48$ factors, each linear in n , and the denominators are constants. $\square$

So a is a polynomial, and no appeal to Ehrhart theory or to any general position argument is needed to see it.

4 The entry's expression

The entry writes $s = p + q + r - 1$ with $(p, q, r) = (n, 8, 6)$ and claims

$$a(n) = \prod_{i=0}^{6-1} \binom{s}{n+i} \frac{i!}{(s-i)^{6-1-i}} = \frac{N(n)}{D(n)}.$$

Lemma 3. *N and D are polynomials in n of degrees 63 and 15 respectively.*

Proof. With $s = n + 8 + 6 - 1$ one has $s - (n + i) = 8 + 6 - 1 - i$, a constant, so

$$\binom{s}{n+i} = \binom{s}{8+6-1-i} = \frac{(s-8-6+2+i)(s-8-6+3+i) \cdots s}{(8+6-1-i)!},$$

a polynomial in n of degree $8 + 6 - 1 - i$. Multiplying over $i = 0, \dots, 6 - 1$ gives $\deg N = \sum_{i=0}^{6-1} (8 + 6 - 1 - i) = 63$. The denominator is $\prod_{i=0}^{6-1} (s - i)^{6-1-i}$, of degree $\sum_{i=0}^{6-1} (6 - 1 - i) = 15$, and it does not vanish for $n \geq 0$. $\square$

Theorem 4. *The entry's expression equals $a(n)$ for every $n \geq 0$.*

Proof. By Theorem 2 and Lemma 3, $a(n)D(n) - N(n)$ is a polynomial in n of degree at most $48 + 15 = 63$. It was evaluated in exact rational arithmetic at $n = 0, 1, \dots, 63$, that is at 64 points, and vanishes at all of them; a nonzero polynomial of degree at most 63 has at most 63 roots, so it is identically zero. Since $D(n) \neq 0$ for $n \geq 0$, the two expressions agree there. $\square$

5 Verification

Three checks, each independent of the algebra above.

First, Theorem 2 reproduces all 12 terms the entry publishes, exactly, in integer arithmetic.

Second, the matrices themselves were counted for the smallest alphabets, directly from the definition: the nondecreasing rows were listed and the chains of 6 of them increasing componentwise counted, with no product formula and no plane partitions anywhere. The counts agree, which is what ties the reversal bijection to the entry's own wording.

Third, the polynomial identity was evaluated over the whole range $n = 0, \dots, 63$ rather than sampled, in exact rational arithmetic.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A140915.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999. (Plane partitions with bounded part size, Section 7.21.)
- [3] P. A. MacMahon, *Combinatory Analysis*, Volume 2, Cambridge University Press, 1916.
- [4] G. E. Andrews, *The Theory of Partitions*, Addison–Wesley, 1976.