

A condition on each element and its neighbours: OEIS A296324

Adrian Perez Fontelles

Independent researcher

6 September 2026

Abstract

OEIS A296324 carries an empirical recurrence of order 40 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. The entry's condition names only cells one step apart, so it is decided by a window of three consecutive lines; the admissible windows are the vertices of a finite digraph and the arrays counted are exactly the walks in it. Hence $a(n)$ is a walk count on $S = 56$ vertices and is C -finite, and whether the conjectured recurrence annihilates it is settled by a finite exact computation, with the Cayley–Hamilton theorem bounding the work.

2020 Mathematics Subject Classification. 05A15, 05B45, 15B36.

1 The conjecture

OEIS A296324 is “Number of $n \times 5$ 0..1 arrays with each 1 horizontally, vertically or antidiagonally adjacent to 2 neighboring 1's.”. It has offset 1 and begins

1, 15, 54, 185, 853, 3569, 14691, 62193, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 6*a(n-1) - 15*a(n-2) + 51*a(n-3) - 121*a(n-4) + 266*a(n-5) - 598*a(n-6) + 1026*a(n-7) - 1958*a(n-8) + 2883*a(n-9) - 4421*a(n-10) + 5480*a(n-11) - 5762*a(n-12) + 4485*a(n-13) - 1246*a(n-14) - 1423*a(n-15) + 3587*a(n-16) - 2563*a(n-17) + 1429*a(n-18) + 788*a(n-19) - 1333*a(n-20) + 2661*a(n-21) - 3283*a(n-22) + 2331*a(n-23) + 1215*a(n-24) - 4008*a(n-25) + 2878*a(n-26) + 724*a(n-27) - 2739*a(n-28) + 1555*a(n-29) + 468*a(n-30) - 1080*a(n-31) + 418*a(n-32) + 204*a(n-33) - 249*a(n-34) + 54*a(n-35) + 44*a(n-36) - 29*a(n-37) + a(n-38) + 4*a(n-39) - a(n-40)$.

As of the “Last modified” line on the live entry (Sep 05 2022 20:23:32, revision 6) this is still recorded as empirical, and nothing on the entry records it as proved.

2 The count is a walk count

The entry counts arrays with 5 columns over the alphabet $\{0, 1\}$, growing one row at a time. Its condition is that every entry equal to 1 must have exactly 2 of its horizontal and vertical and antidiagonal neighbours equal to 1.

Every cell named there lies at most one step away, so three consecutive rows decide the condition on the middle one completely. Carry two consecutive rows as the state; when a new row arrives, the row it follows has all its neighbours present and is tested then, and the row with nothing after it is tested by the end vector. That is also where the boundary is honest: a cell in the first row has nothing before it and a cell in the last has nothing after it, and a missing neighbour is not a neighbour rather than a wrapped-round one.

Vertices with identical futures are merged, which leaves $S = 56$. An admissible array is exactly a walk, so with M the adjacency matrix and ι, τ the vectors recording which states may begin and end an array,

$$a(n) = \iota^\top M^{n-1} \tau,$$

the exponent fixed by the entry's own shape and checked against its published terms. In particular a is C -finite.

3 The proof

Lemma 1. *Let $q(t) = t^{40} - \sum_i c_i t^{40-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+40} - \sum_i c_i M^{j+40-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. A constant factor in front of a divides out of the residual and changes nothing here. $\square$

Theorem 2.

$$a(n) = 6a(n-1) - 15a(n-2) + 51a(n-3) - 121a(n-4) + 266a(n-5) - 598a(n-6) + 1026a(n-7) - 1958a(n-8) + \dots$$

for every $n > 40$.

Proof. The vector $w = q(M)\tau$ was formed by 40 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 40 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

4 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 23 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: the digraph is built by reading the entry's English, and a misreading gives different counts.

Second, the reading was pinned before the digraph was written, by a program that enumerates every array of the given shape one by one and tests the entry's condition at each cell directly. That program shares no code with the transfer matrix, so an error in one does not hide an error in the other.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 1 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A296324.

- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.