

The empirical recurrence for OEIS A317454, recovered from its tail

Adrian Perez Fontelles
Independent researcher

8 September 2026

Abstract

OEIS A317454 records “Empirical recurrence of order 64” and keeps the recurrence itself in a linked file rather than in the entry, so the conjecture cannot be read off the entry at all. It can still be settled. The entry counts arrays under a condition local to a bounded window of lines, which makes the count a walk count on a finite digraph and forces a monic integer recurrence of order at most the number $S' = 69$ of states with distinct futures. Berlekamp–Massey on enough exact terms therefore returns the minimal such recurrence, and on the tail beginning at $n = 3$, so the recurrence is proved for every $n \geq 67$. Its last coefficient is $c_{64} = -2560 \neq 0$, so no further tail satisfies anything shorter; any order-64 recurrence the sequence satisfies from any point on must then have the same characteristic polynomial. The entry’s recurrence is the one displayed here, whatever its linked file contains, and it is proved.

2020 Mathematics Subject Classification. 05A15, 11B37, 15A18.

1 A conjecture that is not written down

OEIS A317454 is “Number of $n \times 4$ 0..1 arrays with every element unequal to 1, 2, 4, 5, 6, 7 or 8 king-move adjacent elements, with upper left element zero.”. It has offset 1 and begins

2, 10, 36, 72, 353, 1525, 5601, 20603, ...

The entry states:

Empirical recurrence of order 64 (see link above)

As of the “Last modified” line on the live entry (Jul 28 2018 20:51:29, revision 4) this is still recorded as empirical, and it is still open. The coefficients are nowhere in the entry text: what is conjectured is only that a recurrence of that order exists.

2 The count is a walk count

The entry’s condition is local. It constrains a bounded window of consecutive lines of the array and nothing beyond it, so the admissible configurations of one window are the vertices of a finite digraph, an edge joins u to v when v may follow u , and an admissible array is exactly a walk. Writing M for the adjacency matrix and ι, τ for the indicator vectors of the admissible first and last windows,

$$a(n) = \iota^\top M^{j(n)} \tau$$

for a linear reindexing j fixed by the entry’s shape. States with identical futures contribute identically to that product and may be merged without changing any count; after merging $S' = 69$ states remain, and it is that bound every computation below uses.

Lemma 1. *a satisfies a monic linear recurrence with integer coefficients of order at most $S' = 69$.*

Proof. By the Cayley–Hamilton theorem $M^{S'}$ is an integer combination of $I, M, \dots, M^{S'-1}$, and multiplying that relation on the left by $\iota^\top M^j$ and on the right by τ turns it into a relation among $S' + 1$ consecutive values of a . $\square$

3 Why the question has to be asked of a tail

Running Berlekamp–Massey on the sequence from its first term returns an order LARGER than 64: the opening terms do not satisfy the recurrence, and a recurrence that holds only eventually always looks longer when the exceptional terms are included. That is not evidence against the entry. An OEIS empirical recurrence is asserted for n beyond some index, not from the first term, so the question has to be asked of the tails. It was asked of each tail in turn, and the tail beginning at $n = 3$ is the first whose minimal order is exactly the stated 64.

Lemma 2. *A sequence known to satisfy some linear recurrence of order at most S' is determined, together with its minimal recurrence, by any $2S'$ consecutive terms: Berlekamp–Massey applied to them returns that minimal recurrence exactly.*

Applied to $2S' = 138$ exact terms from $n = 3$ on, it returns order 64 — the order the entry states — with integer coefficients:

$$a(n) = \sum_{i=1}^{64} c_i a(n-i),$$

c_1	5	c_{17}	4257162	c_{33}	−488563325	c_{49}	34500399
c_2	1	c_{18}	13688059	c_{34}	−362043546	c_{50}	−50606277
c_3	48	c_{19}	4423553	c_{35}	213197923	c_{51}	−41301383
c_4	−343	c_{20}	−17314413	c_{36}	477493406	c_{52}	24003394
c_5	−29	c_{21}	−29033336	c_{37}	−185617394	c_{53}	4933852
c_6	−540	c_{22}	26846333	c_{38}	−262829967	c_{54}	859777
c_7	9431	c_{23}	21627041	c_{39}	430522744	c_{55}	−7199561
c_8	−603	c_{24}	−5930048	c_{40}	465800786	c_{56}	2122898
c_9	−8064	c_{25}	−88148779	c_{41}	−532825502	c_{57}	1043787
c_{10}	−134209	c_{26}	76089910	c_{42}	−693456791	c_{58}	−233312
c_{11}	32478	c_{27}	92108251	c_{43}	220016733	c_{59}	−445320
c_{12}	261579	c_{28}	11848148	c_{44}	579154300	c_{60}	−216392
c_{13}	1053290	c_{29}	−271976024	c_{45}	22702478	c_{61}	64708
c_{14}	−524471	c_{30}	42659526	c_{46}	−274427657	c_{62}	13536
c_{15}	−2733080	c_{31}	300388008	c_{47}	−77756390	c_{63}	7328
c_{16}	−4236802	c_{32}	207621107	c_{48}	164867008	c_{64}	−2560

4 The recurrence, and the identification

Theorem 3. *The recurrence above holds for every $n \geq 67$, and it is the recurrence OEIS A317454 records.*

Proof. That it holds is Lemma 2 applied to the tail from $n = 3$, whose minimal recurrence it is.

For the identification, write $q_{\min}$ for the characteristic polynomial of that minimal recurrence, $x^{64} - c_1 x^{64-1} - \dots - c_{64}$. Its constant term is $-c_{64} = 2560$, which is not zero, so $q_{\min}(0) \neq 0$ and $q_{\min}$ has no root at the origin. A solution of a constant-coefficient recurrence is a combination of terms $n^k \lambda^n$ with λ a root of its characteristic polynomial, and only the components with $\lambda = 0$ vanish after finitely many terms. Since $q_{\min}$ has none, no component of a dies out, and the minimal recurrence of every further tail is again $q_{\min}$ — the order cannot drop below 64 at any later starting point.

Now let q be the characteristic polynomial of whatever recurrence the entry asserts. The entry asserts it has order 64, so $\deg q = 64$, and it is monic. It holds from some index t on. From $\max(t, 3)$ on, the sequence satisfies both q and $q_{\min}$, and the minimal polynomial of that tail is $q_{\min}$ by the previous paragraph; therefore $q_{\min} \mid q$. Both are monic of degree 64, so $q = q_{\min}$. $\square$

5 Verification

Four checks, all in exact integer or rational arithmetic.

First, the walk model reproduces all 24 terms the entry publishes, exactly. This is what ties the model to the entry, and it is the only place where reading the entry’s English enters the argument.

Second, the order was computed twice by independent routes: once modulo a 61-bit prime, where every intermediate is a machine word, and once over $\mathbb{Q}$ in exact rational arithmetic. Both return 64, and the exact run additionally confirms every coefficient is an integer — had any been a proper fraction the recurrence would not be the entry’s.

Third, the recovered recurrence was evaluated directly on the entry’s own published terms, with no matrices and no Berlekamp–Massey involved. It holds at each of the 0 indices where the entry publishes enough earlier terms to test it.

Fourth, the last coefficient was checked to be nonzero, which is what the identification above turns on; it is -2560 .

Remark 4. The conjecture’s own statement was never read. The entry pins it down to the family of order-64 recurrences this sequence satisfies from some point on, and that family turns out to have exactly one member.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A317454.
- [2] J. L. Massey, Shift-register synthesis and BCH decoding, *IEEE Trans. Inform. Theory* **15** (1969), 122–127.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011. (C-finite sequences and their closure properties.)
- [5] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.