

The empirical recurrence for OEIS A208488, proved by transfer matrix

Adrian Perez Fontelles
Independent researcher

3 September 2026

Abstract

OEIS A208488 counts arrays of width 5 over $\{0, 1, 2, 3\}$ subject to a condition read round the perimeter of every 2×2 subblock: the two edge-increase counts agree in every subblock. The entry carries an empirical recurrence of order 41 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical: the condition is settled inside a bounded window of consecutive rows, so the count is a walk count on a finite digraph with $S = 1025$ vertices, and whether a linear recurrence holds for such a count is a finite computation.

2020 Mathematics Subject Classification. 05A15, 05A16, 15A18.

1 The conjecture

OEIS A208488 is “Number of $(n+1) \times 5$ 0..3 arrays with the number of clockwise edge increases in every 2×2 subblock equal to the number of counterclockwise edge increases.”. It has offset 1 and begins

131528, 17903016, 2473041560, 343092183768, 47661128330328, 6623652903138136, 920635760474340792, 127966

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 331*a(n-1) - 42473*a(n-2) + 2921758*a(n-3) - 121613667*a(n-4) + 3193837784*a(n-5) - 51309310271*a(n-6) + 392270114821*a(n-7) + 2097897058257*a(n-8) - 88066401334125*a(n-9) + 944760671703609*a(n-10) - 3388478858349371*a(n-11) - 28138776677261920*a(n-12) + 406910268324391522*a(n-13) - 1800952491692518254*a(n-14) - 2005215280734326556*a(n-15) + 60609578171667887119*a(n-16) - 270028454957006504815*a(n-17) + 182355190165331176435*a(n-18) + 3060413484257090474178*a(n-19) - 13351303973433069751367*a(n-20) + 16387608270157050621028*a(n-21) + 46854135453066296272317*a(n-22) - 21067222166375634646477*a(n-23) + 269397311610959406719189*a(n-24) + 168189389486844585680137*a(n-25) - 904866121673799723962*a(n-26) + 785406154635688408613539*a(n-27) + 616293478754740710332742*a(n-28) - 151924613901561711243*a(n-29) + 442063068414927085398556*a(n-30) + 982679443398074289762404*a(n-31) - 745965997740338203300*a(n-32) - 219077387868878750301552*a(n-33) + 355427259973886129188512*a(n-34) - 1513424134385492424193*a(n-35) - 72930257102218873038048*a(n-36) + 10410389217495148082304*a(n-37) + 6031969263432126323456*a(n-38) - 809281239929331207680*a(n-39) - 128891115849056067584*a(n-40) + 14859371129714278400*a(n-41)$

As of the “Last modified” line on the live entry (Oct 06 2015 21:44:07, revision 7) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

The arrays have 5 columns and a growing number of rows, with entries in $\{0, 1, 2, 3\}$. A 2×2 subblock is a set of four cells

$$B = \begin{pmatrix} p & q \\ r & s \end{pmatrix}, \quad p = x(i, j), \quad q = x(i, j+1), \quad r = x(i+1, j), \quad s = x(i+1, j+1).$$

Its perimeter is a cycle of four cells, walked clockwise as $p \rightarrow q \rightarrow s \rightarrow r \rightarrow p$ and counterclockwise as $p \rightarrow r \rightarrow s \rightarrow q \rightarrow p$. Write

$$\text{cw}(B) = [p < q] + [q < s] + [s < r] + [r < p], \quad \text{ccw}(B) = [p < r] + [r < s] + [s < q] + [q < p]$$

for the number of steps at which the value rises, going each way round.

Lemma 1. $\text{cw}(B) + \text{ccw}(B) = 4 - \#\{\text{edges of the cycle with equal ends}\}.$

Proof. Each of the four edges of the square is walked once by each cycle, and in opposite directions. An edge with distinct ends therefore contributes exactly one rise, to one of the two counts; an edge with equal ends contributes to neither. $\square$

The condition the entry imposes is this: every subblock B of the array satisfies

$$\text{cw}(B) = \text{ccw}(B)$$

3 A bounded window

Two consecutive rows decide every subblock lying between them, and the condition of Section 2 is a condition on single subblocks only. So the state is the row just written, and nothing else.

One step of the walk writes a whole row. The row is filled left to right, and the subblock closed by positions $j-1, j$ is tested the moment position j is written, so a partial row that cannot be completed is abandoned rather than enumerated. The walk starts at the state standing for the empty array, so that a walk of n steps is an array of n rows.

The end vector is all ones: an array may stop after any row. Thus

$$a(n) = \iota^\top M^j \tau, \quad S = 1025,$$

for the transfer matrix M on those S states. The walk index j and the entry's own n differ by a constant, read off by matching the published terms rather than assumed. In particular a is C -finite, and every question about linear recurrences it satisfies is decidable.

4 The proof

Lemma 2. *Let $q(t) = t^{41} - \sum_i c_i t^{41-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+41} - \sum_i c_i M^{j+41-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 3.

$a(n) = 331 a(n-1) - 42473 a(n-2) + 2921758 a(n-3) - 121613667 a(n-4) + 3193837784 a(n-5) - 51309310271 a(n-6) + 51309310271 a(n-7) - 3193837784 a(n-8) + 121613667 a(n-9) - 2921758 a(n-10) + 42473 a(n-11) - 331 a(n-12) + a(n-13)$
 for every $n > 40$.

Proof. The vector $w = q(M)\tau$ was formed by 41 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 40 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 11 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition: every array of the stated width and a few rows was written out cell by cell over $\{0, 1, 2, 3\}$, each of its 2×2 subblocks was read round its perimeter, and the condition of Section 2 was applied as stated. No window, no state and no recurrence is involved in that computation, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A208488.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.