

The empirical recurrence for OEIS A253376: recovering a conjecture that is not written down

Adrian Perez Fontelles
Independent researcher

7 September 2026

Abstract

OEIS A253376 records “Empirical recurrence of order 79”, with the recurrence itself in a linked file rather than in the entry. The conjecture can nevertheless be settled, and the recurrence recovered, without reading that file. The entry counts arrays under a condition local to a bounded window of lines, so the count is a walk count on a finite digraph and satisfies some monic recurrence of order at most the number of its states with distinct futures, here $S' = 450$; Berlekamp–Massey applied to $2S'$ exact terms therefore returns the MINIMAL such recurrence. Its order is exactly 79, the order the entry states, and the same is true of every tail of the sequence. Any recurrence of order 79 that the sequence satisfies from any point on must then have a characteristic polynomial that is a multiple of the minimal one and of the same degree, hence equal to it. So the entry’s recurrence is the one displayed here, whatever its file contains, and it is proved.

2020 Mathematics Subject Classification. 05A15, 11B37, 15A18.

1 A conjecture one cannot read

OEIS A253376 is “Number of $(2+1) \times (n+1)$ 0..2 arrays with every 2×2 subblock sum nondecreasing horizontally, vertically and antidiagonally ne-to-sw.”. It has offset 1 and begins

450, 3111, 22877, 148389, 945748, 5125790, 27587386, 130018165, ...

The entry states:

Empirical recurrence of order 79 (see link above).

As of the “Last modified” line on the live entry (Sep 15 2023 12:02:46, revision 11) this is still recorded as empirical. The recurrence’s coefficients are not in the entry text.

2 The count is a walk count

The entry’s condition is local: it constrains a bounded window of consecutive lines of the array and nothing further. The admissible configurations of one such window are the vertices of a finite digraph, an edge joining u to v when v may follow u , and an admissible array is exactly a walk. Hence, with M the adjacency matrix on $S = 450$ vertices, $a(n) = \iota^\top M^{j(n)} \tau$ for a linear reindexing j fixed by the entry’s shape and checked against its published terms. By the Cayley–Hamilton theorem M^S is an integer combination of $I, M, \dots, M^{S-1}$, so:

Lemma 1. *a satisfies a monic linear recurrence with integer coefficients of order at most $S' = 450$.*

3 Recovering the recurrence

Lemma 2. *A sequence known to satisfy some linear recurrence of order at most S is determined, together with its minimal recurrence, by its first $2S$ terms: Berlekamp–Massey applied to them returns that minimal recurrence exactly.*

Running it on $2S = 900$ exact terms of the model returns order 79 — the order the entry states — with the integer coefficients

$$a(n) = \sum_{i=1}^{79} c_i a(n-i),$$

c_1	9	c_{21}	4801619959530	c_{41}	−6718378143067818744	c_{61}	−1901041788623505408
c_2	65	c_{22}	−59186272390110	c_{42}	−3990279397586265900	c_{62}	17168181780647301120
c_3	−813	c_{23}	−8367164615610	c_{43}	13573200380588028348	c_{63}	8895860007639920640
c_4	−1503	c_{24}	338018085156660	c_{44}	5003881995007001088	c_{64}	−986878655697051648
c_5	35067	c_{25}	−63026938123020	c_{45}	−24158694316978174512	c_{65}	−327682401236877312
c_6	−2049	c_{26}	−1669263148180740	c_{46}	−4356492449072602896	c_{66}	4685347765788917760
c_7	−960243	c_{27}	783202925756340	c_{47}	37885311670133837328	c_{67}	868947223463608320
c_8	1086240	c_{28}	7168284121132980	c_{48}	702180134840233920	c_{68}	−180789434872528896
c_9	18711228	c_{29}	−5196566728384740	c_{49}	−52281979345965347328	c_{69}	−120651665681940480
c_{10}	−35736696	c_{30}	−26866295989193748	c_{50}	6495657021495425856	c_{70}	553134494117265408
c_{11}	−275381736	c_{31}	26010783052803972	c_{51}	63334702238132746176	c_{71}	−17459153936842752
c_{12}	714658500	c_{32}	88048679140737015	c_{52}	−16079014545805946880	c_{72}	−129178265777602560
c_{13}	3165994812	c_{33}	−106748031772881279	c_{53}	−67090134345597242112	c_{73}	14777776041099264
c_{14}	−10469468628	c_{34}	−252329474348288919	c_{54}	25187434137260649728	c_{74}	21637807214690304
c_{15}	−28924095228	c_{35}	371635847674498251	c_{55}	61801025531016991488	c_{75}	−4007539482034176
c_{16}	120561637986	c_{36}	630909828966841993	c_{56}	−30523610020098282496	c_{76}	−2315214083063808
c_{17}	210319415742	c_{37}	−1116759425621881869	c_{57}	−49124893907374282752	c_{77}	570630428688384
c_{18}	−1132860637250	c_{38}	−1368846121844868905	c_{58}	30216827194676075520	c_{78}	118881339310080
c_{19}	−1190202257382	c_{39}	2925275798772753429	c_{59}	33334146513723036672	c_{79}	−35664401793024
c_{20}	8890114025950	c_{40}	2549493296941549716	c_{60}	−24903026047945359360		

Theorem 3. *The recurrence above holds for every n , and it is the recurrence the entry records.*

Proof. It holds by Lemma 2, which returns a recurrence the sequence satisfies from its first term. For the identification: the entry asserts that the sequence satisfies SOME recurrence of order 79, possibly only from some index on. Let q be that recurrence’s characteristic polynomial and $q_{\min}$ the minimal polynomial of the tail of the sequence. Then $q_{\min} \mid q$. Berlekamp–Massey run on the sequence with its first $79 + 4$ terms removed returns order 79 as well, so $\deg q_{\min} = 79 = \deg q$; both are monic, so $q = q_{\min}$, which is the polynomial computed above. $\square$

4 Verification

Three checks.

First, the walk model reproduces all 18 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry, and it is the only place where reading the entry’s English enters.

Second, the recovered recurrence was evaluated directly on those published terms, with no matrices and no Berlekamp–Massey involved, and holds at every index where the entry gives enough earlier terms to test it.

Third, the order was computed twice by independent routes: once modulo a 61-bit prime, where every intermediate is a machine word, and once in exact rational arithmetic; and once more on the sequence with its first $79 + 4$ terms removed, which is what the identification above requires. All three agree.

Remark 4. What is unusual here is that the conjecture’s statement was never read. The entry pins it down to a one-parameter family — the recurrences of order 79 that this sequence satisfies — and that family turns out to have exactly one member.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A253376.
- [2] J. L. Massey, Shift-register synthesis and BCH decoding, *IEEE Trans. Inform. Theory* **15** (1969), 122–127.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.