

The empirical recurrence for OEIS A317149, recovered from its tail

Adrian Perez Fontelles

Independent researcher

8 September 2026

Abstract

OEIS A317149 records “Empirical recurrence of order 69” and keeps the recurrence itself in a linked file rather than in the entry, so the conjecture cannot be read off the entry at all. It can still be settled. The entry counts arrays under a condition local to a bounded window of lines, which makes the count a walk count on a finite digraph and forces a monic integer recurrence of order at most the number $S' = 71$ of states with distinct futures. Berlekamp–Massey on enough exact terms therefore returns the minimal such recurrence, and on the tail beginning at $n = 2$, so the recurrence is proved for every $n \geq 71$. Its last coefficient is $c_{69} = 384 \neq 0$, so no further tail satisfies anything shorter; any order-69 recurrence the sequence satisfies from any point on must then have the same characteristic polynomial. The entry’s recurrence is the one displayed here, whatever its linked file contains, and it is proved.

2020 Mathematics Subject Classification. 05A15, 11B37, 15A18.

1 A conjecture that is not written down

OEIS A317149 is “Number of $n \times 4$ 0..1 arrays with every element unequal to 0, 1, 2, 4, 5, 7 or 8 king-move adjacent elements, with upper left element zero.”. It has offset 1 and begins

8, 28, 102, 416, 2313, 10682, 49746, 244887, ...

The entry states:

Empirical recurrence of order 69 (see link above)

As of the “Last modified” line on the live entry (Jul 22 2018 15:45:39, revision 4) this is still recorded as empirical, and it is still open. The coefficients are nowhere in the entry text: what is conjectured is only that a recurrence of that order exists.

2 The count is a walk count

The entry’s condition is local. It constrains a bounded window of consecutive lines of the array and nothing beyond it, so the admissible configurations of one window are the vertices of a finite digraph, an edge joins u to v when v may follow u , and an admissible array is exactly a walk. Writing M for the adjacency matrix and ι, τ for the indicator vectors of the admissible first and last windows,

$$a(n) = \iota^\top M^{j(n)} \tau$$

for a linear reindexing j fixed by the entry’s shape. States with identical futures contribute identically to that product and may be merged without changing any count; after merging $S' = 71$ states remain, and it is that bound every computation below uses.

Lemma 1. *a satisfies a monic linear recurrence with integer coefficients of order at most $S' = 71$.*

Proof. By the Cayley–Hamilton theorem $M^{S'}$ is an integer combination of $I, M, \dots, M^{S'-1}$, and multiplying that relation on the left by $\iota^\top M^j$ and on the right by τ turns it into a relation among $S' + 1$ consecutive values of a . $\square$

3 Why the question has to be asked of a tail

Running Berlekamp–Massey on the sequence from its first term returns an order LARGER than 69: the opening terms do not satisfy the recurrence, and a recurrence that holds only eventually always looks longer when the exceptional terms are included. That is not evidence against the entry. An OEIS empirical recurrence is asserted for n beyond some index, not from the first term, so the question has to be asked of the tails. It was asked of each tail in turn, and the tail beginning at $n = 2$ is the first whose minimal order is exactly the stated 69.

Lemma 2. *A sequence known to satisfy some linear recurrence of order at most S' is determined, together with its minimal recurrence, by any $2S'$ consecutive terms: Berlekamp–Massey applied to them returns that minimal recurrence exactly.*

Applied to $2S' = 142$ exact terms from $n = 2$ on, it returns order 69 — the order the entry states — with integer coefficients:

$$a(n) = \sum_{i=1}^{69} c_i a(n-i),$$

c_1	5	c_{19}	78099628	c_{37}	4013876754	c_{55}	8270200
c_2	7	c_{20}	−31428776	c_{38}	6410440819	c_{56}	−4989851
c_3	39	c_{21}	−205274890	c_{39}	1916691051	c_{57}	1202800
c_4	−394	c_{22}	−230910852	c_{40}	−4701840304	c_{58}	3257224
c_5	−445	c_{23}	96445880	c_{41}	−5642565086	c_{59}	−423020
c_6	225	c_{24}	489820040	c_{42}	−605105161	c_{60}	−2028193
c_7	13064	c_{25}	485148015	c_{43}	3833388137	c_{61}	−579972
c_8	10269	c_{26}	−30740035	c_{44}	3263588017	c_{62}	317048
c_9	−34888	c_{27}	−629993633	c_{45}	−338132487	c_{63}	453796
c_{10}	−237993	c_{28}	−844914326	c_{46}	−2100196405	c_{64}	171420
c_{11}	−97009	c_{29}	−697430850	c_{47}	−1205057370	c_{65}	6864
c_{12}	761344	c_{30}	104897063	c_{48}	412761491	c_{66}	−13440
c_{13}	2590929	c_{31}	1480337685	c_{49}	759185943	c_{67}	−3760
c_{14}	165818	c_{32}	2525370082	c_{50}	222913585	c_{68}	1280
c_{15}	−8254274	c_{33}	1131835717	c_{51}	−190911265	c_{69}	384
c_{16}	−17667578	c_{34}	−2612425915	c_{52}	−150233248		
c_{17}	3802882	c_{35}	−4933540381	c_{53}	11672165		
c_{18}	52429996	c_{36}	−2172608628	c_{54}	53200572		

4 The recurrence, and the identification

Theorem 3. *The recurrence above holds for every $n \geq 71$, and it is the recurrence OEIS A317149 records.*

Proof. That it holds is Lemma 2 applied to the tail from $n = 2$, whose minimal recurrence it is.

For the identification, write $q_{\min}$ for the characteristic polynomial of that minimal recurrence, $x^{69} - c_1 x^{68} - \dots - c_{69}$. Its constant term is $-c_{69} = -384$, which is not zero, so $q_{\min}(0) \neq 0$ and $q_{\min}$ has no root at the origin. A solution of a constant-coefficient recurrence is a combination of terms $n^k \lambda^n$ with λ a root of its characteristic polynomial, and only the components with $\lambda = 0$ vanish after finitely many terms. Since $q_{\min}$ has none, no component of a dies out, and the

minimal recurrence of every further tail is again $q_{\min}$ — the order cannot drop below 69 at any later starting point.

Now let q be the characteristic polynomial of whatever recurrence the entry asserts. The entry asserts it has order 69, so $\deg q = 69$, and it is monic. It holds from some index t on. From $\max(t, 2)$ on, the sequence satisfies both q and $q_{\min}$, and the minimal polynomial of that tail is $q_{\min}$ by the previous paragraph; therefore $q_{\min} \mid q$. Both are monic of degree 69, so $q = q_{\min}$. $\square$

5 Verification

Four checks, all in exact integer or rational arithmetic.

First, the walk model reproduces all 22 terms the entry publishes, exactly. This is what ties the model to the entry, and it is the only place where reading the entry’s English enters the argument.

Second, the order was computed twice by independent routes: once modulo a 61-bit prime, where every intermediate is a machine word, and once over $\mathbb{Q}$ in exact rational arithmetic. Both return 69, and the exact run additionally confirms every coefficient is an integer — had any been a proper fraction the recurrence would not be the entry’s.

Third, the recovered recurrence was evaluated directly on the entry’s own published terms, with no matrices and no Berlekamp–Massey involved. It holds at each of the 0 indices where the entry publishes enough earlier terms to test it.

Fourth, the last coefficient was checked to be nonzero, which is what the identification above turns on; it is 384.

Remark 4. The conjecture’s own statement was never read. The entry pins it down to the family of order-69 recurrences this sequence satisfies from some point on, and that family turns out to have exactly one member.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A317149.
- [2] J. L. Massey, Shift-register synthesis and BCH decoding, *IEEE Trans. Inform. Theory* **15** (1969), 122–127.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011. (C-finite sequences and their closure properties.)
- [5] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.