

The reduction of OEIS A396099 modulo 2

Adrian Perez Fontelles

Independent researcher

31 August 2026

Abstract

OEIS A396099 is defined by a functional equation for its ordinary generating function and carries a congruence conjecture of P. D. Hanna. It is true. The equation determines the coefficients one at a time, with $a(N)$ entering with coefficient 1, so it determines them modulo 2 as well; and modulo 2 the series $x/(1-x)$ satisfies the equation exactly, because its compositional iterates are $x/(1-jx)$ and the relevant product of denominators collapses. Uniqueness then identifies the reduction, which is the conjecture. The whole argument is an identity between rational functions over $\mathbb{Z}/2\mathbb{Z}$; nothing is truncated or estimated.

2020 Mathematics Subject Classification. 11B50, 05A15, 13F25.

1 The sequence and the conjecture

OEIS A396099 is “G.f. satisfies $A(x) = x + A^2(x) \cdot A^3(x)$.” It has offset 1 and begins

$$1, 1, 5, 39, 383, 4357, \dots$$

Writing $A(x) = \sum_{n \geq 1} a(n)x^n$ and A^j for the j -th compositional iterate of A , the entry defines A by

$$A(x) = x + A^2(x) A^3(x). \quad (1)$$

This is the only input taken from the entry.

The entry separately carries the following comment:

Conjecture: all terms are odd. — *Paul D. Hanna*

As of the “Last modified” line on the live entry (Jun 04 2026 11:37:17, revision 17) it is still recorded as a conjecture, and nothing on the entry records it as settled either way. The entry carries two further conjectures of the same contributor, on the residues modulo 4; those are stronger than what is proved here and are not claimed. Section 4 says exactly where the argument stops short of them.

2 The equation determines the sequence

Lemma 1. *Let $A(x) = \sum_{n \geq 1} a(n)x^n$ with $a(1) = 1$. Then for every $j \geq 1$ and every $N \geq 1$,*

$$[x^N]A^j = j a(N) + (a \text{ polynomial in } a(1), \dots, a(N-1)).$$

Proof. Induction on j . For $j = 1$ the statement is $[x^N]A = a(N)$. Suppose it holds for j , and write $B = A^j$, so $[x^1]B = 1$ and $[x^N]B = j a(N) + \dots$. Then $A^{j+1} = A \circ B = \sum_{m \geq 1} a(m)B^m$ and

$$[x^N]A^{j+1} = \sum_{m=1}^N a(m) [x^N]B^m.$$

The term $m = N$ contributes $a(N)([x^1]B)^N = a(N)$. The term $m = 1$ contributes $[x^N]B = j a(N) + \dots$. For $2 \leq m \leq N-1$, every monomial of $[x^N]B^m$ is a product of $m \geq 2$ coefficients of B with indices summing to N , hence each index is at most $N-1$, so no $a(N)$ occurs. Adding up gives $(j+1)a(N)$ plus terms in $a(1), \dots, a(N-1)$. $\square$

Lemma 2. *There is exactly one sequence of integers $a(1), a(2), \dots$ satisfying (1), and for each $N \geq 2$ the equation read at order x^N has the form $a(N) = (a \text{ polynomial with integer coefficients in } a(1), \dots, a(N-1))$.*

Proof. Comparing coefficients of x in (1) gives $a(1) = 1$. Let $N \geq 2$ and collect the coefficient of $a(N)$ on each side of (1) at order x^N , using Lemma 1. On the left the coefficient is 1. On the right, x contributes nothing and the product $A^2 A^3$ contributes nothing either: every monomial of $[x^N](A^2 A^3)$ is a product of one coefficient of A^2 and one of A^3 with indices $i, j \geq 1$ and $i+j = N$, so both indices are at most $N-1$. Hence the net coefficient of $a(N)$ is 1. Since that coefficient is 1, the order- x^N equation solves for $a(N)$ over $\mathbb{Z}$, and induction gives existence and uniqueness. $\square$

Corollary 3. *Let $m \geq 2$. If $\bar{A} \in (\mathbb{Z}/m\mathbb{Z})[[x]]$ satisfies $\bar{A} = x + O(x^2)$ and (1) with all coefficients read modulo m , then $a(n) \equiv [x^n]\bar{A} \pmod{m}$ for every n .*

Proof. Reduction modulo m is a ring homomorphism, so it carries the integer recursion of Lemma 2 to the same recursion over $\mathbb{Z}/m\mathbb{Z}$, where the coefficient of $a(N)$ is still 1 and hence still invertible. That recursion therefore has a unique solution, and both $a \bmod m$ and the coefficient sequence of $\bar{A}$ solve it. $\square$

3 The reduction modulo 2

Write $f(x) = \frac{x}{1-x}$ over $\mathbb{Z}/2\mathbb{Z}$.

Lemma 4. *$f^j(x) = \frac{x}{1-jx}$ for every $j \geq 1$, and more generally for every integer j , where f^{-1} denotes the compositional inverse.*

Proof. For $j = 1$ this is the definition. If $f^j(x) = x/(1-jx)$ then

$$f^{j+1}(x) = f(f^j(x)) = \frac{\frac{x}{1-jx}}{1 - \frac{x}{1-jx}} = \frac{x}{(1-jx) - x} = \frac{x}{1 - (j+1)x}.$$

The maps $x \mapsto x/(1-jx)$ therefore form a group isomorphic to $(\mathbb{Z}, +)$ under composition, which also gives the statement for $j \leq 0$; in particular $f^{-1}(x) = x/(1+x)$. $\square$

Lemma 5. *Over $\mathbb{Z}/2\mathbb{Z}$, f satisfies $f = x + f^2 f^3$.*

Proof. By Lemma 4, $f^2 f^3 = \frac{x^2}{(1-2x)(1-3x)}$, while $f - x = \frac{x}{1-x} - x = \frac{x^2}{1-x}$. So the claim is that $(1-2x)(1-3x)$ and $1-x$ agree in $(\mathbb{Z}/2\mathbb{Z})[x]$. Expanding,

$$(1-2x)(1-3x) = 1 - 5x + 6x^2,$$

and modulo 2 we have $5 \equiv 1$ and $6 \equiv 0$. Hence both sides equal $1-x$, and the two rational functions agree. $\square$

Theorem 6. Over $\mathbb{Z}/2\mathbb{Z}$ the reduction of A is $x/(1-x)$. Equivalently $a(n) \equiv 1 \pmod{2}$ for every $n \geq 1$, and more generally

$$[x^n]A^j(x) \equiv j^{n-1} \pmod{2}$$

for every $n \geq 1$ and every integer j . This is the conjecture of Section 1.

Proof. By Lemma 5, $f = x/(1-x)$ satisfies the reduced equation, and $f = x + O(x^2)$. By Corollary 3 the reduction of A equals f , so $a(n) \equiv [x^n]f = 1$.

Reduction modulo 2 carries composition of power series with unit linear coefficient to composition, so A^j reduces to f^j for every integer j — for $j < 0$ because A has linear coefficient 1, hence is invertible under composition, and the reduction map is a homomorphism of the composition group. By Lemma 4, $f^j(x) = x/(1-jx) = \sum_{n \geq 1} j^{n-1}x^n$, which gives the second statement. $\square$

Remark 7. The proof used only that $(1-2x)(1-3x) \equiv 1-x$ modulo 2, that is, that 2 divides both $2 \cdot 2$ and $2 \cdot 3$. The largest modulus with that property is $\gcd(2p, p(p+1))$, which is $2p$ for odd p and p for even p ; here $p = 2$ gives 2. This is exactly why the statement is made modulo 2 and not modulo a larger number.

4 Verification

Three checks, each independent of the algebra above.

First, the recursion of Lemma 2 was run in exact rational arithmetic and its output compared against the entry's DATA at every one of the 21 published indices; the values agree exactly, and every one is an integer. A recursion that reproduced only some of them would mean the functional equation had been transcribed wrongly.

Second, at every step the coefficient of $a(N)$ was computed rather than assumed, by evaluating the order- x^N equation at $a(N) = 0$ and at $a(N) = 1$ and subtracting; it came out 1 at every N up to 24, as Lemma 2 requires. The same run was continued to $n = 24$ and every term reduced modulo 2; all residues are 1.

Third, the composition of the proof was carried out independently as a formal power series over $\mathbb{Z}/2\mathbb{Z}$ to order 30, starting from the coefficients of $x/(1-x)$ rather than from the closed form; both sides of (1) agree at every order, and the iterates f^j were confirmed to have coefficients j^{n-1} for $j \leq 6$.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A396099.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999.
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, 2008.