

The empirical recurrence for OEIS A229641, proved by counting defective colourings

Adrian Perez Fontelles
Independent researcher

1 September 2026

Abstract

OEIS A229641 counts the colourings of an array with $K = 3$ colours that are proper except at exactly two adjacent pairs, the colours being introduced in row major order. Neither feature is local: the number of monochromatic adjacent pairs is a property of the whole array, and the row major clause fixes a representative of each relabelling class. Both are removed by machinery already in hand — the running count of mistakes goes into the state, capped at 2, and the colour clause disappears into the identity $3!a(n) = \sum_i \binom{3}{i} D_{3-i} L_i(n)$ with D_m the derangement numbers. What is left is a walk count on $S = 828$ vertices, and the empirical recurrence of order 30 contributed by R. H. Hardin follows from a finite exact computation.

2020 Mathematics Subject Classification. 05A15, 05C15, 15B36.

1 The sequence and the conjecture

OEIS A229641 is “Number of defective 3-colorings of a 5 X n 0..2 array connected horizontally, diagonally and antidiagonally with exactly two mistakes, and colors introduced in row-major 0..2 order.”. It has offset 1 and begins

0, 925, 8800, 89331, 960325, 9722206, 98015235, 960209886, ...

The entry carries the following comment:

Empirical: $a(n) = 18*a(n-1) - 15*a(n-2) - 1191*a(n-3) + 3468*a(n-4) + 31983*a(n-5) - 108707*a(n-6) - 424479*a(n-7) + 1614489*a(n-8) + 2851598*a(n-9) - 13478352*a(n-10) - 7797810*a(n-11) + 66347438*a(n-12) - 12632640*a(n-13) - 190067859*a(n-14) + 147888798*a(n-15) + 287096430*a(n-16) - 412178619*a(n-17) - 138135927*a(n-18) + 522690894*a(n-19) - 168525432*a(n-20) - 273411640*a(n-21) + 227616084*a(n-22) + 9610272*a(n-23) - 73617984*a(n-24) + 21460464*a(n-25) + 7359984*a(n-26) - 4611168*a(n-27) + 186624*a(n-28) + 279936*a(n-29) - 46656*a(n-30)$ for $n > 33$.

As of the “Last modified” line on the live entry (Apr 28 2021 01:22:57, revision 8) the statement is still recorded as empirical, and nothing on the entry records it as settled either way.

Written out, the claim is

$$a(n) = 18a(n-1) - 15a(n-2) - 1191a(n-3) + 3468a(n-4) + 31983a(n-5) - 108707a(n-6) - 424479a(n-7) + 1614489a(n-8) + 2851598a(n-9) - 13478352a(n-10) - 7797810a(n-11) + 66347438a(n-12) - 12632640a(n-13) - 190067859a(n-14) + 147888798a(n-15) + 287096430a(n-16) - 412178619a(n-17) - 138135927a(n-18) + 522690894a(n-19) - 168525432a(n-20) - 273411640a(n-21) + 227616084a(n-22) + 9610272a(n-23) - 73617984a(n-24) + 21460464a(n-25) + 7359984a(n-26) - 4611168a(n-27) + 186624a(n-28) + 279936a(n-29) - 46656a(n-30) \quad (1)$$

for all $n > 33$.

2 Mistakes, and the colour clause

The arrays have 5 rows and $L = n$ columns, entries in $\{0, \dots, 3 - 1\}$. Two cells are adjacent when they differ by one of

$$\mathcal{N} = \{(1, 0), (1, 1), (1, -1)\}$$

(as (column shift, row shift)); each unordered adjacency is represented by exactly one offset, and every offset has a nonnegative column shift, so the pairs inside a column are counted when that column is laid down and the pairs joining two columns when the later one is. Call an adjacent pair a *mistake* when $x_{t,u} = x_{t+d_1, u+d_2}$. The entry asks for exactly $E = 2$ mistakes.

That is a statement about the whole array, so no bounded window sees it; but the running number of mistakes can be carried in the state, capped at 2, the states past 2 being absent because a count that has passed the budget can never come back.

The condition is stated in equalities, so it is invariant under relabelling the colours, and so is the number of mistakes. “Colors introduced in row-major order” keeps exactly one colouring from each relabelling class, so the entry counts equality PATTERNS with at most $K = 3$ classes.

Lemma 1. *Let $L_i(n)$ count the colourings over i colours with exactly 2 mistakes, and $N_j(n)$ the admissible patterns with exactly j classes. Then $L_i = \sum_j N_j i(i-1) \cdots (i-j+1)$ and*

$$3! \sum_{j \leq 3} N_j(n) = \sum_{i=0}^3 \binom{3}{i} D_{3-i} L_i(n), \quad D_m = m! \sum_{t=0}^m \frac{(-1)^t}{t!}.$$

Proof. A colouring is a pattern together with an injection of its classes into the palette, and a pattern with j classes admits $i(i-1) \cdots (i-j+1)$ of them, which is the first identity. Writing the falling factorial as $j! \binom{i}{j}$ and inverting binomially gives $j! N_j = \sum_i (-1)^{j-i} \binom{j}{i} L_i$; summing over $j \leq 3$ and exchanging the order of summation makes the coefficient of L_i equal to $\sum_{t \leq 3-i} (-1)^t / (i! t!)$, which multiplied by $3!$ is $\binom{3}{i} D_{3-i}$, an integer. $\square$

3 Each L_i is a walk count

Fix i and take as vertices the pairs (r, c) with $r \in \{0, \dots, i-1\}^5$ a column and $0 \leq c \leq 2$ a running count. Write $\beta(r, s)$ for the number of mistakes joining columns r and s plus the number of mistakes inside s . Declare $(r, c) \rightarrow (s, c + \beta(r, s))$ an edge when $c + \beta(r, s) \leq 2$, let $\mathbf{s}_i$ mark the vertices whose count is the number of mistakes inside r alone, and $\mathbf{e}_i$ those with $c = 2$.

Lemma 2. $L_i(n) = \mathbf{s}_i^\top P_i^{L-1} \mathbf{e}_i$ for $L \geq 1$, where P_i is the adjacency matrix of that digraph.

Proof. A walk of length $L-1$ determines the sequence of L columns, and conversely; by induction the count component after t columns is the number of mistakes among the first t columns, since each step adds exactly the mistakes joining columns t and $t+1$ together with those inside column $t+1$, and each unordered pair is named by exactly one offset. The walk exists precisely when the running count never exceeds 2, which for a nondecreasing count follows from its final value being 2, and $\mathbf{e}_i$ imposes that final value. The state carries a column, not a boundary, so L columns make a walk of length $L-1$. $\square$

Assembling the 3 chains block-diagonally into N of size $S = 828$ with the weights of Lemma 1 in the start vector $\mathbf{v}$ and the end markers in $\mathbf{u}$,

$$3! a(n) = \mathbf{v}^\top N^{L-1} \mathbf{u}, \quad L = n. \quad (2)$$

4 The criterion, and the computation

Let $q(t) = t^{30} - \sum_i c_i t^{30-i}$ be the characteristic polynomial of (1) and $G = N$.

Lemma 3. *With n_0 the least index for which $L \geq 1$, $o = 1n_0 + 0 - 1$ and $h_j = G^j N^o \mathbf{u}$, we have for $j \geq 30$*

$$3! \left(a(n_0 + j) - \sum_i c_i a(n_0 + j - i) \right) = \mathbf{v}^\top G^{j-30} w, \quad w = h_{30} - \sum_i c_i h_{30-i}.$$

Hence (1) holds from that point on if and only if $\mathbf{v}^\top G^m w = 0$ for all $m \geq 0$, and $m < S$ suffices.

Proof. Substitute (2) and factor; $3!$ is a nonzero scalar. Cayley–Hamilton makes G^S an integer combination of $I, G, \dots, G^{S-1}$, so every later value repeats that combination of earlier ones. $\square$

Theorem 4. *The recurrence (1) holds for every $n > 33$, which is the statement of Section 1.*

Proof. The digraph was built directly from the definitions above. The vector w was formed by 30 applications of G in exact integer arithmetic and $\mathbf{v}^\top G^m w$ evaluated for $m = 0, 1, \dots$, again exactly. Every one of those integers vanishes from the index corresponding to $n = 33 + 1$ onwards, and the run continued until $S + 1$ consecutive zeros had been seen. $\square$

5 Verification

Three checks, each independent of the algebra above.

First, the right-hand side of (2), divided by $3!$, was evaluated at every one of the 17 published indices and agrees with the entry’s DATA exactly, the quotient coming out an integer each time; no shift was fitted, the number of columns being $L = n$ from the entry’s own name and the offset saying which index carries the first published term. Double-counting an unordered pair, or mistaking the budget, changes the counts at once.

Second, the recurrence (1) was evaluated on those published terms in exact integer arithmetic, with no matrices involved, and vanishes wherever it applies.

Third, the annihilation test of Lemma 3 was carried to the full Cayley–Hamilton bound in exact integer arithmetic rather than to a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A229641.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7; falling-factorial inversion, Section 1.9.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.