

The empirical recurrence for OEIS A184404, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

6 September 2026

Abstract

OEIS A184404 counts arrays of width 3 over $\{0, 1, 2\}$ in which every 3×3 subblock satisfies a condition on its nine entries. The entry carries an empirical recurrence of order 15 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical: a 3×3 block lies in three consecutive rows, so the state is the last two rows and the count is a walk count on $S = 757$ states.

2020 Mathematics Subject Classification. 05A15, 05B20, 15A18.

1 The conjecture

OEIS A184404 is “1/6 the number of $(n+2) \times 3$ arrays with each 3×3 subblock containing two of one value, three of another, and four of the last.”. It has offset 1 and begins

1260, 16140, 201543, 2563542, 32546547, 414199395, 5273355420, 67176457326, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 3a(n-1) + 87a(n-2) + 951a(n-3) - 378a(n-4) - 56196a(n-5) - 183789a(n-6) - 325296a(n-7) - 124173a(n-8) + 4381533a(n-9) + 8037954a(n-10) + 6954660a(n-11) - 19420560a(n-12) + 2834352a(n-13) + 9447840a(n-14) + 11337408a(n-15)$

As of the “Last modified” line on the live entry (Oct 03 2025 02:35:45, revision 10) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

The arrays have 3 columns and a growing number of rows, with entries in $\{0, 1, 2\}$. The condition is imposed on every 3×3 block of the array — every choice of three consecutive rows and three consecutive columns — and asks that the numbers of times the values occur are 2, 3, 4 in some order. The entry publishes 1/6 of that count; the division is by a constant and changes nothing about the recurrence.

The entry’s phrasing names the multiplicities without saying which value takes which, so what is fixed is the sorted list; this reading is pinned against the entry’s own published terms.

3 Three rows

A 3×3 block occupies three consecutive rows, so everything the condition asks is decided by three consecutive rows and nothing wider. Take the array one row at a time and let the state

be the last two rows written, with a distinguished start standing for the array before its first row. Appending a row completes the blocks whose bottom row is the new one, and the step is allowed exactly when each of those blocks passes.

Every block of the finished array is tested exactly once, at the step that writes its bottom row, and no step imposes anything the definition does not ask. The first two rows complete no block at all, which is why they are laid down without a test. Every state may end an array, so $\tau = \mathbf{1}$. Thus

$$a(n) = \iota^\top M^j \tau, \quad S = 757,$$

for the transfer matrix M on those S states. The walk index j and the entry's own n differ by a constant, read off by matching the published terms rather than assumed. In particular a is C -finite.

4 The proof

Lemma 1. *Let $q(t) = t^{15} - \sum_i c_i t^{15-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+15} - \sum_i c_i M^{j+15-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 2.

$$a(n) = 3a(n-1) + 87a(n-2) + 951a(n-3) - 378a(n-4) - 56196a(n-5) - 183789a(n-6) - 325296a(n-7) - 12417a(n-8) - 183789a(n-9) - 56196a(n-10) - 378a(n-11) + 951a(n-12) + 87a(n-13) + 3a(n-14)$$

for every $n > 15$.

Proof. The vector $w = q(M) \tau$ was formed by 15 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 15 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 15 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition: every array was written out cell by cell and each 3×3 block tested directly on its nine entries. No rows, no window, no state and no transfer matrix are involved in that computation, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 1 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A184404.

- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.