

# The empirical recurrence for OEIS A266132, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

3 September 2026

## Abstract

OEIS A266132 counts arrays of width 2 in which every entry is equal to the NUMBER of its own neighbours, in named directions, that stand in a stated relation to it. The entry carries an empirical recurrence of order 17 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. The condition is self-referential but local: a cell's neighbours lie in the row above, its own row and the row below, so a window of three consecutive rows decides the condition for the middle one, and the count is a walk count on  $S = 481$  states.

2020 Mathematics Subject Classification. 05A15, 05C30, 15A18.

## 1 The conjecture

OEIS A266132 is “Number of  $2 \times n$  integer arrays with each element equal to the number of horizontal, diagonal and antidiagonal neighbors exactly one smaller than itself.”. It has offset 1 and begins

$$1, 9, 25, 49, 149, 625, 2141, 5433, \dots$$

The entry states, as a conjecture contributed by its author and never marked settled:

$$\begin{aligned} \text{Empirical: } a(n) = & a(n-1) + 8*a(n-3) + 80*a(n-4) - 40*a(n-5) - 52*a(n-6) - 112*a(n-7) \\ & - 1204*a(n-8) + 292*a(n-9) + 64*a(n-10) - 880*a(n-11) + 4016*a(n-12) - 2112*a(n-13) \\ & - 1152*a(n-14) + 1024*a(n-15) - 4608*a(n-16) + 4608*a(n-17) \end{aligned}$$

As of the “Last modified” line on the live entry (Dec 21 2015 12:29:49, revision 4) this is still recorded as empirical, and nothing on the entry records it as proved.

## 2 What is being counted

The array has 2 cells across. The NEIGHBOURS of a cell are the cells reached from it by the offsets

$$(-1, -1), (-1, 0), (-1, 1), (1, -1), (1, 0), (1, 1),$$

those that lie inside the array. The condition is that every entry counts its own neighbours that are exactly one smaller than it: writing  $v = x(i, j)$ ,

$$v = \#\{\text{neighbours } y \text{ of } (i, j) : y = v - 1\}.$$

The entry says “integer arrays” and names no alphabet, but one is forced. An entry counts neighbours in the named directions, of which there are 6, so every entry of the array lies in  $\{0, 1, \dots, 6\}$  and the count is finite. That bound is used below and nothing wider is needed. The entry writes the array with  $n$  as the number of COLUMNS; the walk is run down the transpose, so the offsets above are already the transposed ones. Each direction is taken with both signs, so transposing cannot lose one.

### 3 Three rows decide one

**Lemma 1.** *Every offset above changes the row index by  $-1$ ,  $0$  or  $+1$ , so all the neighbours of a cell of row  $i$  lie in rows  $i-1$ ,  $i$  and  $i+1$ . The condition for the whole of row  $i$  is therefore decided by those three rows, an absent row contributing no neighbours.*

*Proof.* Immediate from the offsets, together with the rule that a neighbour outside the array is not a neighbour.  $\square$

So the state is the pair (row above, current row), the row above allowed to be the symbol  $\perp$ ; a step appends a row and settles the condition for the middle row of the window it completes. Two boundaries have to be right and neither is assumed. The walk starts at the state  $(\perp, \perp)$ , which stands for the empty array, so a walk of no steps is the empty array and one step is a one-row array. And the condition for the LAST row is settled by the end vector, which asks whether that row satisfies its condition with nothing written below it: the end vector is therefore NOT the all-ones vector, and a row that counts correctly only when something is written below it does not finish an array. With  $M$  the adjacency matrix,

$$a(n) = \iota^\top M^j \tau, \quad S = 481.$$

The walk index  $j$  and the entry's own  $n$  differ by a constant, read off by matching the published terms rather than assumed. In particular  $a$  is  $C$ -finite.

### 4 The proof

**Lemma 2.** *Let  $q(t) = t^{17} - \sum_i c_i t^{17-i}$  be the characteristic polynomial of the conjectured recurrence. The residual  $a(n) - \sum_i c_i a(n-i)$  equals  $\iota^\top M^j q(M) \tau$  for the corresponding walk index  $j$ , so the recurrence holds from some point on if and only if  $\iota^\top M^j q(M) \tau = 0$  for all large  $j$ , and it is enough to examine  $j < S$ .*

*Proof.* Factor  $M^j$  out of  $M^{j+17} - \sum_i c_i M^{j+17-i}$ . For the bound,  $M^S$  is an integer combination of  $I, M, \dots, M^{S-1}$  by Cayley–Hamilton, so the numbers  $u_j = \iota^\top M^j q(M) \tau$  obey the monic recurrence given by the characteristic polynomial of  $M$ ;  $S$  consecutive zeros force every later one, and the last nonzero  $u_j$  pins the threshold exactly.  $\square$

**Theorem 3.**

$$a(n) = a(n-1) + 8a(n-3) + 80a(n-4) - 40a(n-5) - 52a(n-6) - 112a(n-7) - 1204a(n-8) + 292a(n-9) + 64a(n-10)$$

for every  $n > 16$ .

*Proof.* The vector  $w = q(M)\tau$  was formed by 17 matrix–vector products in exact integer arithmetic and  $\iota^\top M^j w$  evaluated for  $j = 0, 1, \dots$ , again exactly; those integers vanish from the index corresponding to  $n = 16 + 1$  onwards and the run continued until the working vector was identically zero, which settles every larger  $j$  at once.  $\square$

### 5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 26 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition: enumerate every array of that size, count each cell's neighbours satisfying the relation, and keep the array when every cell equals its own count. No window, no transfer matrix and no transposition are involved, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

## References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A266132.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] J. E. Hopcroft, R. Motwani and J. D. Ullman, *Introduction to Automata Theory, Languages, and Computation*, 3rd ed., Addison–Wesley, 2006.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.