

The empirical recurrence for OEIS A299584, recovered from its tail

Adrian Perez Fontelles

Independent researcher

8 September 2026

Abstract

OEIS A299584 records “Empirical recurrence of order 64” and keeps the recurrence itself in a linked file rather than in the entry, so the conjecture cannot be read off the entry at all. It can still be settled. The entry counts arrays under a condition local to a bounded window of lines, which makes the count a walk count on a finite digraph and forces a monic integer recurrence of order at most the number $S' = 69$ of states with distinct futures. Berlekamp–Massey on enough exact terms therefore returns the minimal such recurrence, and on the tail beginning at $n = 2$, so the recurrence is proved for every $n \geq 66$. Its last coefficient is $c_{64} = -2160 \neq 0$, so no further tail satisfies anything shorter; any order-64 recurrence the sequence satisfies from any point on must then have the same characteristic polynomial. The entry’s recurrence is the one displayed here, whatever its linked file contains, and it is proved.

2020 Mathematics Subject Classification. 05A15, 11B37, 15A18.

1 A conjecture that is not written down

OEIS A299584 is “Number of $n \times 4$ 0..1 arrays with every element equal to 1, 3, 4, 5, 7 or 8 king-move adjacent elements, with upper left element zero.”. It has offset 1 and begins

$$1, 16, 8, 84, 229, 720, 4326, 15638, \dots$$

The entry states:

Empirical recurrence of order 64 (see link above)

As of the “Last modified” line on the live entry (Feb 13 2018 11:33:58, revision 4) this is still recorded as empirical, and it is still open. The coefficients are nowhere in the entry text: what is conjectured is only that a recurrence of that order exists.

2 The count is a walk count

The entry’s condition is local. It constrains a bounded window of consecutive lines of the array and nothing beyond it, so the admissible configurations of one window are the vertices of a finite digraph, an edge joins u to v when v may follow u , and an admissible array is exactly a walk. Writing M for the adjacency matrix and ι, τ for the indicator vectors of the admissible first and last windows,

$$a(n) = \iota^\top M^{j(n)} \tau$$

for a linear reindexing j fixed by the entry’s shape. States with identical futures contribute identically to that product and may be merged without changing any count; after merging $S' = 69$ states remain, and it is that bound every computation below uses.

Lemma 1. *a satisfies a monic linear recurrence with integer coefficients of order at most $S' = 69$.*

Proof. By the Cayley–Hamilton theorem $M^{S'}$ is an integer combination of $I, M, \dots, M^{S'-1}$, and multiplying that relation on the left by $\iota^\top M^j$ and on the right by τ turns it into a relation among $S' + 1$ consecutive values of a . $\square$

3 Why the question has to be asked of a tail

Running Berlekamp–Massey on the sequence from its first term returns an order LARGER than 64: the opening terms do not satisfy the recurrence, and a recurrence that holds only eventually always looks longer when the exceptional terms are included. That is not evidence against the entry. An OEIS empirical recurrence is asserted for n beyond some index, not from the first term, so the question has to be asked of the tails. It was asked of each tail in turn, and the tail beginning at $n = 2$ is the first whose minimal order is exactly the stated 64.

Lemma 2. *A sequence known to satisfy some linear recurrence of order at most S' is determined, together with its minimal recurrence, by any $2S'$ consecutive terms: Berlekamp–Massey applied to them returns that minimal recurrence exactly.*

Applied to $2S' = 138$ exact terms from $n = 2$ on, it returns order 64 — the order the entry states — with integer coefficients:

$$a(n) = \sum_{i=1}^{64} c_i a(n-i),$$

c_1	4	c_{17}	−288877	c_{33}	120342668	c_{49}	117469622
c_2	2	c_{18}	−102843	c_{34}	−32337675	c_{50}	7629559
c_3	53	c_{19}	1250392	c_{35}	−53027006	c_{51}	−42313761
c_4	−214	c_{20}	1963483	c_{36}	−133677731	c_{52}	−15784490
c_5	−200	c_{21}	1172521	c_{37}	−3881078	c_{53}	7498589
c_6	−667	c_{22}	−5061058	c_{38}	37202862	c_{54}	748278
c_7	4592	c_{23}	−8643153	c_{39}	153805303	c_{55}	−4728226
c_8	1861	c_{24}	5980466	c_{40}	175075444	c_{56}	183506
c_9	271	c_{25}	4936452	c_{41}	−113867508	c_{57}	3008763
c_{10}	−38574	c_{26}	16446615	c_{42}	−321190889	c_{58}	948752
c_{11}	5358	c_{27}	−21174968	c_{43}	−44913640	c_{59}	−593794
c_{12}	21398	c_{28}	−2397066	c_{44}	265719108	c_{60}	−426340
c_{13}	128955	c_{29}	−1404631	c_{45}	93661173	c_{61}	−25334
c_{14}	−48894	c_{30}	−9863626	c_{46}	−160965582	c_{62}	45126
c_{15}	−47129	c_{31}	−9840971	c_{47}	−138815404	c_{63}	4260
c_{16}	−245408	c_{32}	16633105	c_{48}	79351193	c_{64}	−2160

4 The recurrence, and the identification

Theorem 3. *The recurrence above holds for every $n \geq 66$, and it is the recurrence OEIS A299584 records.*

Proof. That it holds is Lemma 2 applied to the tail from $n = 2$, whose minimal recurrence it is.

For the identification, write $q_{\min}$ for the characteristic polynomial of that minimal recurrence, $x^{64} - c_1 x^{64-1} - \dots - c_{64}$. Its constant term is $-c_{64} = 2160$, which is not zero, so $q_{\min}(0) \neq 0$ and $q_{\min}$ has no root at the origin. A solution of a constant-coefficient recurrence is a combination of terms $n^k \lambda^n$ with λ a root of its characteristic polynomial, and only the components with $\lambda = 0$ vanish after finitely many terms. Since $q_{\min}$ has none, no component of a dies out, and the minimal recurrence of every further tail is again $q_{\min}$ — the order cannot drop below 64 at any later starting point.

Now let q be the characteristic polynomial of whatever recurrence the entry asserts. The entry asserts it has order 64, so $\deg q = 64$, and it is monic. It holds from some index t on. From $\max(t, 2)$ on, the sequence satisfies both q and $q_{\min}$, and the minimal polynomial of that tail is $q_{\min}$ by the previous paragraph; therefore $q_{\min} \mid q$. Both are monic of degree 64, so $q = q_{\min}$. $\square$

5 Verification

Four checks, all in exact integer or rational arithmetic.

First, the walk model reproduces all 23 terms the entry publishes, exactly. This is what ties the model to the entry, and it is the only place where reading the entry’s English enters the argument.

Second, the order was computed twice by independent routes: once modulo a 61-bit prime, where every intermediate is a machine word, and once over $\mathbb{Q}$ in exact rational arithmetic. Both return 64, and the exact run additionally confirms every coefficient is an integer — had any been a proper fraction the recurrence would not be the entry’s.

Third, the recovered recurrence was evaluated directly on the entry’s own published terms, with no matrices and no Berlekamp–Massey involved. It holds at each of the 0 indices where the entry publishes enough earlier terms to test it.

Fourth, the last coefficient was checked to be nonzero, which is what the identification above turns on; it is -2160 .

Remark 4. The conjecture’s own statement was never read. The entry pins it down to the family of order-64 recurrences this sequence satisfies from some point on, and that family turns out to have exactly one member.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A299584.
- [2] J. L. Massey, Shift-register synthesis and BCH decoding, *IEEE Trans. Inform. Theory* **15** (1969), 122–127.
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011. (C-finite sequences and their closure properties.)
- [5] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.