

# The empirical recurrence for OEIS A218523, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

2 September 2026

## Abstract

OEIS A218523 counts the distinct arrays obtained by applying a fixed local operation to every cell of an array over a bounded alphabet, and carries an empirical recurrence of order 16 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. What is counted is the IMAGE of a map, not its domain, so it is not a walk count in the obvious graph; determinising the machine that emits the derived array row by row turns it into one, on  $S = 155$  states, after which the conjectured recurrence is settled by a finite exact computation.

2020 Mathematics Subject Classification. 05A15, 68Q45, 15B36.

## 1 The conjecture

OEIS A218523 is “Number of  $2Xn$  arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random  $0..3$   $2Xn$  array.”. It has offset 1 and begins

16, 36, 286, 1348, 7902, 42812, 242438, 1349524, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical:  $a(n) = 2*a(n-1) + 17*a(n-2) + 26*a(n-3) - 19*a(n-4) - 123*a(n-5) - 138*a(n-6) - 248*a(n-7) - 329*a(n-8) + 41*a(n-9) + 1108*a(n-10) + 1469*a(n-11) - 54*a(n-12) - 1158*a(n-13) - 316*a(n-14) + 144*a(n-15) + 32*a(n-16)$

As of the “Last modified” line on the live entry (Oct 03 2025 17:41:57, revision 10) this is still recorded as empirical, and nothing on the entry records it as proved.

## 2 What is being counted

Let  $R = \{0, \dots, 3\}^2$  be the set of rows and let the neighbours of a cell be the cells at the offsets

$(-1,0), (-1,1), (1,-1), (1,0)$ , those falling outside the array being simply absent. The entry writes the array with  $n$  as the number of COLUMNS; transposing it exchanges the two coordinates of every neighbour offset, which is done once so that the walk always runs down the rows. For an array  $A = (r_1, \dots, r_n) \in R^n$  let  $\Phi(A)$  be the array whose cell  $(i, j)$  carries the minimum of  $A(i, j)$  and of the values at its neighbours. The entry counts

$$a(n) = |\{\Phi(A) : A \in R^n\}|,$$

the size of an IMAGE. That is the whole difficulty: two different arrays with the same minimum-array must be counted once, and a walk count on the arrays would count them twice.

Because every offset above moves the row index by at most one, row  $i$  of  $\Phi(A)$  depends only on rows  $i-1, i, i+1$  of  $A$ . Write  $\beta(p, c, x)$  for that row, with  $p$  or  $x$  allowed to be  $\perp$  for “outside”; then

$$\Phi(r_1, \dots, r_n) = (\beta(\perp, r_1, r_2), \dots, \beta(r_{n-1}, r_n, \perp)).$$

### 3 The image is a walk count after determinisation

For a set  $D$  of pairs  $(p, c)$  and a derived row  $b$  put

$$\delta(D, b) = \{(c, x) : (p, c) \in D, x \in R, \beta(p, c, x) = b\}, \quad f(D) = |\{\beta(p, c, \perp) : (p, c) \in D\}|,$$

and let  $D_0 = \{(\perp, r) : r \in R\}$ .

**Lemma 1.** *For every  $t \geq 0$ ,  $D_t = \delta(\dots \delta(D_0, b_1) \dots, b_t)$  is exactly the set of pairs  $(r_t, r_{t+1})$  of consecutive rows completing a prefix  $r_1, \dots, r_{t+1}$  whose first  $t$  derived rows are  $b_1, \dots, b_t$ . Consequently, for  $n \geq 1$ ,*

$$a(n) = \sum_{b_1, \dots, b_{n-1}} f(D_{n-1}),$$

*the empty sets contributing nothing.*

*Proof.* Induction on  $t$ :  $D_0$  is the set of pairs  $(\perp, r_1)$ , and  $\delta(D_t, b)$  collects the pairs  $(r_{t+1}, r_{t+2})$  extending a consistent prefix with  $\beta(r_t, r_{t+1}, r_{t+2}) = b$ . An element of the image is a full derived array  $(b_1, \dots, b_n)$ ; its prefix  $(b_1, \dots, b_{n-1})$  is realisable exactly when  $D_{n-1} \neq \emptyset$ , and then the possible last rows are  $\{\beta(p, c, \perp) : (p, c) \in D_{n-1}\}$ , of which there are  $f(D_{n-1})$ . Distinct sequences are distinct derived arrays, so each is counted once.  $\square$

So with  $M$  the adjacency matrix of the digraph whose vertices are the reachable nonempty sets and which carries one edge  $D \rightarrow \delta(D, b)$  for each  $b$  with  $\delta(D, b) \neq \emptyset$  — several  $b$  may lead to the same set, and then the edge is repeated, which is exactly right, since the derived arrays differ —

$$a(n) = \iota^\top M^{n-1} \tau, \quad \iota = e_{D_0}, \quad \tau = f.$$

There are  $S = 155$  reachable sets, generated from  $D_0$  outward, and  $a$  is  $C$ -finite. Note that  $\tau$  is not the all-ones vector: the last derived row is not a step of the walk but a count attached to the state the walk stops in.

### 4 The proof

**Lemma 2.** *Let  $q(t) = t^{16} - \sum_i c_i t^{16-i}$  be the characteristic polynomial of the conjectured recurrence. The residual  $a(n) - \sum_i c_i a(n-i)$  equals  $\iota^\top M^j q(M) \tau$  for the corresponding walk index  $j$ , so the recurrence holds from some point on if and only if  $\iota^\top M^j q(M) \tau = 0$  for all large  $j$ , and it is enough to examine  $j < S$ .*

*Proof.* Factor  $M^j$  out of  $M^{j+16} - \sum_i c_i M^{j+16-i}$ . For the bound,  $M^S$  is an integer combination of  $I, M, \dots, M^{S-1}$  by Cayley–Hamilton, so the numbers  $u_j = \iota^\top M^j q(M) \tau$  obey the monic recurrence given by the characteristic polynomial of  $M$ ;  $S$  consecutive zeros force every later one, and the last nonzero  $u_j$  pins the threshold exactly.  $\square$

**Theorem 3.**

$a(n) = 2a(n-1) + 17a(n-2) + 26a(n-3) - 19a(n-4) - 123a(n-5) - 138a(n-6) - 248a(n-7) - 329a(n-8) + 41a(n-9)$   
for every  $n > 16$ .

*Proof.* The vector  $w = q(M)\tau$  was formed by 16 matrix–vector products in exact integer arithmetic and  $\iota^\top M^j w$  evaluated for  $j = 0, 1, \dots$ , again exactly; those integers vanish from the index corresponding to  $n = 16 + 1$  onwards and the run continued until the working vector was identically zero, which settles every larger  $j$  at once.  $\square$

## 5 Verification

Three checks, each independent of the algebra above.

First, the model reproduces all 20 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: the machine is built by reading the entry’s English, and a misreading gives different counts.

Second, the conjectured recurrence was evaluated directly on those published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Third, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

## References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A218523.
- [2] J. E. Hopcroft, R. Motwani and J. D. Ullman, *Introduction to Automata Theory, Languages, and Computation*, 3rd ed., Addison–Wesley, 2006. (The subset construction, Section 2.3.)
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.