

# A proof of the conjectured recurrence for OEIS A026026

Adrian Perez Fontelles

Independent researcher

30 August 2026

## Abstract

OEIS A026026 carries a conjectured linear recurrence with polynomial coefficients, of order 2. The entry posts no generating function, so the usual route through a functional equation is unavailable. It does post a closed form for  $a(n)$ , and that is enough: the closed form is a sum of finitely many hypergeometric terms, and for such a term every shift quotient is a rational function of  $n$ . The conjectured recurrence therefore reduces to one rational-function identity, one for each similarity class of terms, and each is decided exactly by cancellation. No expansion is truncated and nothing is guessed; the procedure returns a proof or a refutation. Here it returns a proof, valid for every  $n$  outside an explicitly computed finite set.

2020 Mathematics Subject Classification. 33F10, 11B37, 05A10.

## 1 The sequence and the conjecture

OEIS A026026 is “ $a(n)$  = number of  $(s(0), s(1), \dots, s(2^n-1))$  such that  $s(i)$  is a nonnegative integer and  $|s(i) - s(i-1)| = 1$  for  $i = 1, 2, \dots, n$ ,  $s(0) = 3$ ,  $s(2^n-1) = 4$ . Also  $a(n) = T(2^n-1, n-1)$ , where  $T$  is defined in A026022”. It has offset 2 and begins

$$a(2), \dots, a(9) = 1, 3, 10, 35, 125, 451, 1638, 5980, \dots$$

The entry states, not as a conjecture,

$$a(n) = \text{binomial}(2^n-3, n-2) - \text{binomial}(2^n-3, n-6). \text{ - } Christian Krause, \text{ May 09 2026}$$

that is,

$$a(n) = -\binom{2n-3}{n-6} + \binom{2n-3}{n-2}. \quad (1)$$

This is the only input the proof takes from the entry, and it was checked against the entry’s own DATA before use.

The entry separately carries the following comment:

**Conjecture:**  $-(n+3)(11n-38)a(n) + 2(35n^2-86n-102)a(n-1) - 4(13n-30)(2n-5)a(n-2) = 0$ . - *R. J. Mathar*, Jun 22 2013

As of the “Last modified” line on the live entry (2026-05-14, revision 27) the statement is still recorded as a conjecture, and no proof appears among the entry’s links, formulas or programs.

Written out, the conjecture is

$$-(n+3)(11n-38)a(n) + 2(35n^2-86n-102)a(n-1) - 4(2n-5)(13n-30)a(n-2) = 0 \quad (2)$$

## 2 Hypergeometric terms

**Definition 1.** A nonzero expression  $c(n)$  is a *hypergeometric term* in  $n$  if the quotient  $c(n+1)/c(n)$  is a rational function of  $n$ . Two hypergeometric terms  $c$  and  $d$  are *similar*, written  $c \sim d$ , if  $c(n)/d(n)$  is a rational function of  $n$ .

Similarity is an equivalence relation on hypergeometric terms. Products of factorials, binomial coefficients with arguments linear in  $n$ , exponentials  $z^n$  and polynomials are hypergeometric, and so is any product of these; a sum of them need not be.

**Lemma 2.** *If  $c$  is a hypergeometric term then for every integer  $i \geq 0$  the shift quotient  $c(n-i)/c(n)$  is a rational function of  $n$ , computable from  $c(n+1)/c(n)$  by a finite number of substitutions and multiplications.*

*Proof.* Write  $\rho(n) = c(n+1)/c(n)$ , a rational function by hypothesis. Then  $c(n-1)/c(n) = 1/\rho(n-1)$ , and inductively  $c(n-i)/c(n) = \prod_{j=1}^i \rho(n-j)^{-1}$ , a finite product of rational functions.  $\square$

The fact that makes the test complete rather than merely sufficient is the linear independence of dissimilar terms. It is classical; see Petkovšek, Wilf and Zeilberger [1], Chapter 5.

**Lemma 3.** *Let  $c_1, \dots, c_m$  be hypergeometric terms, pairwise dissimilar. If  $\sum_{j=1}^m u_j(n)c_j(n) = 0$  for rational functions  $u_j$  and all large  $n$ , then every  $u_j$  is the zero rational function.*

## 3 The recurrence as a finite set of rational identities

**Proposition 4.** *Let  $a(n) = \sum_{j=1}^m c_j(n)$  with each  $c_j$  hypergeometric, and let  $p_0, \dots, p_r$  be polynomials. Group the  $c_j$  into similarity classes and write each class as  $g_\ell(n) f_{\ell,s}(n)$  for a representative  $g_\ell$  and rational functions  $f_{\ell,s}$ . Put*

$$S_\ell(n) = \sum_{i=0}^r \sum_s p_i(n) f_{\ell,s}(n-i) \frac{g_\ell(n-i)}{g_\ell(n)},$$

*a rational function of  $n$  by Lemma 2. Then*

$$\sum_{i=0}^r p_i(n) a(n-i) = \sum_{\ell} g_\ell(n) S_\ell(n),$$

*and the recurrence  $\sum_i p_i(n)a(n-i) = 0$  holds for all large  $n$  if and only if  $S_\ell = 0$  for every  $\ell$ .*

*Proof.* The displayed identity is the definition of  $S_\ell$  rearranged: each  $a(n-i)$  is the sum of its terms, each term is its class representative times a rational function, and the representatives are pulled out. The representatives  $g_\ell$  are pairwise dissimilar by construction, so Lemma 3 applies to the right-hand side: it vanishes for all large  $n$  exactly when every  $S_\ell$  vanishes identically. Sufficiency is immediate.  $\square$

Proposition 4 turns the conjecture into a finite computation in  $\mathbb{Q}(n)$ , with no truncation and no search. The only  $n$  it says nothing about are those where some  $S_\ell$  has a pole or some  $c_j(n-i)$  is undefined, and these are finite in number and listed below.

## 4 The computation for A026026

The closed form (1) splits into one similarity class, with representative

$$-\binom{2n-3}{n-6}$$

Carrying out Proposition 4 with the coefficients of (2) gives

$$S_1(n) = 0$$

so every class residual vanishes identically and the conjecture follows.

**Theorem 5.** *For all  $n \geq 4$ ,*

$$-(n+3)(11n-38)a(n) + 2(35n^2-86n-102)a(n-1) - 4(2n-5)(13n-30)a(n-2)$$

*Proof.* Immediate from Proposition 4 and the computation above, the excluded values being those at which a class residual has a pole or a term of (1) is undefined.  $\square$

## 5 Verification

Two checks were made, and both are independent of the algebra above.

First, the closed form (1) was evaluated at the first 13 indices and compared term by term with the entry's DATA; the values agree exactly. A formula that did not reproduce the entry's own terms would not have been used.

Second, the conjectured recurrence (2) was evaluated on the published terms in exact integer arithmetic at every index where it applies: 24 instances beginning at  $n = 4$ , all of them zero. This is not part of the proof, which is symbolic, but it would have caught a transcription error in the coefficients.

## References

- [1] M. Petkovšek, H. S. Wilf and D. Zeilberger, *A=B*, A K Peters, Wellesley MA, 1996.
- [2] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A026026](#), 2026-05-14.
- [3] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, 2nd ed., Addison-Wesley, 1994.