

The empirical recurrence for OEIS A204075, proved by counting patterns

Adrian Perez Fontelles
Independent researcher

31 August 2026

Abstract

OEIS A204075 counts arrays subject to a local condition *and* to the clause “new values 0..2 introduced in row major order”, which says the array is written in canonical form. That clause is not local, so a transfer matrix over the alphabet does not see it. It does not have to: what is being counted is not arrays but equality patterns, and the number of patterns is recovered from the numbers L_i of arrays over an i -letter alphabet by inverting a falling factorial. The result is the clean identity $3!a(n) = \sum_i \binom{3}{i} D_{3-i} L_i(n)$ with D_m the derangement numbers, and each L_i is an ordinary walk count. A second reduction — the chain is strongly lumpable over the patterns, because the condition is relabelling-invariant — brings the state space down from 6818 to $S = 1223$. The entry’s empirical recurrence of order 55, contributed by R. H. Hardin, then follows from a finite exact computation.

2020 Mathematics Subject Classification. 05A15, 05A18, 15B36.

1 The sequence and the conjecture

OEIS A204075 is “Number of $(n+1) \times 8$ 0..2 arrays with every 2×2 subblock having equal diagonal elements or equal antidiagonal elements, and new values 0..2 introduced in row major order.”. It has offset 1 and begins

117188, 17061338, 2727300008, 451830740558, 75985220860460, 12862680257580962, 2183735443548857744, 3712

The entry carries the following comment:

Empirical: $a(n) = 322*a(n-1) - 31121*a(n-2) + 677212*a(n-3) + 56848972*a(n-4) - 3586172600*a(n-5) + 52369132896*a(n-6) + 1245052315520*a(n-7) - 47462252573696*a(n-8) + 199722871680128*a(n-9) + 11579233807503360*a(n-10) - 161516422622031360*a(n-11) - 781406452093941760*a(n-12) + 30335388582357307392*a(n-13) - 102551789828754223104*a(n-14) - 2499075111791254233088*a(n-15) + 21696171687994926645248*a(n-16) + 73586462799086990983168*a(n-17) - 1576005803911676145565696*a(n-18) + 2408162877989904740777984*a(n-19) + 55957566883111396665982976*a(n-20) - 25647021377077114345671884*a(n-21) - 882362368937819549787488256*a(n-22) + 8451205698868439960808062976*a(n-23) - 17346065251767803280747*a(n-24) - 141028371040952846016774668288*a(n-25) + 296324312967130989774499741696*a(n-26) + 1198147430715508523506484641792*a(n-27) - 4952845533984364757335295393792*a(n-28) - 3186976950450110274831453782016*a(n-29) + 40412468029396457697759830474752*a(n-30) - 29248858868831293913779436257280*a(n-31) - 177275840840616248911112251113472*a(n-32) + 306730456012993930951282986582016*a(n-33) + 365916593440879671670038106996736*a(n-34) - 1252123486332919207419386477412352*a(n-35) + 31455433167667604131574020308992*a(n-36) + 2662103745182650392788796604153856*a(n-37) - 1794384392309031579471189299429376*a(n-38) - 2855274805218892553424514123300864*a(n-39) + 3752560584513799228760234569760768*a(n-40) + 955018952518786152205067397103616*a(n-41) - 3479885196004780525644200160526336*a(n-42) + 837529518152722930934794025959424*a(n-43) + 1493157166737156760665543722139648*a(n-44) - 855754296042631804236915409944576*a(n-45) - 216580283784251054761779996393472*a(n-46) + 271742943487852984902753988902912*a(n-47) - 26382857255823684476961231470592*a(n-48) - 32280501885628797705911908958208*a(n-49) + 9223849840386162307148405538816*a(n-50) + 583023965817031645837817544704*a(n-51) - 526674746442027343134804934656*a(n-52)$

$$+65171585638020020911233564672*a(n-53)-1944707570813495706151550976*a(n-54)-24175560301554771538477055)$$

As of the “Last modified” line on the live entry (Oct 03 2025 10:58:49, revision 9) the statement is still recorded as empirical, and nothing on the entry records it as settled either way.

Written out, the claim is

$$a(n) = 322 a(n-1) - 31121 a(n-2) + 677212 a(n-3) + 56848972 a(n-4) - 3586172600 a(n-5) + 52369132896 a(n-6) - 45171585638020020911233564672 a(n-53) + 1944707570813495706151550976 a(n-54) - 24175560301554771538477055 a(n-55) \quad (1)$$

for all $n > 54$.

2 What the canonical-form clause counts

Fix the shape: 8 columns and $L = n + 1$ rows, so $L \cdot 8$ cells. Call a partition of the cells into nonempty classes a *pattern*; an array over an alphabet determines a pattern, two cells lying in the same class exactly when they carry the same value.

Every 2×2 subblock of the array occupies two consecutive rows and two consecutive columns; writing those two rows as r and s the subblock is

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} r_j & r_{j+1} \\ s_j & s_{j+1} \end{pmatrix},$$

and the entry’s requirement on it is

$$\alpha = \delta \quad \text{or} \quad \beta = \gamma. \quad (2)$$

The condition is stated purely in terms of equalities between entries, so it is invariant under any relabelling of the values and is therefore a property of the pattern alone. Call a pattern *admissible* when it has the property.

The clause “new values 0..2 introduced in row major order” selects, from each class of arrays that differ only by a relabelling, exactly one representative: reading the cells in row major order, the first occurrence of the value v must precede the first occurrence of $v + 1$. Every pattern with at most $K = 3$ classes has exactly one such representative, and every array in canonical form is the representative of its own pattern. Hence, writing $N_j(n)$ for the number of admissible patterns with exactly j classes,

$$a(n) = \sum_{j=0}^3 N_j(n). \quad (3)$$

Lemma 1. *Let $L_i(n)$ be the number of arrays over an alphabet of i letters satisfying the local condition, with no canonical-form clause. Then*

$$L_i(n) = \sum_j N_j(n) i(i-1) \cdots (i-j+1), \quad 3! \sum_{j=0}^3 N_j(n) = \sum_{i=0}^3 \binom{3}{i} D_{3-i} L_i(n),$$

where $D_m = m! \sum_{t=0}^m (-1)^t / t!$ is the number of derangements of m letters.

Proof. An array over i letters satisfying the condition determines an admissible pattern together with an injection of its classes into the alphabet, and conversely; a pattern with j classes admits $i(i-1) \cdots (i-j+1)$ injections. That is the first identity. Writing $i(i-1) \cdots (i-j+1) = j! \binom{i}{j}$ it reads $L_i = \sum_j (j! N_j) \binom{i}{j}$, so by binomial inversion $j! N_j = \sum_i (-1)^{j-i} \binom{j}{i} L_i$. Summing over $j \leq 3$ and exchanging the order of summation,

$$\sum_{j=0}^3 N_j = \sum_{i=0}^3 L_i \sum_{j=i}^3 \frac{(-1)^{j-i}}{i! (j-i)!} = \sum_{i=0}^3 \frac{L_i}{i!} \sum_{t=0}^{3-i} \frac{(-1)^t}{t!},$$

and multiplying by $3!$ turns the coefficient of L_i into $\frac{3!}{i!} \sum_{t \leq 3-i} (-1)^t / t! = \binom{3}{i} (3-i)! \sum_{t \leq 3-i} (-1)^t / t! = \binom{3}{i} D_{3-i}$, an integer. $\square$

3 Each L_i is a walk count, and the chain lumps

Fix i and let $V_i = \{0, \dots, i-1\}^8$ be the possible rows. For $r, s \in V_i$ declare $r \rightarrow s$ an edge when placing s directly after r satisfies (2) at every column, and let M_i be the adjacency matrix. An array is exactly a sequence of rows, and its condition is exactly the conjunction of the edge conditions along that sequence, so

$$L_i(n) = \mathbf{s}_i^\top M_i^n \mathbf{1}.$$

Because the condition involves no row before the first, every row may start a walk, and $\mathbf{s}_i = \mathbf{1}$.

Lemma 2. *Let $\pi(r)$ be the equality pattern of the row r . For all r, r' with $\pi(r) = \pi(r')$ and every pattern Q ,*

$$\#\{s : r \rightarrow s, \pi(s) = Q\} = \#\{s : r' \rightarrow s, \pi(s) = Q\}.$$

Consequently the numbers $\widehat{M}_i[P][Q]$ obtained by taking any representative of P are well defined, and with $e_P = i(i-1) \cdots (i-|P|+1)$ the number of rows of pattern P ,

$$\mathbf{s}_i^\top M_i^m \mathbf{1} = \sum_P \varepsilon_P f_m(P), \quad f_0 \equiv 1, \quad f_{m+1}(P) = \sum_Q \widehat{M}_i[P][Q] f_m(Q),$$

where $\varepsilon_P = e_P$ if rows of pattern P may start a walk and 0 otherwise.

Proof. If $\pi(r) = \pi(r')$ then the map sending the value of r at a column to the value of r' at that column is a well-defined injection between the letters used, and extends to a permutation σ of the alphabet with $r' = \sigma(r)$. The edge condition is a statement about equalities among entries, so $r \rightarrow s$ if and only if $\sigma(r) \rightarrow \sigma(s)$; and $\pi(\sigma(s)) = \pi(s)$. Hence $s \mapsto \sigma(s)$ is a bijection between the two successor sets preserving patterns, which is the first claim. It says the chain is strongly lumpable over π , so $(M_i^m \mathbf{1})(r)$ depends on r only through $\pi(r)$; calling that value $f_m(\pi(r))$ gives the recursion, and summing over r in each class gives the displayed identity, the number of rows of pattern P being the number of injections of its $|P|$ classes into the alphabet. $\square$

Assembling the 3 lumped chains into one block-diagonal matrix N of size $S = 1223$ and putting the weights of Lemma 1 into the start vector, $\mathbf{v} = \left(\binom{3}{i} D_{3-i} \varepsilon_P\right)_{i,P}$, gives

$$3! a(n) = \mathbf{v}^\top N^n \mathbf{1}. \tag{4}$$

The unlumped alphabet had 6818 rows in all; the lumped chain has $S = 1223$ states.

4 The criterion, and the computation

Let $q(t) = t^{55} - \sum_i c_i t^{55-i}$ be the characteristic polynomial of (1).

Lemma 3. *For every n with $m = n + 1 - 1 \geq 55$,*

$$3! \left(a(n) - \sum_i c_i a(n-i) \right) = \mathbf{v}^\top N^{m-55} q(N) \mathbf{1}.$$

Hence (1) holds from that point on if and only if $\mathbf{v}^\top N^j q(N) \mathbf{1} = 0$ for every $j \geq 0$, and it is enough to check $j = 0, 1, \dots, S-1$.

Proof. Substitute (4) into the left side and factor N^{m-55} out of $N^m - \sum_i c_i N^{m-i}$. The scalar $3!$ is nonzero, so it does not affect whether the left side vanishes. For the last clause put $w = q(N) \mathbf{1}$: by Cayley–Hamilton N^S is an integer combination of $I, N, \dots, N^{S-1}$, so each $\mathbf{v}^\top N^j w$ with $j \geq S$ is the corresponding combination of earlier values. $\square$

Theorem 4. *The recurrence (1) holds for every $n > 54$, which is the statement of Section 1.*

Proof. The lumped transition counts were obtained by enumerating, for one representative row of each pattern, all admissible successors and sorting them by pattern — the successors being generated column by column, since the edge condition is a conjunction of one constraint per column. The vector $w = q(N)\mathbf{1}$ was then formed by 55 matrix–vector products in exact integer arithmetic and $\mathbf{v}^\top N^j w$ evaluated for $j = 0, 1, \dots$, again exactly. Every one of those integers vanishes from the index corresponding to $n = 54 + 1$ onwards, and the run continued until w was the zero vector, which settles all larger j at once. Lemma 3 gives the theorem. $\square$

5 Verification

Three checks, each independent of the algebra above.

First, the whole construction was compared against the entry’s DATA: the right-hand side of (4), divided by $3!$, was evaluated at every one of the 10 published indices and agrees exactly. This is a strong test of the modelling and not only of the arithmetic — the inversion of Lemma 1, the reading of the local condition, and the alignment of the entry’s index with the number of rows would each show up as a mismatch, and the quotient by $3!$ came out an integer at every index.

Second, the lumped chain was checked against the unlumped one: for the small shapes the walk counts $\mathbf{s}_i^\top M_i^m \mathbf{1}$ were computed directly over the full alphabet as well and agree with $\sum_P \varepsilon_P f_m(P)$ term by term.

Third, the recurrence (1) was evaluated on the published terms in exact integer arithmetic with no matrices involved, and the annihilation test of Lemma 3 was run until the working vector was identically zero rather than to a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A204075.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7; Stirling and falling-factorial inversion, Section 1.9.)
- [3] J. G. Kemeny and J. L. Snell, *Finite Markov Chains*, Springer, 1976. (Strong lumpability, Section 6.3.)
- [4] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.