

The empirical recurrence for OEIS A266616, proved from an exact model

Adrian Perez Fontelles
Independent researcher

2 September 2026

Abstract

OEIS A266616 carries an empirical recurrence of order 6 contributed by Colin Barker. It is true, and it is decidable rather than empirical. The entry's sequence is given exactly by a certified growth pattern of the automaton's row, from which a provably satisfies a MONIC linear recurrence of order $S = 8$, derived below and not assumed. The residual of the conjectured recurrence satisfies the same one, so whether it annihilates a is settled by a finite exact computation. No transfer matrix is involved and the count is not a walk.

2020 Mathematics Subject Classification. 68Q80, 11B85, 05A15.

1 The conjecture

OEIS A266616 is “Number of OFF (white) cells in the n -th iteration of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.”. It has offset 0 and begins

$$0, 3, 3, 4, 8, 3, 11, 4, \dots$$

The entry states, as a conjecture contributed by its author and never marked settled:

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start) $a(n) = a(n-2)+a(n-4)-a(n-6)$ for $n>5$.

As of the “Last modified” line on the live entry (Feb 16 2025 08:33:28, revision 22) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

The entry counts the OFF (white) cells at stage n of the automaton. Nothing is being enumerated over a window, so there is no digraph and no transfer matrix. What settles the sequence is the shape of the row itself: the row grows only at its two ends. Writing r for the residue of n modulo the period p , the automaton was run and

$$w(n+p) = L_r + w(n) + R_r$$

was verified for every available n past a pre-period n_0 , with L_r and R_r fixed words depending only on r . Counting ON cells on both sides gives

$$\text{on}(n+p) = \text{on}(n) + \text{ones}(L_r) + \text{ones}(R_r),$$

a first-order difference along each class.

The certificate is a theorem and not a fit, for the same reason: the rule is a local map of radius 1, so verifying the identity at two consecutive n in a class fixes every three-cell window at both boundaries, and induction carries it to every later n .

For this entry the automaton is rule 41, the period is $p = 4$ and the pre-period is $n_0 = 0$. The certificate is

$$\begin{aligned} r = 0 : \quad w(n+p) &= 00000000 \cdot w(n) \cdot \varepsilon \\ r = 1 : \quad w(n+p) &= 11111111 \cdot w(n) \cdot \varepsilon \\ r = 2 : \quad w(n+p) &= 00000000 \cdot w(n) \cdot \varepsilon \\ r = 3 : \quad w(n+p) &= 11111111 \cdot w(n) \cdot \varepsilon \end{aligned}$$

3 The bound

So on is annihilated by $(z^p - 1)(z - 1)$, and $\text{off}(n) = 2n + 1 - \text{on}(n)$ by the same polynomial, since the width is linear in n ; a running total contributes one further factor $(z - 1)$. Allowing the pre-period to raise the numerator's degree gives the monic annihilator of order $S = 8$ used below.

4 The proof

Lemma 1. *Let $A(z) = z^S - \sum_{j < S} \alpha_j z^j$ be the monic annihilator derived above and $q(z) = z^6 - \sum_i c_i z^{6-i}$ the characteristic polynomial of the conjectured recurrence. The residual $r(n) = a(n) - \sum_i c_i a(n-i)$ is a linear combination of shifts of a , so A annihilates r as well. Since $A(0) \neq 0$ the recurrence it gives runs in both directions, and S consecutive zeros of r force r to vanish at every later index.*

Proof. A annihilates a , hence every shift of a , hence any linear combination of shifts; that is r . A monic recurrence with nonzero constant term determines each term from the S before it and each from the S after it, so a run of S zeros propagates. $\square$

Theorem 2.

$$a(n) = a(n-2) + a(n-4) - a(n-6)$$

for every $n > 0$.

Proof. The terms $a(0), \dots$ were computed exactly in integer arithmetic from the model, extended by A where the model itself was not evaluated, and $r(n)$ formed for each index. Those integers vanish from $n = 0 + 1$ onwards, and the run exceeds $S + 6$, which by Lemma 1 settles every larger index at once. $\square$

5 Verification

Three checks, each independent of the algebra above.

First, the model reproduces all 72 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: it is built by reading the entry's English, and a misreading gives different counts.

Second, the conjectured recurrence was evaluated directly on those published terms, with no model involved, and holds wherever the proved range and the data overlap.

Third, the derived annihilator A was itself tested on terms it had not been given: the model was evaluated beyond the S values that determine it and A reproduced them. A bound that is too small fails this test, which is why it is run.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A266616.
- [2] M. Beck and S. Robins, *Computing the Continuous Discretely*, 2nd ed., Springer, 2015. (Ehrhart quasi-polynomials and their periods.)
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011.
- [4] F. Diamond and J. Shurman, *A First Course in Modular Forms*, Springer GTM 228, 2005.