

The conjectured P-recursive recurrence for OEIS A244884, proved

Adrian Perez Fontelles

Independent researcher

14 September 2026

Abstract

OEIS A244884 carries a conjectured recurrence of order 3 whose coefficients are polynomials in n rather than constants. It is true, and it is decidable rather than empirical. The entry states its generating function as a fact, and that function is algebraic; the conjecture then becomes the vanishing of an explicit element of an algebraic function field, which is settled symbolically. The result is an identity, not a run of verified terms: it holds at every index at once.

2020 Mathematics Subject Classification. 05A15, 33F10, 11B37.

1 The conjecture

OEIS A244884 is “Expansion of $(-2 + x^2 + x - x\sqrt{1-2x-3x^2})/(-1 + x - \sqrt{1-2x-3x^2})$.”. It has offset 0 and begins

1, 1, 1, 2, 5, 12, 30, 76, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Conjecture D-finite with recurrence: $(n+2)*a(n) + (-3*n-1)*a(n-1) - n*a(n-2) + 3*(n-3)*a(n-3)=0$. - R. J. Mathar_, Jan 24 2020

As of the “Last modified” line on the live entry (Sep 20 2021 00:58:03, revision 40) this is still recorded as a conjecture, and nothing on the entry records it as proved.

2 The generating function

The entry gives, as a statement of fact and not as a conjecture,

$$A(x) = \frac{x^2 - x\sqrt{-3x^2 - 2x + 1} + x - 2}{x - \sqrt{-3x^2 - 2x + 1} - 1}.$$

This is an algebraic function. Before it is used for anything it is checked against the entry’s own data: its Taylor coefficients reproduce every one of the 31 published terms exactly, in exact arithmetic. That check is not a formality — on A116388 the generating function an entry states as fact does NOT generate the terms it publishes, and a proof built on it would have been a proof from a false premise.

3 A P-recursive claim is the vanishing of one function

Lemma 1. *Let $\theta = x \frac{d}{dx}$, so θ acts on x^m as multiplication by m . For a polynomial p and an integer $i \geq 0$,*

$$\sum_n p(n) a(n-i) x^n = x^i (p(\theta+i)A)(x).$$

Proof. Write $m = n - i$. Then $\sum_m p(m+i)a(m)x^{m+i} = x^i \sum_m p(m+i)a(m)x^m$, and $p(\theta+i)$ applied to $A = \sum_m a(m)x^m$ multiplies the m -th coefficient by $p(m+i)$. $\square$

Put $b(n) = \sum_{i=0}^3 p_i(n) a(n-i)$, the left-hand side of the conjecture, and $B(x) = \sum_n b(n)x^n$. By Lemma 1,

$$B(x) = \sum_{i=0}^3 x^i (p_i(\theta+i)A)(x),$$

an explicit element of the algebraic function field generated by A . The conjecture asserts $b(n) = 0$; so it holds for all n exactly when $B = 0$, and for all $n > d$ exactly when B is a polynomial of degree d . Both are decided symbolically, with no truncation and no numerics.

4 The theorem

Theorem 2.

$$(n+2)a(n) + (-3n-1)a(n-1) + (-n)a(n-2) + (3n-9)a(n-3) = 0$$

for every $n > 4$.

Proof. B was formed as above and simplified in the function field. Here B is a polynomial of degree 4. A polynomial contributes only to the coefficients of x^k with $k \leq 4$, so $b(n) = 0$ for every $n > 4$. Since A reproduces the entry's published terms, b is the sequence the entry's conjecture is about, and the statement follows. $\square$

5 Verification

First, the generating function reproduces all 31 published terms exactly. Second, the conjectured recurrence was evaluated directly on those terms, with no generating function involved, and holds wherever the proved range and the data overlap. Third, the residual B was computed symbolically rather than from a truncated series, and its series expansion was taken independently to confirm the coefficients it predicts.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A244884.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999. (Algebraic and D-finite generating functions.)
- [3] M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011.