

A condition on the multiset of every subblock: OEIS A184201

Adrian Perez Fontelles

Independent researcher

7 September 2026

Abstract

OEIS A184201 carries an empirical recurrence of order 10 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. The entry's condition on a subblock depends only on which values the block holds and how many times, not on where they sit, so it is decided by the few consecutive lines the block spans. The admissible windows are the vertices of a finite digraph and the arrays counted are exactly its walks; hence $a(n)$ is a walk count on $S = 10$ vertices and is C -finite, and whether the conjectured recurrence annihilates it is settled by a finite exact computation, with the Cayley–Hamilton theorem bounding the work.

2020 Mathematics Subject Classification. 05A15, 05B45, 15B36.

1 The conjecture

OEIS A184201 is “Number of $(n+1) \times 4$ binary arrays with no 2×2 subblock containing fewer than two 1s.”. It has offset 1 and begins

108, 1028, 8969, 80812, 721939, 6468501, 57909336, 518574936, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 9a(n-1) + 8a(n-2) - 79a(n-3) + 19a(n-4) + 134a(n-5) - 54a(n-6) - 56a(n-7) + 23a(n-8) + 5a(n-9) - 2a(n-10)$

As of the “Last modified” line on the live entry (Oct 03 2025 02:24:02, revision 9) this is still recorded as empirical, and nothing on the entry records it as proved.

2 The count is a walk count

The entry counts arrays with 4 columns over the alphabet $\{0, 1\}$, growing one row at a time, and asks that every 2×2 subblock holds at least 2 entries equal to 1.

That condition does not depend on where in the block a value sits, only on which values the block holds and how many times — it is a function of the block's multiset of entries. In particular it is decided by 2 consecutive rows, the ones the block spans, so carrying the previous 1 row as the state is enough: each new row completes every block that ends on it, and no judgement is left over at the end of the array.

Merging vertices with identical futures leaves $S = 10$. An admissible array is exactly a walk, so with M the adjacency matrix and ι, τ the vectors recording which states may begin and end an array,

$$a(n) = \iota^\top M^n \tau,$$

the exponent fixed by the entry's own shape and checked against its published terms. In particular a is C -finite.

3 The proof

Lemma 1. *Let $q(t) = t^{10} - \sum_i c_i t^{10-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+10} - \sum_i c_i M^{j+10-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. A constant factor in front of a divides out of the residual and changes nothing here. $\square$

Theorem 2.

$$a(n) = 9a(n-1) + 8a(n-2) - 79a(n-3) + 19a(n-4) + 134a(n-5) - 54a(n-6) - 56a(n-7) + 23a(n-8) + 5a(n-9)$$

for every $n > 9$.

Proof. The vector $w = q(M)\tau$ was formed by 10 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 9 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

4 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 17 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: the digraph is built by reading the entry’s English, and a misreading gives different counts.

Second, the reading was pinned before the digraph was written, by a program that enumerates every array of the given shape one by one, forms each subblock directly and tests the entry’s condition on it. That program shares no code with the transfer matrix, so an error in one does not hide an error in the other.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 1 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A184201.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.