

The empirical recurrence for OEIS A183340, proved by transfer matrix

Adrian Perez Fontelles

Independent researcher

3 September 2026

Abstract

OEIS A183340 counts arrays of width 8 over $\{0,1\}$ under local conditions relating each cell to its neighbours or to the cells preceding it. The entry carries an empirical recurrence of order 32 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical: every condition is settled inside three consecutive array rows, so the count is a walk count on $S = 2081$ states.

2020 Mathematics Subject Classification. 05A15, 05C30, 15A18.

1 The conjecture

OEIS A183340 is “Number of $n \times 8$ binary arrays with each 1 adjacent to exactly one 1 vertically and one 1 horizontally.”. It has offset 1 and begins

1, 19, 63, 193, 1005, 4133, 16029, 68662, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = a(n-1) + 4a(n-2) + 33a(n-3) + 40a(n-4) - 27a(n-5) - 190a(n-6) - 384a(n-7) + 149a(n-8) + 768a(n-9) + 398a(n-10) - 281a(n-11) + 312a(n-12) - 879a(n-13) - 1871a(n-14) + 351a(n-15) + 2551a(n-16) - 928a(n-17) + 499a(n-18) + 267a(n-19) - 947a(n-20) - 295a(n-21) - 55a(n-22) + 284a(n-23) + 192a(n-24) + 136a(n-25) + 98a(n-26) - 86a(n-27) + 17a(n-28) - 4a(n-29) - 19a(n-30) + 3a(n-31) - 12a(n-32)$

As of the “Last modified” line on the live entry (Oct 03 2025 01:35:00, revision 9) this is still recorded as empirical, and nothing on the entry records it as proved.

2 What is being counted

The array has 8 cells across, with entries in $\{0,1\}$. The NEIGHBOURS of a cell are the cells immediately to its left, to its right, above it and below it, those that lie inside the array. The entry asks that

- every cell carrying 1 has exactly 1 of its two VERTICAL neighbours carrying 1 and exactly 1 of its two HORIZONTAL neighbours carrying 1;

and nothing else.

3 Three rows decide everything

Lemma 1. *Every clause above is decided by three consecutive array rows. An adjacency clause for a cell of row i reads rows $i - 1$, i and $i + 1$; a precedence clause for a cell of row i reads rows $i - 2$, $i - 1$ and i , and columns at most two to the left.*

Proof. The neighbours of a cell lie one step away in each of the four directions, and the predecessors one step to the left or one step up; a clause naming two cells in a row or column reaches two steps, never more. $\square$

So the state is the pair (row above, current row), with a row not yet written carried as $\perp$. A step appends a row: the precedence clauses for the NEW row are settled at once, because everything they read is already written, and the adjacency clauses for the MIDDLE row of the window are settled because the row below it has just arrived. An adjacency clause for a row is settled only when the row BELOW it has been written, so the condition for the LAST row of an array belongs to the end vector, which asks whether that row is satisfied with nothing below it. The end vector is therefore not the all-ones vector: a row satisfied only because a further row is written below it does not finish an array. With M the adjacency matrix of the digraph on those states,

$$a(n) = \iota^\top M^j \tau, \quad S = 2081.$$

The walk index j and the entry's own n differ by a constant, read off by matching the published terms rather than assumed. In particular a is C -finite.

4 The proof

Lemma 2. *Let $q(t) = t^{32} - \sum_i c_i t^{32-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+32} - \sum_i c_i M^{j+32-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. $\square$

Theorem 3.

$$a(n) = a(n-1) + 4a(n-2) + 33a(n-3) + 40a(n-4) - 27a(n-5) - 190a(n-6) - 384a(n-7) + 149a(n-8) + 768a(n-9)$$

for every $n > 31$.

Proof. The vector $w = q(M)\tau$ was formed by 32 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 31 + 1$ onwards and the run continued until the working vector was identically zero. $\square$

5 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 23 terms the entry publishes, exactly, in integer arithmetic.

Second, the count was recomputed for the smallest arrays straight from the definition: enumerate every array of that size in the orientation the entry's own name writes and test each

clause on each cell directly. No window, no state and no transfer matrix are involved, and the published terms came back.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 2 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A183340.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.