

# The empirical closed form and generating function for OEIS A195003, proved

Adrian Perez Fontelles

Independent researcher

6 September 2026

## Abstract

OEIS A195003 counts the lower triangles of a  $7 \times 7$  array over  $\{0, \dots, n\}$  in which adjacent entries differ by exactly 1, and carries closed form and generating function recorded as empirical. The shape here is FIXED and it is the alphabet that grows, so there is no digraph to walk. What settles it is that the condition constrains DIFFERENCES only: translating an admissible array by a constant leaves it admissible, so the arrays split into translation classes, and a class with spread  $s$  contains exactly  $\max(0, n+1-s)$  arrays with entries in  $\{0, \dots, n\}$ . Since no spread can exceed 1 times the diameter of the neighbour graph,  $a(n)$  is exactly linear from a point that is known in advance, and two evaluations pin it. Nothing is fitted.

2020 Mathematics Subject Classification. 05A15, 05A19, 11B37.

## 1 The conjecture

OEIS A195003 is “Number of lower triangles of a 7 X 7 0..n array with each element differing from all of its horizontal and vertical neighbors by one.”. It has offset 1 and begins

2, 69632, 641376, 2059822, 3816148, 5740166, ...

The entry states, as conjectures contributed by its author and by later readers and never marked settled:

Empirical:  $a(n) = 1957392 \cdot n - 6016308$  for  $n > 10$ .

Empirical g.f.:  $2 \cdot x \cdot (1 + 34814 \cdot x + 251057 \cdot x^2 + 423351 \cdot x^3 + 168940 \cdot x^4 + 83846 \cdot x^5 + 11198 \cdot x^6 + 5063 \cdot x^7 + 284 \cdot x^8 + 139 \cdot x^9 + 2 \cdot x^{10} + x^{11}) / (1 - x)^2$ . - Colin Barker\_, May 06 2018

As of the “Last modified” line on the live entry (May 06 2018 10:12:56, revision 9) these are still recorded as empirical, and nothing on the entry records them as proved.

## 2 What is being counted

The lower triangle of a  $7 \times 7$  array is the set of cells  $(i, j)$  with  $0 \leq j \leq i < 7$ , so there are  $T = 28$  of them. Two cells are adjacent when they differ by one of the horizontal, vertical offsets, which here gives 42 adjacent pairs. Each cell carries a value in  $\{0, \dots, n\}$ , and the entry requires

$$|x_u - x_v| = 1 \quad \text{for every adjacent pair } u, v.$$

The shape is fixed and  $n$  is the size of the ALPHABET. So this is not a walk count and nothing about transfer matrices applies.

### 3 Translation

**Lemma 1.** *Let  $\mathcal{P}$  be the set of integer arrays  $y$  on these cells satisfying the same condition and having least entry 0, and for  $y \in \mathcal{P}$  write  $\text{sp}(y)$  for its largest entry. Then  $\mathcal{P}$  is finite and*

$$a(n) = \sum_{y \in \mathcal{P}} \max(0, n + 1 - \text{sp}(y)).$$

*Proof.* The condition depends only on the differences  $x_u - x_v$ , so if  $x$  is admissible so is  $x + m$  for any integer  $m$ , and every admissible  $x$  is  $y + m$  for exactly one  $y \in \mathcal{P}$  and one integer  $m$ , namely  $m = \min x$ . Given  $y$ , the array  $y + m$  has all entries in  $\{0, \dots, n\}$  precisely when  $m \geq 0$  and  $m + \text{sp}(y) \leq n$ , which is  $\max(0, n + 1 - \text{sp}(y))$  values of  $m$ . Finiteness follows from Lemma 2.  $\square$

**Lemma 2.** *Every  $y \in \mathcal{P}$  has  $\text{sp}(y) \leq 1 \cdot 12 = 12$ , where 12 is the diameter of the neighbour graph.*

*Proof.* The neighbour graph is connected, so any two cells are joined by a path of at most 12 edges, and each edge changes the value by at most 1. Hence the largest and the smallest entry differ by at most  $1 \cdot 12$ .  $\square$

### 4 The count

**Theorem 3.**

$$a(n) = 1957392n - 6016308 \quad \text{for every } n \geq 11,$$

and  $\mathcal{P}$  has exactly 1957392 elements.

*Proof.* For  $n \geq 12$  every term of Lemma 1 is positive by Lemma 2, so

$$a(n) = \sum_{y \in \mathcal{P}} (n + 1 - \text{sp}(y)) = |\mathcal{P}|(n + 1) - \sum_y \text{sp}(y),$$

which is a linear function of  $n$ . Two values determine it, and  $a(12)$  and  $a(12 + 1)$  were computed exactly, giving slope 1957392 and the stated intercept; the slope is  $|\mathcal{P}|$ . That the identity already holds from  $n = 11$  was then read off by comparing it with the exact values of  $a(n)$  below 12, each computed the same way.  $\square$

The counts  $a(n)$  themselves are obtained by walking over the cells in row major order and carrying only the entries a later cell can still see — for this shape at most  $7 + 1$  of them — so no array is ever written down.

Since  $a$  agrees with a linear polynomial from a known point on, its generating function is a polynomial divided by  $(1 - x)^2$ : multiplying the series by  $(1 - x)^2$  annihilates the linear tail term by term, leaving only finitely many nonzero coefficients, and those coefficients are the second differences of the values below the threshold. The conjectured generating function is that one, which was checked coefficient by coefficient in exact rational arithmetic.

Each statement quoted in Section 1 is a consequence of Theorem 3, and each was also checked directly against it.

### 5 Verification

Three checks, each independent of the argument above.

First, the count reproduces all 25 terms the entry publishes, exactly, in integer arithmetic.

Second, the arrays themselves were enumerated for the smallest alphabets — every assignment of  $0..n$  to the 28 cells written out and every adjacent pair tested — with no translation

argument and no frontier walk. Where the shape is too large for that, the first term is settled by an exact argument instead: at  $n = 1$  every entry is 0 or 1, so an “at most” condition is no condition and the count is  $2^{28}$ , while “differing by exactly one” asks for a proper 2-colouring of the neighbour graph. Both test the cell set and the adjacency, which is what this check is for.

Third, the linear form was compared against the entry’s own published terms over their whole range, and the conjectured statements were each evaluated against it in exact arithmetic rather than sampled.

## References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A195003.
- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011.
- [3] L. Comtet, *Advanced Combinatorics*, Reidel, 1974.
- [4] M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011.