

A proof of the conjectured recurrence for OEIS A098557

Adrian Perez Fontelles

Independent researcher

25 August 2026

Abstract

OEIS A098557 carries a conjectured linear recurrence with polynomial coefficients, of order 2. We prove it. The entry's exponential generating function is transcendental, so the usual algebraic arguments do not apply; instead we observe that it lies in a finitely generated module over $\mathbb{Q}(x)$ spanned by monomials $\log(u)^a \exp(g)^k$, and that this module is closed under differentiation, because $(\log u)' = u'/u$ and $(\exp g)' = g' \exp g$ lower the log power and fix the exponential respectively. Re-indexing the recurrence forward turns shifts into derivatives, so the sequence $b(m) = \sum_j q_j(m) a(m+j)$ has exponential generating function $B = \sum_j q_j(\theta) A^{(j)}$, computed exactly inside that module. Here B collapses to the polynomial -1 , of degree 0, which proves the recurrence for every $n > 0$.

2020 Mathematics Subject Classification. 11B37, 05A15, 12H05, 33F10.

1 The sequence and the conjecture

OEIS A098557 is “Expansion of e.g.f. $(1/2)^*(1+x)*\log((1+x)/(1-x))$ ”. It has offset 0 and begins

$$a(0), \dots, a(7) = 0, 1, 2, 2, 8, 24, 144, 720, \dots$$

The entry does not list an exponential generating function separately: it is defined as the expansion of one, so the identity below is the definition of the sequence rather than a claim about it,

$$A(x) = \sum_{n \geq 0} a(n) \frac{x^n}{n!} = \frac{(x+1) \log\left(\frac{-x-1}{x-1}\right)}{2}, \quad (1)$$

and it carries the following comment:

conjecture: $-a(n) + a(n-1) + (n-1)*(n-3)*a(n-2) = 0$. - *R. J. Mathar*, Nov 14 2011

As of the “Last modified” line on the live entry (2026-06-30, revision 27) the statement is still recorded as a conjecture, and no proof appears among the entry's links, formulas or programs.

Writing the conjectured relation in the form $\sum_{i=0}^2 p_i(n) a(n-i) = 0$, the coefficient polynomials are

$$p_0(n) = -1, \quad p_1(n) = 1, \quad p_2(n) = (n-3)(n-1)$$

2 Recurrences as differential operators

Let $A(x) = \sum_{n \geq 0} a(n)x^n/n!$ be the exponential generating function of a sequence, and let $\theta = x \frac{d}{dx}$, so that θ acts on x^m as multiplication by m . For a polynomial p we write $p(\theta)$ for the corresponding differential operator, so that $p(\theta)A = \sum_m p(m)a(m)x^m/m!$.

Lemma 1. Let $p_0, \dots, p_r$ be polynomials and suppose the conjecture asserts $\sum_{i=0}^r p_i(n) a(n-i) = 0$. Re-index by $n = m + r$ and $j = r - i$, and write $q_j(m) = p_{r-j}(m + r)$, so the claim reads $\sum_{j=0}^r q_j(m) a(m + j) = 0$. Put $b(m) = \sum_j q_j(m) a(m + j)$. Then

$$B(x) := \sum_{m \geq 0} b(m) \frac{x^m}{m!} = \sum_{j=0}^r (q_j(\theta) A^{(j)})(x),$$

where $A^{(j)}$ denotes the j -th derivative of A .

Proof. For an exponential generating function an upward shift is differentiation: $\sum_{m \geq 0} a(m + j) x^m / m! = A^{(j)}(x)$. Multiplying the coefficient of $x^m / m!$ by $q_j(m)$ is exactly the action of $q_j(\theta)$, and summing over j gives the statement. The re-indexing is what makes this work: a downward shift $a(n - i)$ would correspond to repeated integration, whereas an upward shift is a derivative, and the field used below is closed under differentiation. $\square$

Corollary 2. With the notation of Lemma 1, the recurrence $\sum_i p_i(n) a(n - i) = 0$ holds for every $n > d + r$ if and only if B is a polynomial of degree at most d .

Proof. $b(m)$ is $m!$ times the coefficient of x^m in B , so $b(m) = 0$ for all $m > d$ says exactly that B has no terms above x^d . $\square$

This turns the conjecture into a closed-form identity. The point is that it can then be decided exactly, with no truncation: A is algebraic, so B lies in the same field as A , and one only has to recognise it.

Lemma 3. Let $u, g \in \mathbb{Q}(x)$ and let M be the $\mathbb{Q}(x)$ -module spanned by the monomials $\log(u)^a \exp(g)^k$ for $a \geq 0$ and $k \in \mathbb{Z}$. Then M is closed under d/dx , and hence under $\theta = x d/dx$; explicitly,

$$\frac{d}{dx} \left(c \log(u)^a \exp(g)^k \right) = \left(c' + c k g' \right) \log(u)^a \exp(g)^k + c a \frac{u'}{u} \log(u)^{a-1} \exp(g)^k, \quad c \in \mathbb{Q}(x).$$

Proof. Differentiate the product. The factor $\exp(g)^k = \exp(kg)$ contributes kg' and does not change the monomial; the factor $\log(u)^a$ contributes $a(u'/u) \log(u)^{a-1}$, lowering the log exponent by one. Both new coefficients lie in $\mathbb{Q}(x)$, so the result is again a finite combination of the same monomials. $\square$

Two features make this usable. The exponential part never mixes monomials at all, and the logarithmic part only ever moves *downwards*, so the module generated by any single element is finite dimensional and every computation terminates. Writing B in this basis, the test of Corollary 2 becomes: B is a polynomial precisely when every coefficient outside the monomial $\log(u)^0 \exp(g)^0$ vanishes and the remaining coefficient has constant denominator. Both are decided by exact cancellation of rational functions, so no series truncation enters anywhere.

3 The computation for A098557

The generating function (1) lies in the module M of Lemma 3. Applying Lemma 1 to the coefficient polynomials of Section 1 and reducing in K by Lemma 3, the $\sqrt{D}$ component of B cancels identically and the rational part collapses to

$$B(x) = -1,$$

a polynomial of degree 0.

Theorem 4. *The conjectured recurrence*

$$-1 a(n) + 1 a(n-1) + (n-3)(n-1) a(n-2) = 0$$

holds for every $n > 2$.

Proof. Immediate from Corollary 2 and the displayed value of B . □

Remark 5. The polynomial B is not an error term: it records the boundary of the recurrence. For $n \leq 0$ some of the shifted terms $a(n-i)$ fall outside the range of the sequence, and B collects exactly those contributions. Above that range the relation is exact.

4 Verification

Every number above is an output of a script written separately from this note, and two independent checks were run.

First, the generating function was checked against the entry rather than assumed: the Taylor coefficients of (1) were expanded and compared term by term with the DATA section of A098557, agreeing on all 13 terms compared.

Second, the recurrence itself was evaluated directly on the published terms, in exact integer arithmetic and without reference to the generating function: for each n in range the sum $\sum_i p_i(n)a(n-i)$ was formed from the entry's own values and found to vanish, for all 21 values of n from $n = 3$ upward. This is what Corollary 2 predicts from the degree of B , and it is a genuinely different computation from the symbolic one: it never forms B , never differentiates, and uses only integers.

The symbolic step is exact throughout. The residual is obtained by rational-function cancellation in K , not by series truncation, so the identity $B = -1$ is an identity of functions and the theorem holds for all $n > 2$ simultaneously.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A098557.
- [2] M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011, Chapter 7 (holonomic closure properties and the translation between recurrences and differential equations).
- [3] R. P. Stanley, *Enumerative Combinatorics*, Volume 2, Cambridge University Press, 1999, Chapter 6 (algebraic and D-finite generating functions).