

Parities, differences and neighbours: OEIS A229400

Adrian Perez Fontelles

Independent researcher

7 September 2026

Abstract

OEIS A229400 carries an empirical recurrence of order 32 contributed by R. H. Hardin. It is true, and it is decidable rather than empirical. The entry's condition is decided by a bounded amount of state carried down the array; the admissible windows are the vertices of a finite digraph and the arrays counted are exactly the walks in it. Hence $a(n)$ is a walk count on $S = 33$ vertices and is C -finite, and whether the conjectured recurrence annihilates it is settled by a finite exact computation, with the Cayley–Hamilton theorem bounding the work.

2020 Mathematics Subject Classification. 05A15, 05B45, 15B36.

1 The conjecture

OEIS A229400 is “Number of $n \times 6$ 0..2 arrays with top left element 0, horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and antidiagonal differences never 0.”. It has offset 1 and begins

32, 233, 1881, 15952, 138645, 1220881, 10826489, 96353860, ...

The entry states, as a conjecture contributed by its author and never marked settled:

Empirical: $a(n) = 33*a(n-1) - 496*a(n-2) + 4675*a(n-3) - 31879*a(n-4) + 170672*a(n-5) - 754133*a(n-6) + 2833601*a(n-7) - 9205272*a(n-8) + 26062859*a(n-9) - 64494435*a(n-10) + 139473436*a(n-11) - 263178485*a(n-12) + 432422025*a(n-13) - 617475384*a(n-14) + 765111259*a(n-15) - 821876951*a(n-16) + 765111259*a(n-17) - 617475384*a(n-18) + 432422025*a(n-19) - 263178485*a(n-20) + 139473436*a(n-21) - 64494435*a(n-22) + 26062859*a(n-23) - 9205272*a(n-24) + 2833601*a(n-25) - 754133*a(n-26) + 170672*a(n-27) - 31879*a(n-28) + 4675*a(n-29) - 496*a(n-30) + 33*a(n-31) - a(n-32)$.

As of the “Last modified” line on the live entry (Sep 07 2022 11:33:58, revision 6) this is still recorded as empirical, and nothing on the entry records it as proved.

2 The count is a walk count

The entry counts arrays with 6 columns over the alphabet $\{0, 1, 2\}$, growing one row at a time, and asks that the top left entry is 0, and the step to the right may never change the entry by 1 modulo 3, the step downwards may never change the entry by -1 modulo 3, the step antidiagonally may never change the entry by 0 modulo 3.

Every cell the condition names lies within one row of the cell being judged, so three consecutive rows decide everything anchored at the middle one. Carry two consecutive rows as the state; a row is judged when the row after it arrives, and the row with nothing after it is judged

by the end vector. A neighbour off the edge of the array is not a neighbour, which is what the entry means and what makes the first and last rows behave differently from the rest.

Merging vertices with identical futures leaves $S = 33$. An admissible array is exactly a walk, so with M the adjacency matrix and ι, τ the vectors recording which states may begin and end an array,

$$a(n) = \iota^\top M^{n-1} \tau,$$

the exponent fixed by the entry's own shape and checked against its published terms. In particular a is C -finite.

3 The proof

Lemma 1. *Let $q(t) = t^{32} - \sum_i c_i t^{32-i}$ be the characteristic polynomial of the conjectured recurrence. The residual $a(n) - \sum_i c_i a(n-i)$ equals $\iota^\top M^j q(M) \tau$ for the corresponding walk index j , so the recurrence holds from some point on if and only if $\iota^\top M^j q(M) \tau = 0$ for all large j , and it is enough to examine $j < S$.*

Proof. Factor M^j out of $M^{j+32} - \sum_i c_i M^{j+32-i}$. For the bound, M^S is an integer combination of $I, M, \dots, M^{S-1}$ by Cayley–Hamilton, so the numbers $u_j = \iota^\top M^j q(M) \tau$ obey the monic recurrence given by the characteristic polynomial of M ; S consecutive zeros force every later one, and the last nonzero u_j pins the threshold exactly. A constant factor in front of a divides out of the residual and changes nothing here. $\square$

Theorem 2.

$$a(n) = 33a(n-1) - 496a(n-2) + 4675a(n-3) - 31879a(n-4) + 170672a(n-5) - 754133a(n-6) + 2833601a(n-7)$$

for every $n > 32$.

Proof. The vector $w = q(M)\tau$ was formed by 32 matrix–vector products in exact integer arithmetic and $\iota^\top M^j w$ evaluated for $j = 0, 1, \dots$, again exactly; those integers vanish from the index corresponding to $n = 32 + 1$ onwards and the run continued until the working vector was identically zero, which settles every larger j at once. $\square$

4 Verification

Four checks, each independent of the algebra above.

First, the model reproduces all 18 terms the entry publishes, exactly, in integer arithmetic. This is what ties the model to the entry: the digraph is built by reading the entry's English, and a misreading gives different counts.

Second, the reading was pinned before the digraph was written, by a program that enumerates every array of the given shape one by one and tests the entry's condition on it directly. That program shares no code with the transfer matrix, so an error in one does not hide an error in the other.

Third, the conjectured recurrence was evaluated directly on the published terms, with no matrices involved, and holds wherever the proved range and the data overlap.

Fourth, the annihilation test of Lemma 1 was carried out in exact integer arithmetic on the whole vector rather than on a sampled prefix.

References

- [1] The OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026. Sequence A229400.

- [2] R. P. Stanley, *Enumerative Combinatorics*, Volume 1, 2nd ed., Cambridge University Press, 2011. (Transfer-matrix method, Section 4.7.)
- [3] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [4] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013.