Haemocytes of Crassostrea virginica as a dual-purpose source of ultra-long DNA for genome scaffolding and environmental DNA surveillance

Alyssa Strickland[1], Kshitij Amar[2], Bo-Young Lee[1], Caylin Grove[1], Miten Jain[2], Bonnie L. Brown[1*]

[1*]Department of Biological Sciences, University of New Hampshire, Durham, 03874, NH, USA.
[2]Department of Bioengineering, Northeastern University, Boston, 02115, MA, USA.

*Corresponding author(s). E-mail(s): bonnie.brown@unh.edu;

Abstract

Sequencing and assembling ultra-long DNA (ULDNA) scaffolds can extend utility of the existing eastern oyster *Crassostrea virginica*, reference genome. In an attempt to create a long-read oyster reference, methodology to successfully isolate ULDNA from eastern oysters was optimized to sequester ultralong oyster DNA from mucopolysaccharides while avoiding excessive shearing. Analysis using a PromethION™ flow cell yielded read lengths (average N50 = 16.5kb, longest read 508kb) that were comparable to what was expected. After quality filtering ($\boldsymbol{Q \geq 9}$), the number of reads $\mathbf{393 \times 10^3}$ were mapped to 91.4% of the reference genome (649Mbp, N50=14kb). However, the sequencing coverage was not sufficiently deep to accomplish de novo assembly. Of all ULDNA reads, 94.6% were successfully mapped to *C. virginica*. Most of the unmapped reads (5.4% of all reads) were classified to bacteria (56%) and human (33%), revealing, respectively, an unexpected oyster pathogen and signaling sewage contamination in the vicinity of oyster harvest.

1 Introduction

Eastern oysters, *Crassostrea virginica*, are both environmentally and economically important in the estuaries where they reside. Eastern oysters provide reef structures that act as habitats for juvenile marine organisms and aid in protection from predators at vulnerable life stages Coen et al. (1999); Harding and Mann (2001). Additionally, oyster reef structures aid in shoreline protection by reducing storm event wave energy Brandon et al. (2016); La Peyre et al. (2015); Scyphers et al. (2011).

Perhaps the most notable function of eastern oysters is as water filterers. At their pre-decline population levels in Chesapeake Bay, eastern oysters were most likely able to filter the water volume from their entire estuarine systems in a few days Newell (1988). Presently, the universal oyster filtration potential is significantly reduced due to their population decline, resulting in pronounced negative effects to the ecosystem Zu Ermgassen et al. (2013). Oysters filter to consume their food source, phytoplankton, and in the process, they remove other microbes, improve water clarity, and transfer biomass to the estuary floor where in the instance of oyster reefs, they promote more favorable nutrient cycling and ultimately decreasing levels of nitrogen pollution Bricker et al. (2020); Dame et al. (1989); Ehrich and Harris (2015); Grizzle et al. (2008, 2017); Higgins et al. (2011, 2013); Hoellein and Zarnoch (2014); Kellogg et al. (2014); Newell et al. (2005). Since they are continuously filtering at high rates, it can be hypothesized that eastern oysters can act as "natural eDNA samplers." Studies have documented the reduction in nitrogen levels due to oyster filtration (Bricker et al., 2018; Kellogg et al., 2014), bringing forward the idea that bivalve species, such as the eastern oyster, could be used as an ecosystem engineer to improve nutrient levels in the estuaries where they are found. However, with a dramatic decline in eastern oyster populations (Bersoza Hernández et al., 2018a; R. Grizzle Ward, 2016; Mackenzie, 2007; Rothschild et al., 1994), there has been a substantial loss of the ecosystem services that they provide (Zu Ermgassen et al., 2013). A better understanding of eastern oyster responses

to their environment is resulting from advances of omics-techniques. Genomic studies have supported extensive population analyses (Hornick Plough, 2022; A. Strickland Brown, 2024; Varney et al., 2009) transcriptomic studies have uncovered expression patterns in oyster as they respond to changing environmental conditions, stress, and disease (Chapman et al., 2011; Eierman Hare, 2014, 2016; S. Wang et al., 2010). Sequencing and annotation of the eastern oyster genome (Gómez-Chiarri et al., 2015) advanced these studies even further, allowing for more in-depth comparisons than were previously possible. Recent advances in Oxford Nanopore Technologies (ONT) DNA sequencing have allowed for sequencing of ultra-long DNA (ULDNA) from various organisms (Frei et al., 2021; Jain et al., 2018; Lu et al., 2024), but methodology has not yet been optimized for mollusks. Many protocols exist to maximize high molecular weight DNA yields for long-read sequencing (Cahyani et al., 2024; Denton et al., 2024; Gong et al., 2019; Jackson Howard, 2023; Nishii et al., 2023; M. Strickland et al., 2023; Todorovic et al., 2023), but to date there are no reports of methods for mollusk-specific ultra-long reads nor are there molluscan DNA genome assemblies from ultra-long mollusk DNA. In our experience, the lack of high-quality high molecular weight DNA from mollusks is in great part due to co-extraction of mucopolysaccharides during DNA extraction from mollusk tissues. Mucopolysaccharide-rich tissue promotes the oyster's ability to successfully filter feed (Beninger St-Jean, 1997), but is detrimental to extraction of ultra-long, high quality DNA (Adema, 2021). Traditional DNA extraction protocols to remove polysaccharides involve extensive washing and mechanical processing that causes shearing of DNA resulting in short fragments, which eliminates the ability for utilizing ONT's ULDNA technology. Therefore, optimization of methodology for ULDNA extraction from oysters is needed, warranting further methodological testing. This study aimed to (1) optimize the extraction and sequencing of ULDNA from eastern oysters and (2) evaluate the potential of eastern oysters as "natural eDNA samplers." Methodology to successfully isolate high-quality

ULDNA from eastern oysters was generated, which required successful sequestering of mucopolysaccharides from oyster DNA without shearing the DNA. Optimization of methods to collect ULDNA from eastern oysters required careful consideration of tissue and modification of washing steps implemented by Brown (1989) (Brown, 1989) to reduce the possibility of DNA release prior to cell lysis. The oyster tissue chosen for preparation of ULDNA was haemocytes, the primary circulating immune cells in molluscs (Anderson et al., 1992; Estrada et al., 2021). Haemocytes are documented to be the most relevant oyster tissue to target for response to parasitic infection (Anderson et al., 1995; Beaven Paynter, 1999), they are known to express a wide variety of genes and proteins, and their transcriptional profiles have been characterized in a related oyster, Magallana gigas (previously Crassostrea gigas (Salvi Mariottini, 2017)) (Estrada et al., 2021; Gueguen et al., 2009; Jin et al., 2021). Furthermore, these nucleated cells can be harvested from haemolymph withdrawn with relative ease from the adductor muscle sinus, without sacrificing the organism (Anderson et al., 1992). Lastly, haemocytes are easily isolated single cells that facilitate washing and are far easier to lyse than other tissues commonly used for DNA such as mantle, gill, and adductor muscle. The washing steps previously documented to reduce mucopolysaccharides in molluscs (Brown, 1989) were optimized for tissue homogenates and involved density gradients. This work aimed to eliminate all homogenization steps and optimize washing. Assembly of the ultra-long reads to both reference genomes was completed to assess success, and unmapped reads were classified to the species level where possible. Results were compared with known species to assess the potential use of oysters as eDNA samplers.

References

Brandon CM, Woodruff JD, Orton PM, et al (2016) Evidence for elevated coastal vulnerability following large-scale historical oyster bed harvesting. Earth Surface Processes and Landforms 41(8):1136–1143. https://doi.org/10.1002/esp.3931, URL

https://onlinelibrary.wiley.com/doi/10.1002/esp.3931

Bricker SB, Grizzle RE, Trowbridge P, et al (2020) Bioextractive Removal of Nitrogen by Oysters in Great Bay Piscataqua River Estuary, New Hampshire, USA. Estuaries and Coasts 43(1):23–38. https://doi.org/10.1007/s12237-019-00661-8, URL https://doi.org/10.1007/s12237-019-00661-8

Coen LD, Luckenbach MW, Breitburg DL (1999) The Role of Oyster Reefs as Essential Fish Habitat: A Review of Current Knowledge and Some New Perspectives 22:438–454

Dame R, Spurrier J, Wolaver T (1989) Carbon, nitrogen and phosphorus processing by an oyster reef. Marine Ecology Progress Series 54:249–256. https://doi.org/10.3354/meps054249, URL http://www.int-res.com/articles/meps/54/m054p249.pdf

Ehrich MK, Harris LA (2015) A review of existing eastern oyster filtration rate models. Ecological Modelling 297:201–212. https://doi.org/10.1016/j.ecolmodel.2014.11.023, URL https://linkinghub.elsevier.com/retrieve/pii/S0304380014005961

Grizzle RE, Greene JK, Coen LD (2008) Seston Removal by Natural and Constructed Intertidal Eastern Oyster (Crassostrea virginica) Reefs: A Comparison with Previous Laboratory Studies, and the Value of in situ Methods. Estuaries and Coasts 31(6):1208–1220. https://doi.org/10.1007/s12237-008-9098-8, URL http://link.springer.com/10.1007/s12237-008-9098-8

Grizzle RE, Ward KM, Peter CR, et al (2017) Growth, morphometrics and nutrient content of farmed eastern oysters, Crassostrea virginica (Gmelin), in New Hampshire, USA. Aquaculture Research 48(4):1525–1537. https://doi.org/10.1111/are.12988, URL https://onlinelibrary.wiley.com/doi/abs/10.1111/are.12988

Harding J, Mann R (2001) Oyster reefs as fish habitat: Opportunistic use of restored reefs by transient fishes. Journal of Shellfish Research 20(3):951–959. URL https://scholarworks.wm.edu/vimsarticles/1438

Higgins CB, Stephenson K, Brown BL (2011) Nutrient Bioassimilation Capacity of Aquacultured Oysters: Quantification of an Ecosystem Service. Journal of Environmental Quality 40(1):271–277. https://doi.org/10.2134/jeq2010.0203, URL https://acsess.onlinelibrary.wiley.com/doi/10.2134/jeq2010.0203

Higgins CB, Tobias C, Piehler M, et al (2013) Effect of aquacultured oyster biodeposition on sediment N<sub>2 production in Chesapeake Bay. Marine Ecology Progress Series 473:7–27. https://doi.org/10.3354/meps10062, URL http://www.int-res.com/abstracts/meps/v473/p7-27/

Hoellein TJ, Zarnoch CB (2014) Effect of eastern oysters (*Crassostrea virginica*) on sediment carbon and nitrogen dynamics in an urban estuary. Ecological Applications 24(2):271–286. https://doi.org/10.1890/12-1798.1, URL https://esajournals.onlinelibrary.wiley.com/doi/10.1890/12-1798.1

Kellogg ML, Smyth AR, Luckenbach MW, et al (2014) Use of oysters to mitigate eutrophication in coastal waters. Estuarine, Coastal and Shelf Science 151:156–168. https://doi.org/10.1016/j.ecss.2014.09.025, URL https://linkinghub.elsevier.com/retrieve/pii/S0272771414002789

La Peyre MK, Serra K, Joyner TA, et al (2015) Assessing shoreline exposure and oyster habitat suitability maximizes potential success for sustainable shoreline protection using restored oyster reefs. PeerJ 3:e1317. https://doi.org/10.7717/peerj.1317, URL https://peerj.com/articles/1317

Newell RIE (1988) Ecological Changes in Chesapeake Bay: Are They The Result of Overharvesting the American Oyster, Crassostrea virgincia? In: Proceedings of the Chesapeake Bay Research Consortium Conference, Baltimore, MD

Newell RIE, Fisher TR, Holyoke RR, et al (2005) Influence of Eastern Oysters on Nitrogen and Phosphorus Regeneration in Chesapeake Bay, USA. In: Dame RF, Olenin S (eds) The Comparative Roles of Suspension-Feeders in Ecosystems, vol 47. Springer-Verlag, Berlin/Heidelberg, p 93–120, https://doi.org/10.1007/1-4020-3030-4_6, URL http://link.springer.com/10.1007/1-4020-3030-4_6, series Title: NATO Science Series IV: Earth and Environmental Series

Scyphers SB, Powers SP, Heck KL, et al (2011) Oyster Reefs as Natural Breakwaters Mitigate Shoreline Loss and Facilitate Fisheries. PLoS ONE 6(8):e22396. https://doi.org/10.1371/journal.pone.0022396, URL https://dx.plos.org/10.1371/journal.pone.0022396

Zu Ermgassen PSE, Spalding MD, Grizzle RE, et al (2013) Quantifying the Loss of a Marine Ecosystem Service: Filtration by the Eastern Oyster in US Estuaries. Estuaries and Coasts 36(1):36–43. https://doi.org/10.1007/s12237-012-9559-y, URL http://link.springer.com/10.1007/s12237-012-9559-y