



UNIVERSITY OF SCIENCE AND TECHNOLOGY OF HANOI

Mean-field study of the equation of states of nuclear matter and tidal deformation of neutron star

Bachelor Thesis

Author:

Nguyen Hoang Dang KHOA

Supervisors:

Prof. Dao Tien KHOA

Dr. Ngo Hai TAN

June 10, 2021

Contents

List of Abbreviations	2
List of Tables	3
List of Figures	4
1 Introduction	5
2 Hartree-Fock Formalism of Nuclear Matter	8
2.1 Nucleon-Nucleon Interaction	8
2.2 Equation of States of Nuclear Matter	11
2.3 β -Stable Nuclear Matter	12
3 Neutron Star Properties	14
3.1 Tolman-Oppenheimer-Volkoff Equation	14
3.2 Gravito-electric and Gravito-magnetic Tidal Deformation	14
4 Results and Discussions	17
5 Conclusions	24
Bibliography	25

List of Abbreviations

BH	Black hole
BHF	Brueckner-Hartree-Fock
EoS	Equation of States
GE	Gravito-electric
GM	Gravito-magnetic
GR	General Relativity
GRB	Gamma-ray burst
GW	Gravitational wave
HF	Hartree-Fock
NM	Nuclear Matter
NN	Nucleon-Nucleon
NS	Neutron Star
TOV	Tolmann-Oppenheimer-Volkoff

List of Tables

2.1	CDM3Y n interaction's parameters; the 00 and 01 terms are inherited from (Tan et al., 2021), while the 10 and 11 parameters are added by fitting with BHF result.	10
2.2	Yukawa strengths of the M3Y-Paris interaction (Tan et al., 2020 ; Anantaraman et al., 1983).	11

List of Figures

1.1	Neutron star's overall structure. The baryon density decreases (from white to dark gray) as we move outward from the NS center.	6
1.2	Illustration of binary NS merger. (a) The two companion NSs orbit about each others, while gradually losing energy through weak GW and come closer with time. (b) As the two NS get closer, they accelerate and emit stronger GW until (c) colliding, which results in a <i>kilonova</i> , characterized by a short <i>gamma ray burst</i> (GRB). The product of the merger has yet been decided to be a black hole (BH) or another NS.	7
2.1	Energy per baryon E/A of symmetric NM by the 5 CDM3Yn models compared to BHF result (Vidana and Bombaci, 2002). The diamond and square represent the BHF result for $\Delta_n = -\Delta_p = \pm 1$ and $\Delta_n = \Delta_p = \pm 1$ respectively with Δ_τ being the baryon spin polarity.	10
4.1	Proton fraction x_p of β -stable NM at different baryon density and spin polarity for CDM3Yn interactions.	17
4.2	Total mass-energy density of β -stable NM at varying spin polarity with different interaction models.	18
4.3	Total pressure of β -stable NM at several values of Δ with different CDM3Yn models. . .	18
4.4	The relation between gravitational mass M and the radius R of the NS according to the corresponding model and polarity.	19
4.5	GE tidal Love number at 2 nd order k_2 as function of NS mass computed with CDM3Yn models at different spin polarities.	19
4.6	Same as Figure 4.5 for k_3	20
4.7	Same as Figure 4.5 for k_4	20
4.8	GM tidal Love number at 2 nd order j_2 as function of NS mass computed with CDM3Yn models of <i>strictly static fluid</i> at different polarities.	21
4.9	Same as Figure 4.8 for j_3	21
4.10	Same as Figure 4.8 for j_4	22
4.11	GM tidal Love number at 2 nd order j_2 as function of NS mass computed with CDM3Yn models of <i>irrotational fluid</i> at different polarities.	22
4.12	Same as Figure 4.11 for j_3	23
4.13	Same as Figure 4.11 for j_4	23

Chapter 1

Introduction

Neutron stars (NS) are star-like astronomical objects with mass M on the order of solar mass (M_\odot), a radius of $\sim 10 - 12 \text{ km}$ and an average density n several times greater than that of nucleon ($\rho_0 \approx 0.16 \text{ fm}^{-3}$). They are arguably the densest accessible objects, excluding black holes which we know nothing about inside the event horizon, in the universe (Baym and Pethick, 1975). Due to extremely high density, the matter on NS mainly consists of neutrons that are closely packed together with a small percentage of other particles (p, e^-, \dots), similar to a atomic nucleus on macroscopic scale. For this reason, they are also the ideal objects for testing physical theories of dense matter and provide connections between different field of physics, i.e. nuclear physics, elementary particle physics and astrophysics (Lattimer and Prakash, 2004).

During the NS's formation process, protons (p) and electrons (e^-) combined together to form neutrons, i.e.

$$p + e^- \longrightarrow n + \nu_e \quad (1.1)$$

and the star only holds itself against gravity by its own degeneracy pressure and strong force repulsion, which explains why the matter on NS is neutron-dominant and hence the name “neutron stars”. After the NS is formed, energy quickly dissipates through neutrino emission, resulting in a relatively cold NS. In this study, we will only concern with the NS after a considerable time from its formation, when the temperature is considered to be $T = 0 \text{ K}$.

On a NS, the matter exists as an inhomogenous, low-density *crust* and gradually becomes a more uniform *core* the closer to the NS center as in Figure 1.1. In order to study about the properties of NS matter, the problem have to be approached from the nuclear physics point of view, where we study about *nuclear matter* (NM). For a nuclear system as massive as a NS, we consider one with infinite number of nucleons that are in β -stable state with a small portion of leptons, in which the properties of matter are described using an *equation of states* (EoS), i.e. the relation between different state variables (pressure P , mass-energy density ε , ...) of the system. Ideally, the EoS can be derived from the interactions of quarks under strong force in the framework of quantum chromodynamics. However, due to this having yet to be possible at the moment, the EoS of NM is instead interpreted from a nonrelativistic mean-field study approach with several updated versions of the realistic density-dependent CDM3Yn interaction models (Khoa et al., 1995; Khoa and Cuong, 2007) using Hartree-Fock (HF) formalism, which will be implemented further in Chapter 2.

Following the gravitational wave (GW) signals GW170817 (Abbott et al., 2017) and GW190425




Figure 1.1: Neutron star’s overall structure. The baryon density decreases (from white to dark gray) as we move outward from the NS center.

(Abbott et al., 2020) from two binary NS mergers observed by LIGO and Virgo laser interferometer in 17th August 2017 and 25th April 2019 respectively, the tidal deformation of the NS can be further constrained, as well as the mass M and radius R of the NS (Abbott et al., 2018). The NS merger event is illustrated as in Figure 1.2.

Apparently, the EoS of high-density NM plays the most important role in deciding the macroscopic properties of NS. In particular, given the EoS of the crust from the compressible liquid drop model and by using the EoS of the uniform NS core from the result of the HF calculation of cold β -stable NM, the gravitational mass and radius of the star can be decided by the framework of General Relativity (GR) (Tan et al., 2020, 2021), i.e. the Tolman-Oppenheimer-Volkoff (TOV) equation, which will in turn be compared to the observational astrophysical constraints to deduce the most suitable EoS of the constituent NS in this system.

In addition, due to the enormous mass, each NS possesses powerful gravitational field and therefore, they tend to “stretch out” their companion under the tidal effect as in Figure 1.2, while orbiting spirally toward each others and dissipating energy under the form of GW. Particularly, the shape and mass-energy distribution of the NS are tidally deformed from its supposedly spherical shape, resulting in nonzero multipole moments (Hinderer, 2008; Hinderer et al., 2010; Damour and Nagar, 2009). The NS’s reaction, i.e. how strongly it deforms when being under a tidal field, is expressed in terms of the *tidal Love numbers* k_l of several orders l , where in this study, we will evaluate the Love number of NS up to the 4th order, i.e. $l = 2, 3, 4$ (Perot and Chamel, 2021). Apparently, the tidal Love number depends heavily on the EoS of matter and this dependence will be further emphasized in Chapter 3. For NS, the central density can be up to $6\rho_0$ and possesses a Love number of order ~ 0.1 , while our Earth has that of 0.3. In a recent study, the Love number was calculated for spinning black holes, which showed that even with nearly infinite density, they still possess a small Love number of 0.002 (Le Tiec and Casals,




Figure 1.2: Illustration of binary NS merger. (a) The two companion NSs orbit about each others, while gradually losing energy through weak GW and come closer with time. (b) As the two NS get closer, they accelerate and emit stronger GW until (c) colliding, which results in a *kilonova*, characterized by a short *gamma ray burst* (GRB). The product of the merger has yet been decided to be a black hole (BH) or another NS.

2021).

Furthermore, under small perturbation of spacetime, the tidal field can also be separated into two components: the *gravito-electric* (GE) and *gravito-magnetic* (GM) terms (Damour and Nagar, 2009) that are analogous to that in eletromagnetic field. As a result, the deformation of the NS, i.e. Love numbers, in the perturbed tidal field can also be categorized into the corresponding GE (k_l) and GM (j_l) components (Perot and Chamel, 2021), whose result will be presented in more details in Chapter 3. To sum up, this study is dedicated to:

- Include the spin polarization effect to the existing CDM3Y n models (Tan et al., 2021),
- Assess the dependence of NS's gravitational mass and radius to different EoS,
- Investigate the sensitivity of GE and GM tidal deformability and Love numbers to NM properties,
- Compare the NS's calculated properties to the astrophysical constraints obtained experimentally.

Chapter 2

Hartree-Fock Formalism of Nuclear Matter

2.1 Nucleon-Nucleon Interaction

Due to the lack of an exact theory to describe the nucleon-nucleon (NN) interaction, a model needs to be imposed and fit with experimental measurement or theoretical calculation results. Plus, for a system as massive as a NS, deducing the EoS using the *ab initio* method, i.e. solving the Schrödinger equation over all particles, is simply impossible, therefore an *effective interaction* must be used (Greiner and Maruhn, 1996). In this section, we only limit ourselves to two-body interaction, thus, the NN potential can be expressed in the form of

$$v = v(\mathbf{r}, \mathbf{r}', \mathbf{p}, \mathbf{p}', \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') \quad (2.1)$$

where the primed and unprimed variables indicate the properties of 2 nucleons respectively, in which \mathbf{r} is the particle's position, \mathbf{p} is its momentum, $\boldsymbol{\sigma}$ is its intrinsic spin and $\boldsymbol{\tau}$ is its isospin.

The functional form of v in (2.1) cannot freely take any form but is constrained by many invariance requirements (Greiner and Maruhn, 1996)

- **Translational invariance:** The NN potential should only depend on the *relative position* of the two particles but not their explicit positions, thus we can reduce (2.1) to

$$v = v(\mathbf{r} - \mathbf{r}', \mathbf{p}, \mathbf{p}', \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') = v(\mathbf{r}, \mathbf{p}, \mathbf{p}', \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') \quad (2.2)$$

with the last expression, we redefine \mathbf{r} as the relative position vector.

- **Galilei invariance:** The potential should also be invariant under transformation between inertial frames of reference, which requires that only the relative momentum $\mathbf{p} - \mathbf{p}'$ is depended, i.e.

$$v = v(\mathbf{r}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') \quad (2.3)$$

where here we denote \mathbf{p} as the relative momentum.

- **Rotational invariance:** The potential should be constructed such that the total angular momentum is zero.

- **Isospin invariance:** The NN interaction potential needs to be invariant under rotation in isospin space, in other words, it can only depend on the isospin-independent terms and the terms with $\boldsymbol{\tau} \cdot \boldsymbol{\tau}'$. Therefore, we can split the potential into

$$v = v_0(\mathbf{r}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\sigma}') + v_1(\mathbf{r}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\sigma}') \hat{\boldsymbol{\tau}} \cdot \hat{\boldsymbol{\tau}}' \quad (2.4)$$

- **Parity invariance:** The NN interaction potential is also expected to be invariant under the action of parity operator, i.e. changing the sign of spatial coordinates

$$v(\mathbf{r}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') = v(-\mathbf{r}, -\mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') \quad (2.5)$$

- **Time reversal invariance:** Finally, the interaction should stay the same after switching the time arrow direction

$$v(\mathbf{r}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') = v(\mathbf{r}, -\mathbf{p}, -\boldsymbol{\sigma}, -\boldsymbol{\sigma}', \boldsymbol{\tau}, \boldsymbol{\tau}') \quad (2.6)$$

Having the above considerations, developing further the M3Y-Paris interaction, which was used by the HF study of NM (Loan et al., 2011; Tan et al., 2016, 2020, 2021) and the folding model study of NN scattering (Khoa et al., 1997; Khoa and Satchler, 2000),

$$v = v_{00}(r) + v_{10}(r) \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + v_{01}(r) \boldsymbol{\tau} \cdot \boldsymbol{\tau}' + v_{11}(r) (\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}') (\boldsymbol{\tau} \cdot \boldsymbol{\tau}') \quad (2.7)$$

by adding a density-dependent form factor to each term gives the CDM3Yn interaction model

$$\begin{aligned} v(\rho, r) = & F_{00}(\rho) v_{00}(r) + F_{10}(\rho) v_{10}(r) \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' \\ & + F_{01}(\rho) v_{01}(r) \boldsymbol{\tau} \cdot \boldsymbol{\tau}' + F_{11}(\rho) v_{11}(r) (\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}') (\boldsymbol{\tau} \cdot \boldsymbol{\tau}') \end{aligned} \quad (2.8)$$

where each radial term is the superposition of 3 Yukawa potentials

$$v_{\sigma\tau}(r) = \sum_{k=1}^3 Y_{\sigma\tau}(k) \frac{\exp(-\mu_k r)}{\mu_k r} \quad (2.9)$$

and the form factor $F_{\sigma\tau}(\rho)$ shared the functional form (Khoa et al., 1997; Tan et al., 2020, 2021; Thân, 2010)

$$F_{\sigma\tau}(\rho) = C_{\sigma\tau} [1 + \alpha_{\sigma\tau} \exp(-\beta_{\sigma\tau} \rho) + \gamma_{\sigma\tau} \rho] \quad (2.10)$$

with parameters given in Table 2.1. The parameters of F_{00} were adjusted to give the corresponding incompressibility K of symmetric NM at saturation density ρ_0 and the binding energy $E_0 \approx 15.8 \text{ MeV}$, while the 10 term is modified from (Thân, 2010) to reproduce $E_{\text{sym}}(\rho_0) \approx 30 \text{ MeV}$, $L \approx 50 \text{ MeV}$ and to be in agreement with the ab-initio results (Akmal et al., 1998; Gandolfi et al., 2010) at higher density (Tan et al., 2021). On the other hand, the spin-dependent terms, 10 and 11, are hereby included in the 5 models by fine tuning the parameters to yield the same result as the Brueckner-Hartree-Fock (BHF) study of spin polarized NM (Vidana and Bombaci, 2002) as in Figure 2.1.




Figure 2.1: Energy per baryon E/A of symmetric NM by the 5 CDM3Y n models compared to BHF result (Vidana and Bombaci, 2002). The diamond and square represent the BHF result for $\Delta_n = -\Delta_p = \pm 1$ and $\Delta_n = \Delta_p = \pm 1$ respectively with Δ_τ being the baryon spin polarity.

Table 2.1: CDM3Y n interaction's parameters; the 00 and 01 terms are inherited from (Tan et al., 2021), while the 10 and 11 parameters are added by fitting with BHF result.

Interaction	$\sigma\tau$	$C_{\sigma\tau}$	$\alpha_{\sigma\tau}$	$\beta_{\sigma\tau}$ (fm^3)	$\gamma_{\sigma\tau}$ (fm^3)	K (MeV)
CDM3Y3	00	0.2985	3.4528	2.6388	-1.5	217
	01	0.2343	5.3336	6.4738	4.3172	
	10	0.3890	3.5635	-2.6717	20.3624	
	11	0.8802	4.0433	12.3262	0.3662	
CDM3Y4	00	0.3052	3.2998	2.3180	-2.0	228
	01	0.2129	6.3581	7.0584	5.6091	
	10	0.2593	6.0016	-2.3377	18.8725	
	11	0.8329	3.5941	9.2012	0.2690	
CDM3Y5	00	0.2728	3.7367	1.8294	-3.0	241
	01	0.2204	6.6146	7.9910	6.0040	
	10	0.4106	5.6265	-1.6698	-1.9866	
	11	0.6815	2.5833	5.1700	0.2578	
CDM3Y6	00	0.2658	3.8033	1.4099	-4.0	252
	01	0.2313	6.6865	8.6775	6.0182	
	10	0.5186	9.9402	1.6698	2.9799	
	11	0.6058	3.1947	4.4512	0.0822	
CDM3Y8	00	0.2658	3.8033	1.4099	-4.3	257
	01	0.2643	6.3836	9.8950	5.4249	
	10	0.5997	9.1900	0.7514	-4.7181	
	11	0.3786	3.9435	2.7012	0.3512	

2.2 Equation of States of Nuclear Matter

In HF formalism, the total HF energy of the system can be expressed as

$$E_{HF} = \sum_{\sigma\tau} \sum_{\mathbf{k}}^{\frac{k_F^{\sigma\tau}}{2}} \frac{\hbar^2 k^2}{2m_\tau} + \frac{1}{2} \sum_{\mathbf{k}\sigma\tau} \sum_{\mathbf{k}'\sigma'\tau'} \left[\langle \mathbf{k}\sigma\tau, \mathbf{k}'\sigma'\tau' | v^D | \mathbf{k}\sigma\tau, \mathbf{k}'\sigma'\tau' \rangle + \langle \mathbf{k}\sigma\tau, \mathbf{k}'\sigma'\tau' | v^{EX} | \mathbf{k}'\sigma\tau, \mathbf{k}\sigma'\tau' \rangle \right] \quad (2.11)$$

where the single-particle wave function is plane wave

$$|\mathbf{k}\sigma\tau\rangle = \frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{\sqrt{\Omega}} \chi_\sigma \chi_\tau \quad (2.12)$$

Ω being the spatial volume of the system, $k_F^{\sigma\tau} = (6\pi^2 \rho_{\sigma\tau})^{1/3}$ is the Fermi momentum corresponding to spin σ and isospin τ , $v^{D(EX)}$ is the direct (exchange) part of the interaction determined from the singlet and triplet-even (odd) of the central NN force. Adopting the same functional form of (2.8), the direct and exchange interaction is written as

$$v^{D(EX)}(\rho_b, r) = F_{00}(\rho_b) v_{00}^{D(EX)}(r) + F_{10}(\rho_b) v_{10}^{D(EX)}(r) \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + F_{01}(\rho_b) v_{01}^{D(EX)}(r) \boldsymbol{\tau} \cdot \boldsymbol{\tau}' + F_{11}(\rho_b) v_{11}^{D(EX)}(r) (\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}') (\boldsymbol{\tau} \cdot \boldsymbol{\tau}') \quad (2.13)$$

and

$$v_{\sigma\tau}^{D(EX)}(r) = \sum_{k=1}^3 Y_{\sigma\tau}^{D(EX)}(k) \frac{\exp(-\mu_k r)}{\mu_k r} \quad (2.14)$$

with the Yukawa strengths given in Table 2.2 and the density-dependent form factor parameters are in Table 2.1. Note that in (2.13), ρ_b denotes the *baryon density*, this will be used in order to distinguish with the lepton density in the later section.

Table 2.2: Yukawa strengths of the M3Y-Paris interaction ([Tan et al., 2020](#); [Anantaraman et al., 1983](#)).

k	μ_k (fm^{-1})	Y_{00}^D (MeV)	Y_{10}^D (MeV)	Y_{01}^D (MeV)	Y_{11}^D (MeV)	Y_{00}^{EX} (MeV)	Y_{10}^{EX} (MeV)	Y_{01}^{EX} (MeV)	Y_{11}^{EX} (MeV)
1	4.0	11061.625	938.875	313.625	-969.125	-1524.25	-3492.75	-4118.0	-2210.0
2	2.5	-2537.5	-36.0	223.5	450.0	-518.75	795.25	1054.75	568.75
3	0.7072	0.0	0.0	0.0	3.4877	-7.8474	2.6157	2.6157	-0.8719

Multiply (2.11) with Ω^{-1} , the energy density of the NM is separated into the kinetic term ε_{kin} and the potential terms $\varepsilon_{\sigma\tau}$, i.e.

$$\varepsilon_{HF} = \frac{E_{HF}}{\Omega} = \varepsilon_{kin} + F_{00}(\rho_b) \varepsilon_{00} + F_{01}(\rho_b) \varepsilon_{01} + F_{10}(\rho_b) \varepsilon_{10} + F_{11}(\rho_b) \varepsilon_{11} \quad (2.15)$$

The final expressions of each terms of the energy density are

$$\varepsilon_{kin} = \frac{3}{10} \sum_{\sigma\tau} \frac{\hbar^2 (k_F^{\sigma\tau})^2}{m_\tau} \rho_{\sigma\tau} \quad (2.16)$$

$$\varepsilon_{00} = \frac{1}{2} \left[\rho_b^2 J_{00}^D + \int A_{00}^2 v_{00}^{EX}(r) d^3r \right] \quad (2.17)$$

$$\varepsilon_{10} = \frac{1}{2} \left[\rho_b^2 J_{10}^D \left(\Delta_n \frac{1+\delta}{2} + \Delta_p \frac{1-\delta}{2} \right)^2 + \int A_{10}^2 v_{10}^{EX}(r) d^3r \right] \quad (2.18)$$

$$\varepsilon_{01} = \frac{1}{2} \left[\rho_b^2 J_{01}^D \delta^2 + \int A_{01}^2 v_{01}^{EX}(r) d^3r \right] \quad (2.19)$$

$$\varepsilon_{11} = \frac{1}{2} \left[\rho_b^2 J_{11}^D \left(\Delta_n \frac{1+\delta}{2} - \Delta_p \frac{1-\delta}{2} \right)^2 + \int A_{11}^2 v_{11}^{EX}(r) d^3r \right] \quad (2.20)$$

where $\Delta_\tau = (\rho_{\uparrow\tau} - \rho_{\downarrow\tau})/\rho_\tau$ is the polarity of nucleon, $\delta = (\rho_n - \rho_p)/\rho_b$ is the asymmetry of NM, $J_{\sigma\tau}^D = \int v_{\sigma\tau}^D(r) d^3r$ is the volume integral of the direct interaction and

$$\begin{aligned} A_{00} &= \rho_{\uparrow n} \hat{j}_1(k_F^{\uparrow n} r) + \rho_{\downarrow n} \hat{j}_1(k_F^{\downarrow n} r) + \rho_{\uparrow p} \hat{j}_1(k_F^{\uparrow p} r) + \rho_{\downarrow p} \hat{j}_1(k_F^{\downarrow p} r) \\ A_{10} &= \rho_{\uparrow n} \hat{j}_1(k_F^{\uparrow n} r) - \rho_{\downarrow n} \hat{j}_1(k_F^{\downarrow n} r) + \rho_{\uparrow p} \hat{j}_1(k_F^{\uparrow p} r) - \rho_{\downarrow p} \hat{j}_1(k_F^{\downarrow p} r) \\ A_{01} &= \rho_{\uparrow n} \hat{j}_1(k_F^{\uparrow n} r) + \rho_{\downarrow n} \hat{j}_1(k_F^{\downarrow n} r) - \rho_{\uparrow p} \hat{j}_1(k_F^{\uparrow p} r) - \rho_{\downarrow p} \hat{j}_1(k_F^{\downarrow p} r) \\ A_{11} &= \rho_{\uparrow n} \hat{j}_1(k_F^{\uparrow n} r) - \rho_{\downarrow n} \hat{j}_1(k_F^{\downarrow n} r) - \rho_{\uparrow p} \hat{j}_1(k_F^{\uparrow p} r) + \rho_{\downarrow p} \hat{j}_1(k_F^{\downarrow p} r) \end{aligned} \quad (2.21)$$

with $\hat{j}_1(x) = 3j_1(x)/x$ and $j_1(x)$ being the 1st order spherical Bessel function.

2.3 β -Stable Nuclear Matter

After the HF calculation, we were able to obtain a numerical HF energy density $\varepsilon(\rho_n, \rho_p, \Delta_n, \Delta_p)$. However, it is in fact impossible for a NS to exist while consisting of purely nucleon. In order to compensate for this issue, leptons (e^- and μ^-) have to be introduced to the matter constituents and the $npe\mu$ matter has to satisfy the β -stable condition (Glendenning, 2012), i.e.

- Charge balance

$$\rho_p = \rho_e + \rho_\mu \quad (2.22)$$

- Chemical potential balance

$$\mu_n - \mu_p = \mu_e = \mu_\mu \quad (2.23)$$

where $\mu_i = \frac{\partial \varepsilon}{\partial \rho_i}$ ($i = n, p, e, \mu$) is the chemical potential of the i particle.

The total energy density of the $npe\mu$ matter is thus

$$\varepsilon = \varepsilon_{HF} + \rho_n m_n c^2 + \rho_p m_p c^2 + \varepsilon_e + \varepsilon_\mu \quad (2.24)$$

which leads to the nucleon chemical potential of the form

$$\mu_\tau(\rho_n, \rho_p, \Delta_n, \Delta_p) = \frac{\partial \varepsilon}{\partial \rho_\tau} = \frac{\partial \varepsilon_{HF}}{\partial \rho_\tau} + m_\tau c^2 \quad (2.25)$$

Let $\hat{\mu} = \mu_n - \mu_p$ be the leptons' chemical potential, (2.22) is equivalent to¹

$$3\pi^2(\hbar c)^3 \rho_p - \hat{\mu}^3 - [\hat{\mu}^2 - (m_\mu c^2)^2]^{3/2} \theta(\hat{\mu} - m_\mu c^2) = 0 \quad (2.26)$$

from which the proton fraction $x_p = \rho_p/\rho_b$ can be obtained as shown in Figure 4.1. Furthermore, under strong magnetic field like that of a magnetar, we can approximate $\Delta_n \approx -\Delta_p \approx \Delta$ and reduce the EoS to depend on just the baryon polarity Δ alone, and the more baryon polarized, the stronger the magnetic field of the NS.

For a fixed value of Δ , we are able to obtain a density function of the form $\rho_n(\rho_b, \Delta)$ and $\rho_p(\rho_b, \Delta)$, which in turn gives the lepton chemical potential $\hat{\mu}(\rho_b, \Delta) = \hat{\mu}(\rho_n, \rho_p)$. On the other hand, the leptons' densities are then [Loan et al. \(2011\)](#)

$$\rho_e(\rho_b, \Delta) = \frac{\hat{\mu}^3(\rho_b, \Delta)}{3\pi^2(\hbar c)^3} \quad \text{and} \quad \rho_\mu(\rho_b, \Delta) = \frac{[\hat{\mu}^2(\rho_b, \Delta) - (m_\mu c^2)^2]^{3/2}}{3\pi^2(\hbar c)^3} \theta(\hat{\mu}(\rho_b, \Delta) - m_\mu c^2) \quad (2.27)$$

Consider the e^- and μ^- to be systems of relativistic Fermi gas, then their respective energy densities and pressure contributions are ($l = e, \mu$) ([Moustakidis and Panos, 2009](#))

$$\varepsilon_l(\rho_b, \Delta) = \frac{2}{(2\pi)^3} \int_0^{[3\pi^2 \rho_l(\rho_b, \Delta)]^{1/3}} \sqrt{\hbar^2 c^2 k^2 + m_l^2 c^4} d^3 \mathbf{k} \quad (2.28)$$

and

$$P_l(\rho_b, \Delta) = \frac{1}{3} \frac{2}{(2\pi)^3} \int_0^{[3\pi^2 \rho_l(\rho_b, \Delta)]^{1/3}} \frac{\hbar^2 c^2 k^2}{\sqrt{\hbar^2 c^2 k^2 + m_l^2 c^4}} d^3 \mathbf{k} \quad (2.29)$$

Plus, from the HF formalism with NM, the baryon pressure is given by

$$P_b = \rho_b^2 \frac{\partial(\varepsilon_{HF}/\rho_b)}{\partial \rho_b} \quad (2.30)$$

Finally, we obtain the total energy density dependence on baryon density as

$$\varepsilon(\rho_b, \Delta) = \varepsilon_{HF}(\rho_b, \Delta) + \rho_n(\rho_b, \Delta) m_n c^2 + \rho_p(\rho_b, \Delta) m_p c^2 + \varepsilon_e(\rho_b, \Delta) + \varepsilon_\mu(\rho_b, \Delta) \quad (2.31)$$

and the total pressure of NS matter

$$P(\rho_b, \Delta) = P_b(\rho_b, \Delta) + P_e(\rho_b, \Delta) + P_\mu(\rho_b, \Delta) \quad (2.32)$$

and this completes the EoS of cold β -stable NS matter (Figure 4.2 and 4.3).

¹ $\theta(x)$ is the Heaviside function, i.e. it returns 1 for $x \geq 0$ and 0 otherwise.

Chapter 3

Neutron Star Properties

3.1 Tolman-Oppenheimer-Volkoff Equation

Suppose the NS to be static and spherically symmetric, the metric elements are then ([Glendenning, 2012](#))

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = e^{2\nu(r)}c^2dt^2 - e^{2\lambda(r)}dr^2 - r^2d\theta^2 - r^2\sin^2\theta d\phi^2 \quad (3.1)$$

Consider the NS matter to be perfect fluid, the energy-momentum tensor of its has the form of

$$T^{\mu\nu} = -Pg^{\mu\nu} + (P + \varepsilon)u^\mu u^\nu \quad (3.2)$$

where $u^\mu = dx^\mu/d\tau$ is the local fluid 4-velocity. Solving the Einstein's field equation

$$G^{\mu\nu} = -\frac{8\pi G}{c^4}T^{\mu\nu} \quad (3.3)$$

gives the Tolman-Oppenheimer-Volkoff (TOV) equation ([Glendenning, 2012](#))

$$\frac{dP}{dr} = -\frac{G\varepsilon(P)m}{c^2r^2} \left(1 + \frac{P}{\varepsilon(P)}\right) \left(1 + \frac{4\pi Pr^3}{mc^2}\right) \left(1 - \frac{2Gm}{c^2r}\right)^{-1} \quad (3.4)$$

$$\frac{dm}{dr} = \frac{4\pi r^2\varepsilon(P)}{c^2} \quad (3.5)$$

where $\varepsilon(P)$ is the EoS obtained from the CDM3Yn interaction calculated previously. Additional boundary conditions are

$$P(0) = P_c; \quad P(R) = 0; \quad m(0) = 0; \quad m(R) = M$$

and by varying the center pressure P_c , a relation of the gravitational mass M and radius R of the NS can be obtained.

3.2 Gravito-electric and Gravito-magnetic Tidal Deformation

In close orbit with another compact companion in a binary system, the NS is tidally deformed by strong gravitational interaction. Analogous to the classical theory of electromagnetism, the tidal field

experienced by it can be decomposed into 2 types: the *gravito-electric* and *gravito-magnetic* components with respective *relativistic tidal moment* (Damour and Nagar, 2009)

$$\mathcal{E}_L = \partial_{L-1} E_{a_l} \quad \text{and} \quad \mathcal{M}_L = c^2 \partial_{L-1} B_{a_l} \quad (3.6)$$

where E_{a_l} and B_{a_l} are the a_l component of the externally generated local GE and GM field, L represents the multi-index (a_1, a_2, \dots, a_l) and l being the order of the moment. As a result, the deformation of NS is parameterized by the GE and GM *tidal deformabilities* λ_l and σ_l , i.e. in leading order (Damour and Nagar, 2009)

$$\mathcal{Q}_L = \lambda_l \mathcal{E}_L, \quad (3.7)$$

$$\mathcal{S}_L = \sigma_l \mathcal{M}_L \quad (3.8)$$

with \mathcal{Q}_L being the induced mass multipole moment, i.e. the deviation of the mass distribution from spherically symmetry at order l , while \mathcal{S}_L is the current multipole moment in adiabatic approximation (Damour and Nagar, 2009; Perot and Chamel, 2021). From the deformabilities, the dimensionless GE and GM *tidal Love numbers* are defined as (Perot and Chamel, 2021)

$$k_l = \frac{1}{2}(2l-1)!! \frac{G\lambda_l}{R^{2l+1}} \quad \text{and} \quad j_l = 4(2l-1)!! \frac{G\sigma_l}{R^{2l+1}} \quad (3.9)$$

These parameters are related to the GE and GM *tidal deformability parameters* as

$$\Lambda_l = \frac{2}{(2l-1)!!} k_l \left(\frac{c^2 R}{GM} \right)^{2l+1} \quad (3.10)$$

$$\Sigma_l = \frac{1}{4(2l-1)!!} j_l \left(\frac{c^2 R}{GM} \right)^{2l+1} \quad (3.11)$$

which can be potentially extracted from the signal of GW. In order to properly calculate these parameters, let $H_l(r)$ and $\tilde{H}_l(r)$ characterize small perturbations of the static metric. These functions have to satisfy (Perot and Chamel, 2021; Damour and Nagar, 2009)

$$\begin{aligned} & H_l''(r) + H_l'(r) \left[1 - \frac{2Gm(r)}{c^2 r} \right]^{-1} \left\{ \frac{2}{r} - \frac{2Gm(r)}{c^2 r^2} - \frac{4\pi G}{c^4} r [\varepsilon(r) - P(r)] \right\} \\ & + H_l(r) \left[1 - \frac{2Gm(r)}{c^2 r} \right]^{-1} \left\{ \frac{4\pi G}{c^4} \left[5\varepsilon(r) + 9P(r) + c^2 \frac{d\varepsilon}{dP} [\varepsilon(r) + P(r)] \right] \right. \\ & \left. - \frac{l(l+1)}{r^2} - 4 \left[1 - \frac{2Gm(r)}{c^2 r} \right]^{-1} \left[\frac{Gm(r)}{c^2 r^2} + \frac{4\pi G}{c^4} r P(r) \right]^2 \right\} = 0 \end{aligned} \quad (3.12)$$

for GE perturbations and

$$\begin{aligned} & \tilde{H}_l''(r) - \tilde{H}_l'(r) \left[1 - \frac{2Gm(r)}{c^2 r} \right]^{-1} \frac{4\pi G}{c^4} r [P(r) + \varepsilon(r)] \\ & - \tilde{H}_l(r) \left[1 - \frac{2Gm(r)}{c^2 r} \right]^{-1} \left\{ \frac{l(l+1)}{r^2} - \frac{4Gm(r)}{c^2 r^3} + \theta \frac{8\pi G}{c^4} [P(r) + \varepsilon(r)] \right\} = 0 \end{aligned} \quad (3.13)$$

for GM perturbations; the value of $\theta = 1$ is for static fluid while irrotational fluid adopts the value $\theta = -1$. These two equations are integrated along with the TOV equation (3.4). In addition, we have the compactness parameters $C = GM/(Rc^2)$ and define

$$y_l = \frac{RH'_l(R)}{H_l(R)} \quad \text{and} \quad \tilde{y}_l = \frac{R\tilde{H}'_l(R)}{\tilde{H}_l(R)}. \quad (3.14)$$

The explicit expressions of the first few orders of the GE and GM Love numbers are ([Perot and Chamel, 2021](#))

$$k_2 = \frac{8}{5}C^5(1-2C)^2[2(y_2-1)C - y_2 + 2] \left\{ 2C[4(y_2+1)C^4 + 2(3y_2-2)C^3 - 2(11y_2-13)C^2 + 3(5y_2-8)C - 3(y_2-2)] + 3(1-2C)^2[2(y_2-1)C - y_2 + 2] \log(1-2C) \right\}^{-1}, \quad (3.15)$$

$$k_3 = \frac{8}{7}C^7(1-2C)^2[2(y_3-1)C^2 - 3(y_3-2)C + y_3 - 3] \times \left\{ 2C[4(y_3+1)C^5 + 2(9y_3-2)C^4 - 20(7y_3-9)C^3 + 5(37y_3-72)C^2 - 45(2y_3-5)C + 15(y_3-3)] + 15(1-2C)^2[2(y_3-1)C^2 - 3(y_3-2)C + y_3 - 3] \log(1-2C) \right\}^{-1}, \quad (3.16)$$

$$k_4 = \frac{32}{147}C^9(1-2C)^2[12(y_4-1)C^3 - 34(y_4-2)C^2 + 28(y_4-3)C - 7(y_4-4)] \times \left\{ 2C[8(y_4+1)C^6 + 4(17y_4-2)C^5 - 12(83y_4-107)C^4 + 40(55y_4-116)C^3 - 10(191y_4-536)C^2 + 105(7y_4-24)C - 105(y_4-4)] + 15(1-2C)^2[12(y_4-1)C^3 - 34(y_4-2)C^2 + 28(y_4-3)C - 7(y_4-4)] \log(1-2C) \right\}^{-1}, \quad (3.17)$$

$$j_2 = \frac{24}{5}C^5[2(\tilde{y}_2-2)C - \tilde{y}_2 + 3] \left\{ 2C[2(\tilde{y}_2+1)C^3 + 2\tilde{y}_2C^2 + 3(\tilde{y}_2-1)C - 3(\tilde{y}_2-3)] + 3[2(\tilde{y}_2-2)C - \tilde{y}_2 + 3] \log(1-2C) \right\}^{-1}, \quad (3.18)$$

$$j_3 = \frac{64}{21}C^7[8(\tilde{y}_3-2)C^2 - 10(\tilde{y}_3-3)C + 3(\tilde{y}_3-4)] \times \left\{ 2C[4(\tilde{y}_3+1)C^4 + 10\tilde{y}_3C^3 + 30(\tilde{y}_3-1)C^2 - 15(7\tilde{y}_3-18)C + 45(\tilde{y}_3-4)] + 15[8(\tilde{y}_3-2)C^2 - 10(\tilde{y}_3-3)C + 3(\tilde{y}_3-4)] \log(1-2C) \right\}^{-1}, \quad (3.19)$$

$$j_4 = \frac{80}{147}C^9[40(\tilde{y}_4-2)C^3 - 90(\tilde{y}_4-3)C^2 + 63(\tilde{y}_4-4)C - 14(\tilde{y}_4-5)] \times \left\{ 2C[4(\tilde{y}_4+1)C^5 + 18\tilde{y}_4C^4 + 90(\tilde{y}_4-1)C^3 - 5(137\tilde{y}_4-334)C^2 + 105(7\tilde{y}_4-26)C - 210(\tilde{y}_4-5)] + 15[40(\tilde{y}_4-2)C^3 - 90(\tilde{y}_4-3)C^2 + 63(\tilde{y}_4-4)C - 14(\tilde{y}_4-5)] \log(1-2C) \right\}^{-1}$$

Chapter 4

Results and Discussions




Figure 4.1: Proton fraction x_p of β -stable NM at different baryon density and spin polarity for CDM3Y n interactions.




Figure 4.2: Total mass-energy density of β -stable NM at varying spin polarity with different interaction models.




Figure 4.3: Total pressure of β -stable NM at several values of Δ with different CDM3Y n models.




Figure 4.4: The relation between gravitational mass M and the radius R of the NS according to the corresponding model and polarity.






Figure 4.5: GE tidal Love number at 2nd order k_2 as function of NS mass computed with CDM3Y n models at different spin polarities.

Figure 4.6: Same as Figure 4.5 for k_3 .Figure 4.7: Same as Figure 4.5 for k_4 .




Figure 4.8: GM tidal Love number at 2nd order j_2 as function of NS mass computed with CDM3Yn models of *strictly static fluid* at different polarities.








Figure 4.9: Same as Figure 4.8 for j_3 .

Figure 4.10: Same as Figure 4.8 for j_4 .Figure 4.11: GM tidal Love number at 2nd order j_2 as function of NS mass computed with CDM3Y n models of *irrotational fluid* at different polarities.

Figure 4.12: Same as Figure 4.11 for j_3 .Figure 4.13: Same as Figure 4.11 for j_4 .

Chapter 5

Conclusions

Bibliography

- Abbott, B., Abbott, R., Abbott, T., Abraham, S., Acernese, F., Ackley, K., Adams, C., Adhikari, R., Adya, V., Affeldt, C., et al. (2020). GW190425: observation of a compact binary coalescence with total mass $\sim 3.4M_{\odot}$. *The Astrophysical Journal Letters*, 892(1):L3.
- Abbott, B. P., Abbott, R., Abbott, T., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R., Adya, V., et al. (2017). GW170817: observation of gravitational waves from a binary neutron star inspiral. *Physical Review Letters*, 119(16):161101.
- Abbott, B. P., Abbott, R., Abbott, T., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R. X., Adya, V. B., et al. (2018). GW170817: Measurements of neutron star radii and equation of state. *Physical review letters*, 121(16):161101.
- Akmal, A., Pandharipande, V., and Ravenhall, D. a. (1998). Equation of state of nucleon matter and neutron star structure. *Physical Review C*, 58(3):1804.
- Anantaraman, N., Toki, H., and Bertsch, G. (1983). An effective interaction for inelastic scattering derived from the Paris potential. *Nuclear Physics A*, 398(2):269–278.
- Baym, G. and Pethick, C. (1975). Neutron stars. *Annual Review of Nuclear Science*, 25(1):27–77.
- Damour, T. and Nagar, A. (2009). Relativistic tidal properties of neutron stars. *Physical Review D*, 80(8):084035.
- Gandolfi, S., Illarionov, A. Y., Fantoni, S., Miller, J., Pederiva, F., and Schmidt, K. (2010). Microscopic calculation of the equation of state of nuclear matter and neutron star structure. *Monthly Notices of the Royal Astronomical Society: Letters*, 404(1):L35–L39.
- Glendenning, N. K. (2012). *Compact stars: Nuclear physics, particle physics and general relativity*. Springer Science & Business Media.
- Greiner, W. and Maruhn, J. A. (1996). *Nuclear models*. Springer.
- Hinderer, T. (2008). Tidal Love numbers of neutron stars. *The Astrophysical Journal*, 677(2):1216.
- Hinderer, T., Lackey, B. D., Lang, R. N., and Read, J. S. (2010). Tidal deformability of neutron stars with realistic equations of state and their gravitational wave signatures in binary inspiral. *Physical Review D*, 81(12):123016.

- Khoa, D. T. and Cuong, D. C. (2007). Mean-field description of the nucleus-nucleus optical potential. *arXiv preprint arXiv:0708.3711*.
- Khoa, D. T. and Satchler, G. (2000). Generalized folding model for elastic and inelastic nucleus–nucleus scattering using realistic density dependent nucleon–nucleon interaction. *Nuclear Physics A*, 668(1-4):3–41.
- Khoa, D. T., Satchler, G., and Von Oertzen, W. (1997). Nuclear incompressibility and density dependent NN interactions in the folding model for nucleus-nucleus potentials. *Physical Review C*, 56(2):954.
- Khoa, D. T., Von Oertzen, W., Bohlen, H., Bartnitzky, G., Clement, H., Sugiyama, Y., Gebauer, B., Ostrowski, A., Wilpert, T., Wilpert, M., et al. (1995). Equation of State for Cold Nuclear Matter from Refractive O 16+ O 16 Elastic Scattering. *Physical review letters*, 74(1):34.
- Lattimer, J. M. and Prakash, M. (2004). The physics of neutron stars. *science*, 304(5670):536–542.
- Le Tiec, A. and Casals, M. (2021). Spinning black holes fall in love. *Physical Review Letters*, 126(13):131102.
- Loan, D. T., Tan, N. H., Khoa, D. T., and Margueron, J. (2011). Equation of state of neutron star matter, and the nuclear symmetry energy. *Physical Review C*, 83(6):065809.
- Moustakidis, C. C. and Panos, C. (2009). Equation of state for β -stable hot nuclear matter. *Physical Review C*, 79(4):045806.
- Perot, L. and Chamel, N. (2021). Role of dense matter in tidal deformations of inspiralling neutron stars and in gravitational waveforms with unified equations of state. *Physical Review C*, 103(2):025801.
- Tan, N. H., Khoa, D. T., and Loan, D. T. (2020). Spin-polarized β -stable neutron star matter: The nuclear symmetry energy and GW170817 constraint. *Physical Review C*, 102(4):045809.
- Tan, N. H., Khoa, D. T., and Loan, D. T. (2021). Equation of state of asymmetric nuclear matter and the tidal deformability of neutron star. *arXiv preprint arXiv:2104.09121*.
- Tan, N. H., Loan, D. T., Khoa, D. T., and Margueron, J. (2016). Mean-field study of hot β -stable protoneutron star matter: Impact of the symmetry energy and nucleon effective mass. *Physical Review C*, 93(3):035806.
- Thân, H. S. (2010). *UFR SCIENTIFIQUE D’ORSAY INSTITUT DE PHYSIQUE NUCLEAIRE D’ORSAY*. PhD thesis, Citeseer.
- Vidana, I. and Bombaci, I. (2002). Equation of state and magnetic susceptibility of spin polarized isospin asymmetric nuclear matter. *Physical Review C*, 66(4):045801.