

UNCERTAINTY ON THE GALAXY-HALO CONNECTION FOR LYMAN- α EMITTERS AT $Z = 3.1$

JULIÁN E. MEJÍA-RESTREPO^{1,3}, JAIME E. FORERO-ROMERO²

¹Departamento de Astronomía, Universidad de Chile, Camino el Observatorio 1515, Santiago, Chile

²Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio Ip, Bogotá, Colombia

³FACom-Instituto de Física-FCEN, Universidad de Antioquia, Calle 70 No. 52-21, Medellín, Colombia

Submitted for publication in *ApJ*

ABSTRACT

We study the impact of cosmic variance and observational uncertainties in constraining the mass and occupation fraction (f_{occ}) of dark matter halos hosting Ly α Emitting Galaxies at high redshift (LAEs). We use a N-body simulation to construct mock catalogs with the same typical size of observed fields at $z = 3.1$ ($\sim 1\text{deg}^2$) to match the observed angular correlation function (ACF) and number density of LAEs. In our model a dark matter halo with mass in the range $M_{\text{min}} < M_h < M_{\text{max}}$ can only host one detectable LAE at most. By following a thorough Markov Chain Monte-Carlo exploration of the parameter space determined by M_{min} and M_{max} , our analysis shows that f_{occ} is uniquely determined by M_{min} regardless of M_{max} using the relation $f_{\text{occ}} \sim 0.1 (M_{\text{min}}/10^{10.5})^{0.93}$. However, the current observational data only allows us to put weak constraints on M_{min} , M_{max} and consequently on f_{occ} . Particularly, we find that $10^{9.6}h^{-1}M_{\odot} \leq M_{\text{min}} \leq 10^{11.0}h^{-1}M_{\odot}$, $10^{10.9}h^{-1}M_{\odot} \leq M_{\text{max}} \leq 10^{13.0}h^{-1}M_{\odot}$ and $0.02 \leq f_{\text{occ}} \leq 0.30$. Nevertheless, we also find that the upcoming large surveys can help to put tighter constraints on M_{min} and f_{occ} through the width measurement of the LAE number distribution function obtained over several fields of the same size of current observations. Analogously, the improvement in the precision of the measured ACF will assist in constraining M_{max} .

Subject headings: Cosmology: theory - large-scale structure of Universe - Methods: data analysis - numerical - N-body simulations

1. INTRODUCTION

(●REFS)ADD NEWER REFERENCES Lyman- α emitting galaxies (LAEs) are central to a wide range of subjects in extragalactic astronomy. LAEs can be used as probes of reionization (Dijkstra et al. 2011), tracers of large scale structure (Koehler et al. 2007), signposts for low metallicity stellar populations, markers of the galaxy formation process at high redshift (Dayal et al. 2009; Forero-Romero et al. 2012) and tracers of active star formation (Guaita et al. 2013).

In most of those cases, capitalizing the observations requires understanding how LAEs are formed within an explicit cosmological context. Under the current structure formation paradigm the dominant matter content of the Universe is dark matter (DM). Each galaxy is thought to be hosted by a larger dark matter structure known as a halo. (Peebles 1980; Springel et al. 2005). Understanding the cosmological context of LAEs thus implies studying the galaxy-halo connection. Galaxy formation models suggest that the physical processes that regulate the star formation cycle are dependent on halo mass (Behroozi et al. 2013). The mass becomes the most important element in the halo-galaxy connection.

The goal becomes finding the typical DM halo mass of halos hosting LAEs. In the case of LAEs there are different ways to find this mass range. One approach is theoretical, using general astrophysical principles to find the relationship between halo mass, intrinsic Ly α luminosities and observed Ly α luminosities. This approach is usually implemented through semi-analytic models (Garel et al. 2012; Orsi et al. 2012) and full N-body hydrodynamical simulations (Laursen & Sommer-Larsen 2007; Dayal et al. 2009; Forero-Romero et al. 2011; Ya-

jima et al. 2012).

The downside of these calculations is the uncertainty in the estimation of the escape fraction of Ly α photons. Given the resonant nature of the Ly α line, the escape fraction is sensitive to the dust contents, density, temperature, topology and kinematics of the neutral Hydrogen in the interstellar medium (ISM). The process of finding a consensus on the expected value for the Ly α escape fraction in high redshift galaxies is still matter of open debate (Neufeld 1991; Verhamme et al. 2006; Forero-Romero et al. 2011; Dijkstra & Kramer 2012; Laursen et al. 2013; Orsi et al. 2012).

A different approach to infer the typical mass of halos hosting LAEs is based on the spatial clustering information. This approach uses the fact that in CDM cosmologies the spatial clustering of galaxies on large scales is entirely dictated by the halo distribution (Colberg et al. 2000), which in turn has a strong dependence on halo mass. Using measurements of the angular correlation function of LAEs, observers have put constraints on the typical mass and occupation fraction of the putative halos hosting these galaxies (Hayashino et al. 2004; Gawiser et al. 2007b; Nilsson et al. 2007; Ouchi et al. 2010; Bielby et al. 2016). In these studies the observations are done on fields of $\sim 1\text{deg}^2$ and the conclusions derived on the halo host mass do not elaborate on the uncertainty resulting from the cosmic variance on these fields.

In this letter we investigate the impact of cosmic variance in constraining the mass and occupation fraction of halos hosting LAEs at $z = 3$. We build mock surveys from a cosmological N-body to compare them against the observations in Bielby et al. (2016) using the angular correlation function. We use a simple model to populate a




FIG. 1.— Cumulative halo number density distribution function over 27 mock fields. Each line corresponds to a different model with increasing values of M_{\min} . Different models produce different number density distributions. The width increases with M_{\min} . halo in the simulation with a LAE assuming a minimum (M_{\min}) and maximum mass (M_{\max}) for the dark matter halos hosting LAEs without predicting a Ly α luminosity. This approach bypasses all the physical uncertainties associated to star formation and radiative transfer. We use the Markov Chain Monte Carlo technique to obtain the likelihood of the parameters given the observational constraints.

Throughout this letter we assume a Λ CDM cosmology with the following values for the cosmological parameters, $\Omega_m = 0.30711$, $\Omega_\Lambda = 0.69289$ and $h = 0.70$, corresponding to the matter density, vacuum density and the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2. METHODOLOGY

The base of our method is the comparison between observations and mock catalogs. This approach allows us to take explicitly into account cosmic variance. The comparison has four key elements. First, the observations we take as a benchmark. Second, the N-body simulation and the halo catalogs we use to build the mocks. Third, the parameters describing our model to assign a LAE to a halo. Fourth, the statistical method we adopt to compare observations and simulations. We describe in detail these four elements in the following subsections.

2.1. Observational constraints

Bielby et al. (2016) used narrow band imaging to detect 643 LAE candidates at $z \sim 3$ with equivalent widths of $\gtrsim 65\text{\AA}$ and a flux limit of $2 \times 10^{17} \text{ erg/cm}^2/\text{s}$ ($L \sim 7 \times 10^{42} \text{ erg/s}$). Using spectroscopy they found a 22% contamination fraction. Their observations cover 5 (out of 9) independent and co-spatial fields of the VLT LBG Redshift Survey (VLRS). The total observed area corresponds to 1.07 deg^2 that translates to $\sim 80^2 h^{-1} \text{ Mpc}^2$ in a comoving scale. Bielby et al. (2016) used the NB497 narrow-band filter whose 77\AA FWHM and 154\AA FWTM correspond to a total observational depth of $44 h^{-1} \text{ Mpc}$ and $82 h^{-1} \text{ Mpc}$ comoving, respectively.

2.2. Simulation and halo catalog

We use results from the Bolshoi simulation (Klypin et al. 2011, 2014) which was performed in a cubic volume of $250 h^{-1} \text{ Mpc}$ comoving on a side. The dark matter distribution was sampled using 2048^3 particles. The cosmological parameters are consistent with Planck results (Planck Collaboration et al. 2014) with a matter density $\Omega_m = 0.307$, cosmological constant $\Omega_\Lambda = 0.693$, dimensionless Hubble constant $h = 0.678$, slope of the power spectrum $n = 0.96$ and normalization of the power spectrum $\sigma_8 = 0.823$. This translates into a particle mass of $m_p = 1.5 \times 10^8 h^{-1} \text{ M}_\odot$.

We use halo catalogs constructed with a Bound-Density-Maxima (BDM) algorithm. The catalogs were obtained from the publicly available Multidark database¹ (Riebe et al. 2013). For each halo in the box we extract its comoving position and mass. We focus our work on halos more massive than $1.5 \times 10^9 h^{-1} \text{ M}_\odot$ resolved with at least 10 particles. We do not take into account sub-halos.

We split the simulation volume at $z \sim 3$ into 27 smaller mock volumes mimicking the area and depth reported in Bielby et al. (2016) and described in §2.1.

2.3. A model to populate halos with LAEs

We build the simplest possible model to assign a LAE to each DM halo without trying to compute a LAE luminosity.

We first assume that a dark matter halo can host one detectable LAE at most. This assumption is consistent with theoretical analysis of the correlation function (Jose et al. 2013) and observations that confirm a lack of class pairs in LAEs Bond et al. (2009). Then we say that a halo will host a LAE with probability f_{occ} if and only if the halo mass is in the range $M_{\min} < M_h < M_{\max}$. In what follows we refer to a given set of values f_{occ} , M_{\min} , M_{\max} as a model, \mathcal{M} . We also use a variable ΔM to represent the mass range width, $\Delta M \equiv \log M_{\max} - \log M_{\min}$.

The parameter f_{occ} can be thought as the LAE occupation fraction of dark matter halos. An astrophysical interpretation of f_{occ} convolves at least three phenomena: the actual presence of a star forming galaxy in a halo, the escape fraction of Ly α radiation and its detectability as a LAE. We do not try to disentangle these complex phenomena. We instead opt for an arithmetic interpretation by setting f_{occ} as the ratio between the observed number of LAEs and the number of halos within the considered mass range, that is $f_{\text{occ}} \equiv N_{\text{LAEs}}/N_{\text{halos}}$.

Fig. 1 shows the cumulative halo number density for all 27 subvolumes in the simulation, with a normalization by the average number density among fields. Each line represents a different model \mathcal{M} with fixed $f_{\text{occ}} = 1$ and $\Delta M = 1.0$; and varying M_{\min} . This Figure shows that the halo number density varies across sub-volumes, as an expression of cosmic variance. As a consequence, the f_{occ} also varies across the mock fields by the same factor.

We do not explore any physical model to disentangle these two effects. This means that our model does not assign a luminosity or escape fraction to each LAE. We avoid this in order to maintain theoretical uncertainties

¹ <http://www.multidark.org/MultiDark/>




FIG. 2.— One and two dimensional projections of the posterior probability distributions of M_{\min} , M_{\max} and f_{occ} . The models with $\log f_{\text{occ}} > 0.00$ ($f_{\text{occ}} > 1$) correspond to models where the number density of halos is smaller than the number of observed LAEs but we consider them as consistent because of the uncertainty in the median number density due to cosmic variance. See Fig. 1 and §2.4 for details.

to a minimum. This flexibility allows us to explore a wide range of possible masses for the host halos without any strong theoretical prejudice regarding the details of star formation and Ly α escape fraction.

We also artificially contaminate our mock catalogs with 22% randomly distributed data points to mimic the fraction of interlopers in observations Bielby et al. (2016). On top of that we apply rejection sampling to our LAE selection taking the transmission function of the NB479 filter used in their observations as a radial selection factor.

In what follows we note by the letter \mathcal{M} a model defined by a particular choice of the two parameters M_{\min} , M_{\max} . For each model \mathcal{M} we define \tilde{f}_{occ} as the median occupation fraction within the the mock fields and $\Delta M \equiv M_{\max} - M_{\min}$.

2.4. Exploring and selecting consistent models

We use the angular correlation function to compare the observations with our mock catalogs. This is done through a thorough exploration of the parameter space of the models \mathcal{M} by means of a Monte Carlo Markov Chain minimization using the EMCEE python package (Foreman-Mackey et al. 2013). We put a flat prior on $\log M_{\min}$ and $\log M_{\max}$ to vary between 9.2 up to 13.4, corresponding to the halo mass range of the simulation at $z = 3$. Given that the typical scatter in N_{halos} is about 0.3 dex (i.e. about a factor of 2), our parameter space is restricted to models where the median number of dark matter halos is larger than $N_{\text{LAE}}/3$. Because of this possibility of having N_{halos} smaller than N_{LAE} there will be some models where $f_{\text{occ}} \equiv N_{\text{LAE}}/N_{\text{halo}} > 1$. Those models only reflect the uncertainty in the number density of LAEs due to cosmic variance.




FIG. 3.— $\log M_{\min}$ vs $\log M_{50\%}$ (black), $\log M_{16\%}$ - $\log M_{84\%}$ (blue) and $\log M_{2.5\%}$ - $\log M_{97.5\%}$ (red) for different values of $\log M_{\max}$. The points are color coded by $\Delta \log M$ [$M_{\odot} h^{-1}$]. The green line diagonal line accounts for the 1:1 relation and the green vertical lines represent the 16, 50 and 84 percentiles of the posterior probability distribution of M_{\min} .

The MCMC exploration is done using a total of 24 seeds and 400 iterations (9600 models) to sample the posterior PDF, $P(M_{\min}, M_{\max}, f_{\text{occ}} | \mathcal{M}) \propto \exp(-\chi^2_{\mathcal{M}}/2)$, with

$$\chi^2_{\mathcal{M}} = \sum_{\theta} \left[\frac{(\text{ACF}_{\mathcal{M}}(\theta) - \text{ACF}_{\text{obs}}(\theta))^2}{\sigma^2_{\mathcal{M}}(\theta) + \sigma^2_{\text{obs}}(\theta)} \right] \quad (1)$$

where $\text{ACF}_{\mathcal{M}}$ and ACF_{obs} are the ACF of the explored model \mathcal{M} and the observational ACF reported by Bielby et al. (2016) respectively. $\sigma_{\mathcal{M}}$ is the associated 1- σ scatter of the $\text{ACF}_{\mathcal{M}}$ as a product of cosmic variance and σ_{obs} is the observational error associated to ACF_{obs} . The $\text{ACF}_{\mathcal{M}}$ s are computed using the Landy & Szalay estimator (Landy & Szalay 1993).

3. RESULTS AND DISCUSSION

3.1. Constraining M_{\min} , M_{\max} and f_{occ} in Dark Matter Halos hosting LAEs

In Fig. 2 we present the one and two dimensional projections of the posterior probability distributions of M_{\min} , M_{\max} and f_{occ} . We find that the M_{\min} and M_{\max} parameter space cannot be tightly constrained mainly because of cosmic variance and the large observational Poissonian uncertainty in the ACF. We however found that $9.6 < \log M_{\min} < 11.0$ and $10.9 < \log M_{\max} < 13.0$. We interestingly find that f_{occ} is basically completely determined by M_{\min} going from $f_{\text{occ}}=0.015$ when $\log M_{\min}=9.6$ to $f_{\text{occ}}=0.30$ when $\log M_{\min}=11.0$ ($f_{\text{occ}} \sim 0.1 (M_{\min}/10^{10.5})^{0.93}$). We remark that those models where $\log f_{\text{occ}} > 0.00$ ($f_{\text{occ}} > 1$) correspond to models where the number density of halos is smaller than the number of observed LAEs but are still consistent because of the uncertainty in the median number density of LAEs in the universe due to cosmic variance (see Fig. 1 and §2.4).

3.2. Halo mass distribution within models

Previous attempts to constrain the DMH mass of LAEs by means of the ACF present their result in terms of the median mass of LAEs that is derived from the derived

correlation length by means of semi-analytical prescriptions (e.g. Hayashino et al. 2004; Gawiser et al. 2007a; Ouchi et al. 2010; Bielby et al. 2016). Instead, in this work we directly compare the ACF of our simulated mock catalogs with the observational ACF and present our results in terms of a minimum and maximum mass DMH hosting LAEs. It is therefore necessary to connect our approach by computing the median mass of the DMH in each of the models \mathcal{M} . In Fig. 3 we show the 50 ($\log M_{50}$, black dots), the 84 ($\log M_{84}$, blue diamonds) and 95 ($\log M_{95}$, red squares) percentiles of the LAE halo mass as a function of $\log M_{\min}$ for each of the models that we run in our MCMC simulation. The points are color coded according to their $\Delta \log M$ associated value. We can see that the median mass (M_{50}) and M_{84} are not very sensitive to $\log M_{\max} = \log M_{\min} + \Delta \log M$ specially when $\Delta \log M \gtrsim 1.0\text{dex}$. We particularly found that $\log M_{\min} \lesssim \log M_{50} \lesssim \log M_{\min} + 0.2\text{dex}$ and that $\log M_{\min} + 0.1\text{dex} \lesssim \log M_{50} \lesssim \log M_{\min} + 0.5\text{dex}$ regardless of the value of $\Delta \log M$. The latter is a consequence of the very steepen distribution of the dark matter mass function and is at the same time the reason for the almost perfect one to one relation between M_{\min} and f_{occ} . However, it can also be seen in Fig. 3 that $\log M_{84}$ is very sensitive to $\log M_{\max}$ ($\log M_{\min} + 0.2\text{dex} \lesssim \log M_{50} \lesssim \log M_{\min} + 1.5\text{dex}$). Thereby, any differences in the clustering strength of models sharing the same M_{\min} but different M_{\max} should be mainly driven by the $\sim 16\%$ most massive halos of each model \mathcal{M} .

In order to estimate the effect of most massive halos in $\text{ACF}_{\mathcal{M}}$ in Fig. 4 we show the computed $\text{ACF}_{\mathcal{M}}$ of models with $\log M_{\min} = 0.5$ and different values of $\Delta \log M$. We can see that the clustering gets stronger for larger values of $\Delta \log M$. Nevertheless, due to the large impact of cosmic variance at the volume of the current observations all the models are basically consistent within errors. The last result together with the large Poissonian observational error in the ACF explain the current difficulty to put tighter constraints in $\log M_{\max}$ in our model.

In Fig. 3 the green vertical lines represent the 16, 50 and 84 percentiles of the posterior probability distribution of M_{\min} that we previously obtained. From this figure we find that the median dark matter halo mass is $\tilde{M}_h = 10^{10.6^{+0.7}_{-0.6}}$. This result is consistent with previous estimations of the median dark matter halo mass reported by Bielby et al. (2016) ($\tilde{M}_h = 10^{11.0 \pm 0.6}$), Gawiser et al. (2007b) ($\tilde{M}_h = 10^{10.9 \pm 0.9}$) and Ouchi et al. (2010) ($\tilde{M}_h = 6.7^{+42.0}_{-6.7} \times 10^{10}$) using semi-analytical approaches.

3.3. Constraining Dark matter halos mass with cosmic variance

Fig. 1 shows the halo number distribution (HND) in the mock fields of the simulation for different models \mathcal{M} . By simple inspection one can infer an increase in the distribution with as $\log M_{\min}$ increase. This trend is confirmed in Fig. 5 where we plot the central 1- σ (blue diamonds) and 2- σ (red dots) widths of the NDH as a function of $\log M_{\min}$. We particularly found that when we consider survey fields of $\sim 1\text{deg}^2$, the central 1- σ (2- σ) width of the HND increases monotonically from 0.05dex (0.10dex) when $\log M_{\min} = 9.5$ to 0.20dex (0.35dex) when $\log M_{\min} = 12.0$. The latter result opens the possibility to constrain the $\log M_{\min}$ (as well as the median




FIG. 4.— Angular Correlation functions for $\log M_{\min}[\text{M}_{\odot}h^{-1}] = 10.5$ and different values of ΔM . The shaded region represents the 1-sigma deviation due to cosmic variance. Radical different models in ΔM are consistent with observations once cosmic variance is modelled in detail.




FIG. 5.— Width of the halo number distribution over the 54 mock field of the simulation ($\Delta \log(N_{\text{halos}}/\bar{N}_{\text{halos}})$) of the central 68 (blue diamonds) and 95 (red circles) percentiles vs M_{\min} . Points are color coded according to their associated value of $\Delta M \equiv M_{\min} - M_{\text{min}}$. The green line diagonal line accounts for the 1:1 relation and the green vertical lines represent the 16, 50 and 84 percentiles of the posterior probability distribution of M_{\min} .

mass) of LAEs by simply measure the width of the distribution of observed LAE along several observational fields mapping $\sim 1\text{deg}^2$ in area and the observational depth determined by the NB479 filter.

4. CONCLUSIONS

In this letter we studied the impact of cosmic variance and observational uncertainties in constraining the

mass range and occupation fraction of dark matter halos hosting LAEs. To this end, we used the BolshoiP N-body simulation to construct 27 mock fields with the same typical size of observed fields at $z = 3.1$ ($\sim 1\text{deg}^2$). In our model a dark matter halo with mass in the range $M_{\min} < M_h < M_{\max}$ can only host one detectable LAE at most. We explored the parameter space determined by M_{\min} and M_{\max} using Monte Carlo Markov-Chain minimization to match the observed ACF and mean number density of LAEs.

Our analysis allowed us to put weak constraints on M_{\min} , M_{\max} and f_{occ} where $10^{9.7}h^{-1}\text{M}_{\odot} \leq \log M_{\min} \leq 10^{11.2}h^{-1}\text{M}_{\odot}$, $10^{10.9}h^{-1}\text{M}_{\odot} \leq \log M_{\max} \leq 10^{13.0}h^{-1}\text{M}_{\odot}$ and $0.02h^{-1}\text{M}_{\odot} \leq f_{\text{occ}} \leq 0.5$, spanning three orders of magnitude in halo mass. Previous works (Hayashino et al. 2004; Gawiser et al. 2007b; Ouchi et al. 2008; Bielby et al. 2016) have found typical masses within somehow narrower mass ranges ($\sim 10^{10.5}-10^{12}$). The main reason for our weaker constraints and large discrepancies with previous works resides in the cosmic variance on the typical field size in observations. A thorough exploration of cosmic variance impact as we present in this letter had not been presented so far in the literature.

Our analysis also allowed us to draw three results that can be used to put tighter constraints on M_{\min} , M_{\max} and f_{occ} once upcoming large LAE surveys, as the HETDEX project (Adams et al. 2011) and the HSC ultra deep survey, are available :

1. f_{occ} and the median mass of LAEs are almost uniquely determined by M_{\min} regardless of M_{\max} .
2. measuring the width of the LAE number distribution function obtained over several fields of $\approx 1\text{deg}^2$ one will be able to tightly constrain M_{\min} , M_{med} and f_{occ} up to a factor of 2.
3. M_{\max} drives the ACF strength at short angular distances ($< 1\text{arcmin}$), therefore precise measurements of the ACF at such scales are crucial to accurately determine M_{\max} .

ACKNOWLEDGMENTS

JM acknowledges “CONICYT-PCHA/doctorado Nacional para extranjeros/2013-63130316” for their PhD scholarship support.

The authors gratefully acknowledge the Gauss Centre for Supercomputing e.V. (www.gauss-centre.eu) and the Partnership for Advanced Supercomputing in Europe (PRACE, www.prace-ri.eu) for funding the MultiDark simulation project by providing computing time on the GCS Supercomputer SuperMUC at Leibniz Supercomputing Centre (LRZ, www.lrz.de). The Bolshoi simulations have been performed within the Bolshoi project of the University of California High-Performance Astro-Computing Center (UC-HiPACC) and were run at the NASA Ames Research Center.

REFERENCES

- Adams, J. J., Blanc, G. A., Hill, G. J., Gebhardt, K., Drory, N., Hao, L., Bender, R., Byun, J., Ciardullo, R., Cornell, M. E., Finkelstein, S. L., Fry, A., Gawiser, E., Gronwall, C., Hopp, U., Jeong, D., Kelz, A., Kelzenberg, R., Komatsu, E., MacQueen, P. J., Murphy, J., Odoms, P. S., Roth, M., Schneider, D. P., Tufts, J. R., & Wilkinson, C. P. 2011, *ApJS*, 192, 5
- Behroozi, P. S., Wechsler, R. H., & Conroy, C. 2013, *ApJ*, 770, 57
- Bielby, R. M., Tummuangpak, P., Shanks, T., Francke, H., Crighton, N. H. M., Bañados, E., González-López, J., Infante, L., & Orsi, A. 2016, *MNRAS*, 456, 4061
- Bond, N. A., Gawiser, E., Gronwall, C., Ciardullo, R., Altmann, M., & Schawinski, K. 2009, *ApJ*, 705, 639

- Colberg, J. M., White, S. D. M., Yoshida, N., MacFarland, T. J., Jenkins, A., Frenk, C. S., Pearce, F. R., Evrard, A. E., Couchman, H. M. P., Efstathiou, G., Peacock, J. A., Thomas, P. A., & Virgo Consortium. 2000, *MNRAS*, 319, 209
- Dayal, P., Ferrara, A., Saro, A., Salvaterra, R., Borgani, S., & Tornatore, L. 2009, *MNRAS*, 400, 2000
- Dijkstra, M., & Kramer, R. 2012, *MNRAS*, 424, 1672
- Dijkstra, M., Mesinger, A., & Wyithe, J. S. B. 2011, *MNRAS*, 414, 2139
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Forero-Romero, J. E., Yepes, G., Gottlöber, S., Knollmann, S. R., Cuesta, A. J., & Prada, F. 2011, *MNRAS*, 415, 3666
- Forero-Romero, J. E., Yepes, G., Gottlöber, S., & Prada, F. 2012, *MNRAS*, 419, 952
- Garel, T., Blaizot, J., Guiderdoni, B., Schaerer, D., Verhamme, A., & Hayes, M. 2012, *MNRAS*, 422, 310
- Gawiser, E., Francke, H., Lai, K., Schawinski, K., Gronwall, C., Ciardullo, R., Quadri, R., Orsi, A., Barrientos, L. F., Blanc, G. A., Fazio, G., & Feldmeier, J. J. 2007a, *ApJ*, 671, 278
- Gawiser, E., Francke, H., Lai, K., Schawinski, K., Gronwall, C., Ciardullo, R., Quadri, R., Orsi, A., Barrientos, L. F., Blanc, G. A., Fazio, G., Feldmeier, J. J., Huang, J.-s., Infante, L., Lira, P., & Padilla, N. 2007b, *ApJ*, 671, 278
- Guaita, L., Francke, H., Gawiser, E., Bauer, F. E., Hayes, M., Östlin, G., & Padilla, N. 2013, *A&A*, 551, A93
- Hayashino, T., Matsuda, Y., Tamura, H., Yamauchi, R., Yamada, T., Ajiki, M., Fujita, S. S., Murayama, T., Nagao, T., Ohta, K., Okamura, S., Ouchi, M., Shimasaku, K., Shioya, Y., & Taniguchi, Y. 2004, *AJ*, 128, 2073
- Jose, C., Srianand, R., & Subramanian, K. 2013, *ArXiv e-prints*
- Klypin, A., Yepes, G., Gottlober, S., Prada, F., & Hess, S. 2014, *ArXiv e-prints*
- Klypin, A. A., Trujillo-Gomez, S., & Primack, J. 2011, *ApJ*, 740, 102
- Koehler, R. S., Schuecker, P., & Gebhardt, K. 2007, *A&A*, 462, 7
- Landy, S. D., & Szalay, A. S. 1993, *ApJ*, 412, 64
- Laursen, P., Duval, F., & Östlin, G. 2013, *ApJ*, 766, 124
- Laursen, P., & Sommer-Larsen, J. 2007, *ApJ*, 657, L69
- Neufeld, D. A. 1991, *ApJ*, 370, L85
- Nilsson, K. K., Möller, P., Möller, O., Fynbo, J. P. U., Michałowski, M. J., Watson, D., Ledoux, C., Rosati, P., Pedersen, K., & Grove, L. F. 2007, *A&A*, 471, 71
- Orsi, A., Lacey, C. G., & Baugh, C. M. 2012, *MNRAS*, 425, 87
- Ouchi, M., Shimasaku, K., Akiyama, M., Simpson, C., Saito, T., Ueda, Y., Furusawa, H., Sekiguchi, K., Yamada, T., Kodama, T., Kashikawa, N., Okamura, S., Iye, M., Takata, T., Yoshida, M., & Yoshida, M. 2008, *ApJS*, 176, 301
- Ouchi, M., Shimasaku, K., Furusawa, H., Saito, T., Yoshida, M., Akiyama, M., Ono, Y., Yamada, T., Ota, K., Kashikawa, N., Iye, M., Kodama, T., Okamura, S., Simpson, C., & Yoshida, M. 2010, *ApJ*, 723, 869
- Peebles, P. J. E. 1980, *The large-scale structure of the universe*
- Planck Collaboration, Ade, P. A. R., Aghanim, N., Armitage-Caplan, C., Arnaud, M., Ashdown, M., Atrio-Barandela, F., Aumont, J., Baccigalupi, C., Banday, A. J., & et al. 2014, *A&A*, 571, A16
- Riebe, K., Partl, A. M., Enke, H., Forero-Romero, J., Gottlber, S., Klypin, A., Lemson, G., Prada, F., Primack, J. R., Steinmetz, M., & Turchaninov, V. 2013, *Astronomische Nachrichten*, 334, 691
- Springel, V., White, S. D. M., Jenkins, A., Frenk, C. S., Yoshida, N., Gao, L., Navarro, J., Thacker, R., Croton, D., Helly, J., Peacock, J. A., Cole, S., Thomas, P., Couchman, H., Evrard, A., Colberg, J., & Pearce, F. 2005, *Nature*, 435, 629
- Verhamme, A., Schaerer, D., & Maselli, A. 2006, *A&A*, 460, 397
- Yajima, H., Choi, J.-H., & Nagamine, K. 2012, *MNRAS*, 427, 2889