

Pú ert jörðin

JEFR¹, MCRG¹, JNGC², MD³

¹ Bogota

² Tucson

³ Oslo

Star-forming Compact Dwarf Galaxies (CDGs) resemble the expected pristine conditions of the first galaxies. Until these early galaxy generations are observationally detected, CDGs are the best systems to test our ideas on primordial galaxy formation and evolution. Here we report on one of such CDGs, Tololo 1214-277, which presents features in its Lyman- α emission that had evaded theoretical interpretation so far. We show that these special features, a symmetric triple peaked emission line, are naturally explained by gas rotation. We find that the Lyman- α emission region in Tololo 1214-277 should have a rotational velocity of $V_r = 300 \text{ km s}^{-1}$ and a neutral Hydrogen column density of $\log N_{HI}/\text{atoms cm}^{-2} = 20.5$. Using archival observational information we find that the diameter for that region should be in the range of $110 \text{ pc} < D < 340 \text{ pc}$ and its total dynamical mass should be between $2.1 \times 10^9 M_\odot < M_D < 6.6 \times 10^9 M_\odot$. This dynamical mass is at least 16 ± 9 times larger than the neutral mass hydrogen. We argue that a possibility to explain the excess in dynamical mass is the presence of a super-massive black hole.

The first generation of galaxies trace our cosmic origins. Their characteristics are the first step on the evolution of galaxies such as the Milky Way. In the standard Big Bang cosmology the only elements that were created in the nucleosynthesis process were Hydrogen, Helium and Lithium. Heavier elements must have been created in stellar evolution process. Therefore the first generation of galaxies should be metal free and rich in Hydrogen. This kind of primordial galaxies have not been detected yet. However, dwarf star forming galaxies with a low metallicity content are close template to understand the early galaxy evolution process.

Almost fifty years ago [7] it was realized that young galaxies could be detected through a strong Lyman- α line emission. This theoretical prediction was only confirmed thirty year later, not on first generation galaxies but on distant relatively young systems. Currently, Lyman Alpha Emitting (LAE) galaxies are commonly targetted in surveys. The presence of the Ly- α emission line provides confirmation of the distance of a galaxy and also provides clues about the stellar population powering the emission.

Ly- α emission is not exclusive of distant galaxies. Any galaxy with low dust content and ongoing star formation has the right conditions to show this line. There are, for instance, local Universe surveys that target Ly- α emission in nearby dwarf star forming galaxies. [6]. A nearby LAE sample allow for the study of other indicators that might be more difficult to obtain for distant

galaxies such as morphology, dust attenuation, neutral hydrogen contents and ionization state.

However, the physical interpretation of Ly- α observations is not straightforward. This is due to the resonant nature of the Ly- α line. A Ly- α photon follows a diffusion-like process before escaping the galaxy or being absorbed by dust. The resulting line profile becomes sensitive to the dynamical, chemical and thermal conditions in the interstellar medium. Only in very few cases of high symmetry there are analytical solutions available. Interpretation of observations require state-of-the-art models use monte-carlo radiative transfer simulations.

Recent models computational models [2] have shown that rotation also imprints an effect on the Lyman-alpha morphology. The most important consequence of rotation is that spherical symmetry is broken. The line morphology now depends on the viewing angle respect to the rotation axis. For a line of sight perpendicular to the rotation axis the intensity and the line center and the line width increase with rotational velocity. When the rotational velocity is close to the half-line width of the static line, the line becomes single peaked as it is observed in Tololo 1214-277, a unique feature that other theoretical models find impossible to reproduce.

Tololo 1214-277 is a compact star forming dwarf galaxies (CDGs) that presents a strong Ly- α emission [10] with two features that had so far evaded a natural physical interpretation. First, it has symmetric profile without any systemic displacement from the expected line-center. Second, it shows a triple peak, while the most common observational feature corresponds either to a single or asymmetric double peak. We have attempted an interpretation of the Ly- α line in Tololo 1214-277 by including the effect of gas bulk rotation. We find that this model provides a satisfactory explanation of Tololo 1214-277's puzzling features.

Dots in Figure 1. represent the observational data for Tololo 1214-277 with the overplot from our best fit model from the full radiative transfer simulation. The parameters for the best fit are $v_{max} = 300 \text{ km s}^{-1}$, $\tau = 1 \times 10^7$, $T = 1.5 \times 10^4 \text{ K}$. The optical depth and temperature can be translated into a column density of N_{HI} . Our model is also able to constrain the angle between the rotation axis and the observational line-of-sight to $\theta = 65^\circ \pm$. To constrain the model uncertainties we perform a Markov Chain Monte Carlo calculation using an approximate analytical solution for the Ly- α spectrum in the bulk rotation case.

Radio surveys have put an upper limit to the neutral hydrogen mass in Tololo 1214-277 of $M < 2.65 \times 10^8 M_\odot$. Together with our velocity and column density results, under the assumption of spherical symmetry, we can constrain the total dynamical mass of the HI region to be in the range $2.1 \times 10^9 M_\odot < M_D < 6.6 \times 10^9 M_\odot$ and the diameter of HI region should have a diameter of $0.11 < D/\text{kpc} < 0.34$. This makes the dynamical mass at least 7 to 25 times larger than the HI mass.

We consider three possibilities to contribute to this high dynamical mass: stars, dark matter and a supermassive black hole. The stellar mass can be




Figure 1: Nice fit

constrained by measurements of the stellar continuum, this can only represent xxM_{\odot} of the total contribution. Considering a standard variations for the dark matter halo, it can only contribute to $xx M_{\odot}$ in the region of interest. A super massive black hole remains an open possibility. Recent observations of ultra-compact dwarf galaxies [9] have confirmed the presence of super-massive black holes containing 15% of the total objects mass, suggesting that there is a large population of undetected black holes.

A future observational test of this possibility in Tololo 1214-277 would require integral field unit measurements spatially resolving its spatial extent. Tololo 1214-277 spans a region of ≈ 4 arcminutes. An instrument such as the Multi Unit Spectroscopic Explorer with its nominal 0.2 arcseconds spatial sampling over a 1.0 arcminute field in wide-field mode could provide a detailed mapping of different ionization lines and infer a kinematic map.

Why are not more examples of triple peaked emission in observed Lyman-alpha emitting galaxies? The ubiquity of galaxy outflows overimposed to the rotation transforms the line into the common double or single peaked observations. Therefore, the absence of HI outflows in Tololo 1214-277 is another special feature that permitted a symmetric triple peaked Ly α line to emerge.

The implications of our finding for the study of Lyman-alpha emitters, including the first generation of galaxies are manifold. First of all, it provides support to including rotation as a feature to interpret observed spectra. Secondly, it provides a motivation to explore under what conditions rotation dominated dynamics are expected in comparison with outflow dominated dynamics. Finally, if the hypothesis of a supermassive black hole in Tololo 1214-277 proves to be consistent with future observational kinematic maps, it would provide a new information to test and probe theories on the coevolution of galaxies and black holes.

References

- [1] Klaus J. Fricke, Yuri I. Izotov, Polychronis Papaderos, Natalia G. Guseva, and Trinh X. Thuan. An imaging and spectroscopic study of the very metal-deficient blue compact dwarf galaxy Tol 1214-277. *The Astronomical Journal*, 121(1):169, 2001.
- [2] J. N. Garavito-Camargo, J. E. Forero-Romero, and M. Dijkstra. The Impact of Gas Bulk Rotation on the Ly α Line. *ApJ*, 795:120, November 2014.
- [3] D. G. Hummer and P. J. Storey. Recombination-line intensities for hydrogenic ions. I - Case B calculations for H I and He II. *MNRAS*, 224:801–820, February 1987.
- [4] Y. I. Izotov, P. Papaderos, N. G. Guseva, K. J. Fricke, and T. X. Thuan. Deep VLT spectroscopy of the blue compact dwarf galaxies Tol 1214-277 and Tol 65. *A&A*, 421:539–554, July 2004.

- [5] J. M. Mas-Hesse, D. Kunth, G. Tenorio-Tagle, C. Leitherer, R. J. Terlevich, and E. Terlevich. Ly α Emission in Starbursts: Implications for Galaxies at High Redshift. *ApJ*, 598:858–877, December 2003.
- [6] G. Östlin, M. Hayes, F. Duval, A. Sandberg, T. Rivera-Thorsen, T. Marquart, I. Orlitová, A. Adamo, J. Melinder, L. Guaita, H. Atek, J. M. Cannon, P. Gruyters, E. C. Herenz, D. Kunth, P. Laursen, J. M. Mas-Hesse, G. Micheva, H. Otí-Floranes, S. A. Pardy, M. M. Roth, D. Schaerer, and A. Verhamme. The Ly α Reference Sample. I. Survey Outline and First Results for Markarian 259. *ApJ*, 797:11, December 2014.
- [7] R. B. Partridge and P. J. E. Peebles. Are Young Galaxies Visible? *ApJ*, 147:868, March 1967.
- [8] S. A. Pustilnik and J.-M. Martin. H I study of extremely metal-deficient dwarf galaxies. I. The Nançay radio telescope observations of twenty-two objects. *A&A*, 464:859–869, March 2007.
- [9] A. C. Seth, R. van den Bosch, S. Mieske, H. Baumgardt, M. D. Brok, J. Strader, N. Neumayer, I. Chilingarian, M. Hilker, R. McDermid, L. Spitler, J. Brodie, M. J. Frank, and J. L. Walsh. A supermassive black hole in an ultra-compact dwarf galaxy. *Nature*, 513:398–400, September 2014.
- [10] T. X. Thuan and Y. I. Izotov. Nearby Young Dwarf Galaxies: Primordial Gas and Ly α Emission. *ApJ*, 489:623–635, November 1997.

$\alpha(2000)^a$	12h17min17.1s
$\delta(2000)^b$	-28d02m32s
l, b (deg)	294, 34
m_V	17.5
$v(\text{km s}^{-1})$	7795

Table 1: Observational characteristics of TOL1214-277 [10]

Tololo 1214-277 characteristics

Tololo 1214-277 receding velocity is $7785 \pm 50 \text{ km s}^{-1}$, which translates into a distance of 106.6 Mpc (Hubble constant $73 \text{ Mpc km}^{-1} \text{ s}^{-1}$) Its metallicity is $\sim Z_{\odot}/24$ [4] as derived from optical spectroscopy.

The observed flux for the Lyman alpha line is $\sim 8.1 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ [10] and a Equivalent Width of 70 Å and its $H\beta$ flux is $1.62 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ [4] which gives a $\text{Ly}\alpha/H\beta$ flux ratio of 4.9 ± 0.1 . Comparing this ratio with the theoretical expectation from case B recombination of 23.3 [3] one can estimate an escape fraction of 20% for $\text{Ly}\alpha$ radiation.

The optical emission comes from a region with approximate diameter ?? [1].

MCMC constraints

Physical Interpretation

Interpretation by [5].

There is an upper limit for the integrated flux of $< 0.10 \text{ Jy km s}^{-1}$, which translates into a upper limit for the HI mass of $M < 2.65 \times 10^8 M_{\odot}$ [8]. From the optical depth of 10^7 and the non-detection in the HI line, we have an upper limit for the size where the $\text{Ly}\alpha$ emission comes of $D < 0.34 \text{ kpc}$.

For an homogeneous sphere the HI optical depth from its can be written as $\tau = \sigma_0 n D / 2$, where $\sigma_0 = 5.898 \times 10^{-14} \text{ cm}^{-2}$ is the Lyman α optical depth at the line's center, n is the number density and D is the sphere's diameter. From this we can impose additional constraints on D from the typical values of the Hydrogen number density and our constrain on $\tau = 10^7$. Using a range of $1 < n / \text{atoms/cm}^{-3} < 10^{-3}$. This gives us a range of $0.11 < D / \text{kpc} < 100$. Together from the total HI mass we have thus that the HI region should have a diameter of $0.11 < D / \text{kpc} < 0.34$.

This can be rewritten in terms of the gas' temperature T and column density N_H as $\tau = 3.31 \times 10^{-14} (10^4 \text{ K} / T)^{1/2} (N_H / \text{atoms cm}^{-2})$.

This allows us to approximate the total hydrogen mass as

$$M_H = m_H N_H D^2 = 226 \times \tau \left(\frac{T}{10^4 \text{ K}} \right)^{1/2} \left(\frac{D}{\text{kpc}} \right)^2 M_{\odot} \quad (1)$$

On the other hand, we have an estimate for the dynamical mass from the galaxy size D and its rotational velocity V :

$$M_T = \frac{V^2 D}{G} = 2.16 \times 10^5 \left(\frac{V}{\text{km s}^{-1}} \right)^2 \left(\frac{D}{\text{kpc}} \right) M_\odot \quad (2)$$

From this limit and the rotational velocity of 300 km/s and the limit in the size D we have limits of for the dynamical mass of $2.1 \times 10^9 M_\odot < M_D < 6.6 \times 10^9 M_\odot$. which is at least 7 to 25 times larger than the HI mass.

Joint Effect of Rotation and Outflow

The almost complete absence of outflows is a central requirement for reproducing the lyman alpha line in Tololo 1214-277. In this section we present some results on the exploration of the presence of outflows join with the bulk rotation. We defer a detailed exploration of this effects for another publication.

Figure XX shows the results of four different models. a) pure rotation ($V_{\text{rot}} = 100 \text{ km s}^{-1}$), b) pure outflow ($V_{\text{out}} = 100 \text{ km s}^{-1}$), c) mixed outflow with rotation ($V_{\text{rot}} = 100 \text{ km s}^{-1}$)($V_{\text{rot}} = 25 \text{ km s}^{-1}$).