

Dark Matter’s influence on Evolution of MBHB in Dwarf Galaxies: A Case Study of Leo I dSph

Muhammad Junaid^{1,2*}, Fazeel Mahmood Khan^{1,2}

¹*Department of Space Science, Institute of Space Technology, Islamabad 44000, Pakistan*

²*Space and Astrophysics Research Lab (SARL), National Centre of GIS and Space Applications (NCGSA), Islamabad 44000, Pakistan*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

In this study, we investigate the dynamical evolution of a massive binary black hole (MBHB) in the Leo I dwarf spheroidal galaxy model and examine how dark matter along with stellar matter’s gravitational interactions influence its long-term behavior. Using high-resolution direct N-body simulations, we follow the orbital evolution of the binary within a realistic model of the Leo I stellar and dark matter distribution. We found that the binary separation decreases from an initial 300-parsec orbit to roughly 1 parsec over a period of about 2 Gyr, primarily driven by dynamical friction and stellar hardening. The orbital evolution then stalls at this scale, illustrating the well-known “final parsec problem”. During this phase, the binary also develops increasing orbital eccentricity and produces a modest redistribution of the inner mass profiles in some cases. We then further estimate the final stage of the system’s evolution using gravitational-wave emission models and find that the binary is unlikely to merge within a Hubble time. The prolonged dynamical friction phase appears to be related to the low stellar and dark matter densities in Leo I. These results suggest that massive binary black holes in dwarf spheroidal galaxies such as Leo I will not contribute to the gravitational-waves detectable from LISA even if dark matter is considered.

Key words: black hole physics – dwarf galaxies – gravitational waves – methods: numerical

1 INTRODUCTION

1.1 Leo I

Leo I is a dwarf spheroidal (dSph) galaxy and a well-studied member of the Local Group, gravitationally associated with the Milky Way subsystem. Early investigations of its stellar kinematics and metallicities established it as a chemically and dynamically distinct system [Koch et al. \(2007\)](#), while studies of its chemical abundances provided further constraints on its stellar populations and evolutionary history [Bosler et al. \(2007\)](#). Structural analyses have revealed complex, multi-age stellar populations [Okamoto et al. \(2011\)](#), and Leo I has been classified as an unquenched, very low-mass galaxy, suggesting an evolutionary path distinct from other dwarf galaxies [W. McQuinn et al. \(2015\)](#). Its dark matter halo profile is consistent with a central core, with a total mass comparable to other Milky Way dSph satellites [Koch et al. \(2007\)](#); [Mateo et al. \(1998\)](#). Broader studies have also placed Leo I within the context of dSph progenitor populations and the evolutionary puzzle of the Local Group [Grebel et al. \(2003\)](#), while its kinematics have been used to probe dark matter halo structure at the low-mass end of the galaxy luminosity function [Salucci et al. \(2012\)](#).

A particularly significant finding emerged from integrated-light measurements and previously published stellar velocity data, which revealed a consistent increase in velocity dispersion toward the center of Leo I, with an integrated-light value of 11.76 ± 0.66 km/s within $75''$ [Bustamante-Rosell et al. \(2021\)](#). Axisymmetric, orbit-based dynamical models accounting for the stellar mass-to-light ratio, dark matter halo structure, and tidal effects from the Milky Way at large radii consistently indicate a central black hole mass of $M_{\bullet} \sim 3.3 \pm 2 \times 10^6 M_{\odot}$ [Bustamante-Rosell et al. \(2021\)](#). The absence of a black hole is ruled out at greater than 95% confidence. This makes Leo I the first dSph galaxy in which a central black hole has been detected using spatially resolved kinematics, with a mass comparable to both the total stellar mass of the system and the Milky Way’s central black hole [Bustamante-Rosell et al. \(2021\)](#).

The detected black hole mass places Leo I well off the standard $M_{\bullet}-M_{\star}$ scaling relation by more than two orders of magnitude [Pacucci et al. \(2023\)](#). One proposed explanation is that Leo I’s stellar system was originally more massive and consistent with the scaling relation, but lost a substantial fraction of its stellar mass through extreme tidal stripping during close passages within the Milky Way’s virial radius [Pacucci et al. \(2023\)](#). Analytical models and N-body simulations estimate a stellar mass loss of 32%–78% depending on pericenter distance and velocity dispersion, with up

* E-mail: mjunaidtariq94@gmail.com

to 90% possible with additional passages [Bustamante-Rosell et al. \(2021\)](#); [Pacucci et al. \(2023\)](#). A tidal stream aligned along the line of sight toward Leo I is noted as potential evidence of this stripping event, though its detection in current Gaia data remains challenging [Pacucci et al. \(2023\)](#).

1.2 Dark Matter

Dark matter is a non-luminous component that interacts gravitationally but does not emit or absorb electromagnetic radiation. Within the Λ CDM framework, it constitutes approximately 25% of the total mass-energy content of the universe. Its existence was first postulated by Fritz Zwicky in 1933 to account for the anomalous orbital velocities of galaxies within the Coma cluster [Berera \(2011\)](#). Since then, a broad range of observational evidence has supported its presence, including galaxy rotation curves, gravitational lensing, and the large-scale structure of the universe [Abramowski et al. \(2015\)](#); [Esmaili et al. \(2012\)](#).

The nature of dark matter remains one of the central open questions in modern physics. Leading theoretical candidates include weakly interacting massive particles (WIMPs), axions, and sterile neutrinos [Abe et al. \(2020\)](#); [Pefianco \(2022\)](#). On cosmological scales, dark matter provides the gravitational scaffolding for structure formation and the cosmic web [Spergel & Steinhardt \(2000\)](#), while at galactic scales it governs rotation curves, mass distributions, and dynamical stability [Bertone et al. \(2005\)](#); [Kronawitter et al. \(2000\)](#); [Guo & Wu \(2009\)](#). Direct detection experiments, space-based observatories, and high-energy colliders continue to search for dark matter interactions with ordinary matter.

1.3 Evidence for Dark Matter

Three key observational pillars support the existence of dark matter:

- (i) **Galaxy Rotation Curves:** Observations of spiral galaxy rotation curves, systematically pursued from the late 1970s, reveal flat velocity profiles at large radii inconsistent with the distribution of luminous matter alone. This flatness implies the presence of a significant unseen mass component extending beyond the visible disk.
- (ii) **Gravitational Lensing:** The bending and distortion of light from background sources by foreground mass concentrations provides a direct probe of total matter content independent of luminosity. Both strong and weak lensing observations have confirmed substantial dark matter distributions in galaxy clusters and at cosmological scales [Bartelmann & Schneider \(2001\)](#); [Calore \(2020\)](#); [Pires et al. \(2010\)](#).
- (iii) **Cosmic Microwave Background (CMB):** Fluctuations in the CMB and their correlation with large-scale structure are consistent with the gravitational influence of dark matter on density perturbations in the early universe [Aslanyan et al. \(2016\)](#).

1.4 The Core-Cusp Problem

A persistent tension exists between Λ CDM predictions and observations of dwarf galaxy interiors. N-body simulations of structure formation predict that dark matter halos should follow a Navarro-Frenk-White (NFW) density profile, characterized by a central cusp where $\rho_{\text{DM}} \propto r^{-1}$. In contrast, observed rotation curves of gas-rich dwarf irregular galaxies favor constant-density cores in the inner halo, inconsistent with the predicted cusp.

Hydrodynamical simulations that incorporate baryonic processes offer a resolution through what is termed cusp-core transformation. As gas cools and condenses at galactic centers, stellar feedback drives repeated outflows that transfer energy and angular momentum to the dark matter, gradually flattening the cuspy NFW profile into a shallower core over multiple cycles of gas inflow and outflow [Read et al. \(2016\)](#). This gravitational heating mechanism, sometimes called dark matter heating, reproduces observed core profiles in low-mass galaxies within a Λ CDM framework.

1.5 Supermassive Black Hole Binaries

The closest known supermassive black hole (SMBH) binary was identified in the radio galaxy 0402+379. Multi-frequency Very Long Baseline Array (VLBA) observations identified two central compact flat-spectrum components [Maness et al. \(2004\)](#), subsequently confirmed as two SMBHs within a single galaxy by [Rodríguez et al. \(2006\)](#). Their projected separation is 7 pc, and the combined system mass is estimated at $1.5 \times 10^8 M_{\odot}$ [Maness et al. \(2004\)](#); [Rodríguez et al. \(2006\)](#). Galaxy mergers are widely regarded as the primary formation channel for SMBH binaries, making their study central to understanding galaxy evolution.

1.6 Evolution of SMBH Binaries

The dynamical evolution of SMBH pairs following a galaxy merger proceeds through three broad phases, as outlined by [Begelman et al. \(1980\)](#):

- (i) **Dynamical Friction:** The galactic cores migrate toward the merger center under dynamical friction from background stars, gas, and dark matter. The cores merge through violent relaxation, forming a new nucleus. A bound binary forms once the SMBH separation falls below the gravitational influence radius.
- (ii) **Binary Hardening:** The binary continues to shrink through dynamical friction and three-body gravitational slingshot interactions with stars on intersecting orbits. These stars carry away energy and angular momentum and are ejected at velocities comparable to the binary orbital velocity.
- (iii) **Gravitational Wave Inspiral:** At sufficiently small separations, gravitational wave emission dominates the energy and angular momentum loss, driving rapid coalescence of the two black holes.

1.7 Stellar Dynamical Hardening

Following the dynamical friction phase, the SMBH binary enters the hardening regime. For a binary with component masses $M_{\bullet 1}$ and $M_{\bullet 2}$, total mass $M_{\bullet} = M_{\bullet 1} + M_{\bullet 2}$, mass ratio $q = M_{\bullet 2}/M_{\bullet 1}$, and reduced mass $\mu = M_{\bullet 1} M_{\bullet 2}/M_{\bullet}$, the gravitational influence radius is defined as $r_h = GM_{\bullet}/\sigma^2$, where σ is the one-dimensional stellar velocity dispersion. Binaries are classified as hard or soft relative to the mean stellar kinetic energy per unit mass, following Heggie's law: hard binaries statistically harden further while soft binaries soften [Quinlan \(1996\)](#). The hardening rate is driven by three-body encounters and is characterized by the dimensionless hardening parameter H [Quinlan \(1996\)](#).

1.8 Gravitational Wave Emission

As the binary separation decreases sufficiently, gravitational wave emission takes over as the dominant energy loss mechanism. The

characteristic frequency of gravitational waves is inversely proportional to the black hole mass, placing SMBH mergers in the nanohertz to millihertz frequency band. This is below the sensitivity range of ground-based detectors and requires space-based observatories such as LISA or Pulsar Timing Arrays (PTAs). Black hole mergers in the mass range 10^3 – $10^7 M_\odot$ are expected to be detectable by LISA with a mass precision of approximately 5% (Bellovary et al. 2019). The orbital evolution under gravitational wave emission follows Peters' (1964) formulation Peters (1964), which describes the decay of both semi-major axis and eccentricity as functions of the binary parameters, with coalescence timescale T_{GW} dependent on the initial separation, eccentricity, and component masses.

The manuscript is structured as follows: Section 2 outlines the modelling and simulation framework and tools, Section 3 summarizes the data obtained, and Section 4 evaluates the implications of the work.

2 INITIAL MODEL SETUP

2.1 Galaxy Mass Distribution Functions

Galaxies exhibit a wide range of structures and dynamics. Understanding the distribution of mass within galaxies is crucial for unraveling their formation and evolution. One of the key aspects in modeling the mass distribution of galaxies is the use of spherical models. These models provide valuable insights into the gravitational potential, dynamics, and structural properties of galaxies. The study of nuclear star clusters, globular clusters, massive black holes, and dark matter halo within the context of spherical systems has significantly advanced our understanding of galaxy formation and evolution. Galaxy models for spherical systems are crucial for understanding the structure and dynamics of galaxies. Various models have been developed to describe the distribution of mass in galaxies, each with its unique characteristics and applications.

The Dehnen model is a significant spherical galaxy model that has been widely used to represent the mass distribution in galaxies. This model, introduced by Dehnen (1993), is precious for studying the dynamics of spherical stellar systems. The Dehnen model is characterized by its flexibility in describing the density profiles of galaxies, making it applicable to a wide range of galactic systems. It is also known as the η – *Tremaine* model. The density profile by Dehnen is:

$$\rho_{\text{dehnen}}(r) = \rho_0 \left(\frac{r}{r_s} \right)^{-\gamma} \left(1 + \frac{r}{r_s} \right)^{(\gamma-4)} \quad (1)$$

here r_s is the scale radius, and ρ_0 is the normalization density which is $(3 - \gamma)M/4\pi r_s^3$. The Dehnen cumulative mass profile is defined by:

$$M(r) = M \left[\frac{r}{r + r_s} \right]^{3-\gamma} \quad (2)$$

where M is the total mass of the component to be modeled and γ is the cusp exponent which should be $0 \leq \gamma < 3$. The models by Jaffe and Hernquist correspond to $\gamma = 2$ and 1, respectively (Dehnen 1993). These models have been utilized to study the gravitational potential and dynamics of spherical stellar systems, contributing to our understanding of galaxy structure and evolution.

J. Read presented the core NFW profile in which the inner dark matter profile that can transition from cusp to core is dependent on a function, this makes the profile closer to what we deduce from observations. Read et al. (2016)

Modified Leo-I model					
Component	M ($10^6 M_\odot$)	M (MU)	r_s (kpc)	N	γ
Black hole	3.3	0.22	1×10^{-5}	1	0
Stellar matter	4.2	0.28	0.1	7,17,948	0.6
Dark matter	7.5	0.5	0.2	1,282,051	0.6
Full model	15	1	-	2,000,000	-

Table 1. Parameters for Leo-I model: M is mass, r_s is scale radius, N is the number of particles and γ is the Dehnen slope.

2.2 AGAMA

AGAMA, which stands for Action-based Galaxy Modeling Architecture, is a powerful tool used to generate stable galaxy models using Schwarzschild modeling. This python package is designed to construct self-consistent, multi-component galaxy models in dynamic equilibrium by using the distribution function in space and solving for the virial theorem; $2K + U = 0$, where K is the kinetic energy and U is the gravitational potential energy of the system. This approach allows for the construction of detailed models that accurately represent the mass distribution, kinematics, and orbital properties of stars within galaxies. AGAMA is unique in its ability to handle a wide range of galaxy components, including dark matter, stellar disks, bulges, and halos, allowing for the creation of complex and realistic galaxy models. The code uses a flexible and efficient algorithm to generate stable models that closely match observational data, making it a valuable tool for studying the formation and evolution of galaxies. It takes different parameters to construct a galaxy model: mass of the component or normalized density, axis ratios (p and q), density function, scale radius, slopes (α, β, γ) or the Sersic index (n), and cut-off radius with cut-off strength Vasiliev (2018).

2.3 Model Generation

To model the galaxy's overall mass distribution, we used Dehnen's distribution function and used AGAMA. Our model successfully reproduced the mass profiles of Bustamnte's data. We fitted both its slope parameter and the masses of the NFW model on the Dehnen model for dark matter in Leo-I. We then limit the profile to approximately 300 parsecs of the galaxy's center, consistent with the results of the previous literature. Beyond 300 parsecs, the cumulative mass profile flattened out while the density declined sharply as we intended. We constrained our model to only 300 parsecs to study the high-resolution dynamics of Leo-I with respect to binary black holes.

We chose 2 million particles and gave each stellar and dark matter particle equal mass. The mass of a star and dark matter particle is 3.89×10^{-7} in model units and $5.835 M_\odot$ in physical units. As the dark matter's mass is greater than our model's stellar mass, the number of dark matter particles will also be higher than the number of stars.

This approach allowed us to generate a modified model for the central region of the Leo-I dSph galaxy in equilibrium using AGAMA. The model's density and cumulative mass profile aligned with analytical analysis, testing whether its properties could be reproduced through similar assumptions about its black holes, dark matter, and stellar components as shown in (1) for the whole model and its components.

Since Leo-I is a dwarf spheroidal, the axis ratios of both of our components (stellar and dark matter) were chosen and shown in




Figure 1. Density profiles of the Modified Leo I model components over the radial range (r) from 0.1 pc to 1000 pc. The black hole is modeled with a Plummer profile, while the stellar and dark matter components follow Dehnen profiles. (see Table 1 for model parameters). The full model (dash-dot line) is the sum of all three components. Dotted vertical lines indicate the scale radii of the stellar and dark matter distributions.

Component	$p = b/a$	$q = c/a$
Stellar matter	0.9	0.8
Dark matter	0.9	0.8

Table 2. Axis ratios for modified Leo-I model

Quantity	Model Units	Physical Units
Mass	1 [M]	$1.5 \times 10^7 M_\odot$
Length	1 [L]	1 kpc
Velocity	1 [V]	7.97 km/s
Time	1 [T]	121.7372 Myrs

Table 3. Model units and their equivalence in physical units.

table 2. We also generated a visual representation of our modified Leo-I mass model using the Glnemo2 N-body visualization tool. Glnemo2 renders the positions of each particle.

2.4 Units

We adopted the model units in such a way that $M_{\text{gal}} = G = r_o = 1$. The model units can be scaled to physical units by changing the galaxy's mass M_{gal} and scale radius r_o to physical units, when combined with the relations below we can convert time and velocity from model to physical units (Gualandris & Merritt (2008)).

$$[T] = \left(\frac{GM_{\text{gal}}}{r_o^3} \right)^{-1/2} = 1.5 \text{ Myr} \left(\frac{M_{\text{gal}}}{10^{11} M_\odot} \right)^{-1/2} \left(\frac{r_o}{\text{kpc}} \right)^{3/2} \quad (3)$$

$$[V] = \left(\frac{GM_{\text{gal}}}{r_o} \right)^{1/2} = 655 \text{ km s}^{-1} \left(\frac{M_{\text{gal}}}{10^{11} M_\odot} \right)^{1/2} \left(\frac{r_o}{\text{kpc}} \right)^{-1/2} \quad (4)$$

Our model's data is in model units [MU] and their physical equivalence is given in the table below at table (3):

2.5 Secondary Black Hole

We inserted a secondary intermediate-mass black hole (IMBH) by modifying the second particle's mass, position, and velocity in our model. We chose the mass of the secondary black hole to be 100 times and 40 times less than that of our primary SMBH; hence, the mass ratios are $q_1 = 0.01$ and $q_2 = 0.025$, corresponding to secondary black hole masses of $3.3 \times 10^4 M_\odot$ and $8.25 \times 10^4 M_\odot$, respectively. We introduced the black hole at a distance of $r_{\text{ini}} = 300 \text{ pc}$. We chose the secondary black hole's velocity as 50% of the circular velocity at that radius for a particle this massive. The position was added in the x direction and velocity in the y component. This was deliberately done to make the initial orbit elliptical and in xy-plane.

$$v_{\text{cir}} = \sqrt{\frac{GM_{\text{enc}}(r_{\text{ini}})}{r_{\text{ini}}}} \quad (5)$$

2.6 N-body Code

N-body simulations are fundamental tools for studying the gravitational interactions and dynamics of astrophysical systems. These simulations involve the numerical integration of the equations of motion for a large number of particles, typically representing stars, dark matter, or other celestial objects, under the influence of gravitational forces. To perform these simulations, various N-body codes and integration methods have been developed, each with unique features and applications. In astrophysics, a system with N-particles interacting with each other gravitationally has a force F_i acting on each particle with mass m_i :

$$\mathbf{F}_i = -m_i \sum_{j=1, j \neq i}^N \frac{m_j (\mathbf{r}_i - \mathbf{r}_j)}{(|\mathbf{r}_i - \mathbf{r}_j|^2 + \epsilon_{ij}^2)^{3/2}} \quad (6)$$

here, ϵ is called the softening parameter. It is used to stop close body encounters and from further binaries to be formed. The value of softening parameter between two particles is then calculated to be:

$$\epsilon_{ij} = \sqrt{\frac{\epsilon_i^2 + \epsilon_j^2}{2}} \quad (7)$$

Gravitational forces on each particle from all the particles in the system are computed by N-body code, and for a number of time steps, the computational difficulty scales as N^2 .

2.6.1 Hermite Integrator Scheme

The Hermite integrator is a numerical method commonly used in N-body simulations to accurately solve the equations of motion for a system of particles under the influence of gravitational forces. This integration method is particularly well-suited for simulating the dynamics of astrophysical systems, such as galaxies and star clusters, where the gravitational interactions between a large number of particles need to be accurately calculated over extended periods of time. The Hermite integrator is known for its high accuracy and efficiency in long-term gravitational interactions. It achieves this by using higher-order terms in the Taylor expansion of the equations of motion, allowing for more precise predictions of particle trajectories and gravitational forces. Additionally, the Hermite integrator can adapt its time step to account for variations in the

Softening length (ϵ) in kpc	
ϵ_{SMBH}	0.0
$\epsilon_{\star}$	1×10^{-4}
ϵ_{DM}	1×10^{-4}

Table 4. Softening length for different components particles of modified Leo-I model in kpc.

gravitational forces, ensuring stable and accurate simulations over extended timescales. The Hermite fourth-order integrator is called 4^{th} -order because the predictor term considers the third-order polynomial, and then after obtaining the acceleration, adds a 4^{th} -order corrector term. Harfst et al. (2007).

2.6.2 φ -GPU

The parallel processing capabilities of GPUs allow for the simultaneous execution of a large number of computational tasks, making them well-suited for accelerating N-body simulations, which involve the gravitational interactions of a vast number of particles.

φ -GPU is a computational framework designed for performing N-body simulations, particularly focusing on the dynamics and interactions of particles in astrophysical systems. It is based on φ -GRAPE6a (GRAvity PipelinE) code which can do the same function but on special boards. This framework leverages the computational power of Graphics Processing Units (GPUs) to speed up the simulations, enabling the efficient and high-performance modeling of complex gravitational systems. The φ -GPU framework harnesses the parallel processing power of GPUs to efficiently calculate the gravitational forces and trajectories of particles in N-body simulations. By distributing the computational workload across numerous processing cores within the GPU, φ -GPU significantly speeds up the simulation time, enabling researchers to study the dynamics and evolution of astrophysical systems with high precision and computational efficiency. The φ -GPU code builds upon a CPU serial N-body code, written in C++. It fully parallelizes the code using MPI. Additionally, it accesses GPUs directly through the NVIDIA CUDA library for native GPU support. Berczik et al. (2011)

Before running long simulations, we performed a stability analysis of our Leo-I model using the φ -GPU N-body solver. As φ -GPU does not incorporate regularization methods for close encounters, gravitational softening was required for all particle interactions. A softening length of 1×10^{-4} in our model units was adopted. This value was chosen to be sufficiently small to maximize the collisional nature of the dense stellar and dark matter components while resolving interactions down to tens of parsecs. The softening for the SMBH particle is equal to zero as shown in table (4)

The model was evolved in the first run of the simulations for 1.5 model time units, equivalent to 180 million years. Density profiles and cumulative mass profiles were calculated at several timestamps throughout the simulation. We find that the model remains stable over this period, with no significant deviations from the initial conditions.

3 RESULTS

The following sections present the time-resolved properties of the binary system and its impact on the host galaxy. We investigate

the orbital evolution of a massive black hole binary (MBHB) in the central 300 pc of a Leo I for mass ratio $q = 1 : 100$ as well as $q = 1 : 40$.

3.1 Binary Evolution Overview

Figure 2 presents a comprehensive view of the MBHB orbital evolution over 2.5 Gyr of simulation time for Leo I model. The upper-left panel shows the relative separation, for $q = 0.01$, relative separation gradually shrinks from ~ 300 pc to sub-parsec scales. The binary reaches ~ 10 pc after ~ 1.5 Gyr, and 5 pc near the 2 Gyr mark. But for $q = 0.025$ model, relative separation of BBH reaches 10 pc within 1 Gyr but after this the binary stalls. At no point does the separation drop below 0.1 pc, suggesting long-term stalling instead of moving towards gravitational-wave (GW) coalescence. The upper-right panel shows the semi-major axis (a), which steadily declines following binary formation ($T_b \sim 0.5$ Gyr). Before this time the semi-major axis is estimated using the periapsis and apoapsis values for binary and later the solid line represents semi-major from the simulation's results. Similar to the relative separation, semi-major axis of the $q = 0.01$ system drops gradually to the final parsec meanwhile $q = 0.025$ system reaches that mark quicker by almost 1 Gyr. The flattening trend beyond 1.5 Gyr suggests inefficient hardening. After that the semi-major axis in both simulations stalls from further dropping. In gravitational emission regime of BBH evolution, the average energy emitted in terms of GWs is given by Peters (1964):

$$\frac{dE_{\text{rad}}}{dt} = \frac{32}{5} \frac{G^4}{c^5} \frac{M_1^2 M_2^2 (M_1 + M_2)}{a^5} F(e) \quad (8)$$

where the factor

$$F(e) = (1 - e^2)^{-7/2} \left(1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right) \quad (9)$$

where $F(e)$ is a function dependent on the binary's eccentricity. So, the change in semi-major axis and eccentricity due to GW will be:

$$\left. \frac{da}{dt} \right|_{\text{GW}} = -\frac{64}{5} \frac{G^3}{c^5} \frac{M_1 M_2 (M_1 + M_2)}{a^3} F(e) \quad (10)$$

$$\left. \frac{de}{dt} \right|_{\text{GW}} = -\frac{304}{15} \frac{G^3}{c^5} \frac{M_1 M_2 (M_1 + M_2)}{a^4} e(1 - e^2)^{-5/2} \left(1 + \frac{121}{304} e^2 \right) \quad (11)$$

So, the estimation of semi-major axis using the loss of orbital energy via GW suggests that the binaries will not merge by reaching the Schwarzschild radius of the primary black hole ($r_s = 3.16 \times 10^{-7}$ pc) within the Hubble time as the change is very small over time. The bottom-left panel shows inverse semi-major axis ($1/a$), a useful parameter for analysis of binary hardening. Its decreasing slope implies progressive stalling. The hardening rate is given by:

$$s_{\text{NB}} = \left(\frac{d}{dt} \left(\frac{1}{a} \right) \right)_{\text{NB}} = -\frac{1}{a^2} \left(\frac{da}{dt} \right)_{\text{NB}} \quad (12)$$

$$\left(\frac{da}{dt} \right)_{\text{NB}} = -s_{\text{NB}} \cdot a^2, s_{\text{NB}} \approx \text{const} \quad (13)$$

$$\left(\frac{da}{dt} \right)_{\text{total}} = -(s_{\text{NB}} \cdot a^2) + \left(\frac{da}{dt} \right)_{\text{GW}} \quad (14)$$

A hardening rate $s_{\text{NB}} \approx 0.31 \text{ kpc}^{-1} \text{ Myr}^{-1}$ for $q = 0.01$ and




Figure 2. Orbital evolution of the massive black hole binary. Top-left: relative separation (ΔR). Top-right: semi-major axis (a). Bottom-left: inverse semi-major axis ($1/a$), inside plot zooming to the region showing the slopes used to calculate hardening rates. Bottom-right: eccentricity (e). Time is measured from $t = 0$ when the secondary black hole reaches apoapsis at 300 pc. The plus markers denote values derived from apoapsis and periapsis estimates, while the corresponding solid lines represent values directly extracted from the simulation. Gravitational wave (GW) evolution is overplotted for initial eccentricities of $e_0 = 0.75$ (bold solid line) and $e_0 = 0.99$ (dashed line). Green and gold curves correspond to the $q = 0.01$ model, while blue and copper indicate the $q = 0.025$ model. The vertical grey dashed line marks the age of the Universe (13.8 Gyr), and the horizontal black line in the upper panels indicates the Schwarzschild radius of the primary black hole.

$s_{\text{NB}} \approx 0.076 \text{ kpc}^{-1} \text{ Myr}^{-1}$ for $q = 0.025$ is inferred by the results of the simulation by calculating the slope of inverse semi-major axis around 2 to 2.5 Gyr. These extremely low values of hardening rates are then used to calculate semi-major axis and eccentricity till 14 Gyrs. We estimated semi-major axis and eccentricities from $a_0 \approx 2.5 \text{ pc}$ and $e_0 = 0.75$ for both models and as well as at extreme eccentricity, $e \approx 0.99$. The bottom-right panel displays the orbital eccentricity, estimated using apoapsis/periapsis extraction till the hard binary forms. Eccentricity grows gradually to $e \sim 0.85$ for $q = 0.01$ and $e \sim 0.95$ for $q = 0.025$ model by the end of the simulation. The greater value enhances GW energy loss but remains insufficient for coalescence within a Hubble time, given the stalled separation.

3.2 Axis ratio evolution

The initial shape of the galaxy model is nearly spherical with axis ratios $b/a = 0.9$, $c/a = 0.8$. Over the course of the simulation, the

inner galaxy evolves toward a more spherical shape, with c/a increasing to ~ 0.85 . This flattening trend, shown in Fig. 3, is likely induced by angular momentum redistribution from BBH–star interactions. A more spherical core may reduce the population of centrophilic orbits, suppressing further loss cone refilling.

3.3 Core Evolution

The MBHB’s hardening ejects stars via three-body encounters, depleting both stellar and dark matter densities in the central $\sim 50 \text{ pc}$. Figure 4 shows cumulative mass and density profiles at multiple timestamps. Over the 2 Gyr evolution, the core experiences a tenfold drop in density. The resulting mass deficit halts further hardening once the binary reaches $\sim 0.4 \text{ pc}$, leading to long-term stalling. Figure 5 shows how gamma (inner slope) evolved from shallow cusp and initial value of 0.6 throughout the simulation period. For less massive MBHB model, it remains almost stable and ends at closer




Figure 3. Time evolution of axis ratios in the $q = 0.01$ model (left) and $q = 0.025$ model (right): b/a as "+" and c/a as "x" and inner region (right). The system becomes more spherical over time, potentially limiting loss cone refilling.

to 0.8, but for heavier MBHB, γ first rises and peaks at almost 1 Gyr and then continues to drop and finally ends at 0.5 for dark matter and 0 or stellar matter, displaying core formation.

4 CONCLUSION AND DISCUSSION

4.1 Conclusion

We investigated the dynamical evolution of a massive black hole binary (MBHB) at the centre of a dwarf spheroidal galaxy, using Leo I as a well-constrained observational template. A self-consistent, multi-component N-body model of Leo I was constructed with AGAMA Vasiliev (2018), incorporating the observed stellar and dark matter distributions together with the central black hole inferred from kinematic data Bustamante-Rosell et al. (2021). A stability analysis confirmed that the model preserves its initial density and mass profiles over 180 Myr, validating it as a reliable initial condition for binary evolution experiments. A secondary IMBH was then introduced at 300 pc with mass ratios $q_1 = 0.01$ and $q_2 = 0.025$ relative to the primary, and the subsequent evolution was followed for ~ 2.5 Gyr using the direct N-body code φ -GPU Berczik et al. (2011).

Our principal findings are as follows.

- (i) **Binary formation and dynamical friction.** Both models form a gravitationally bound hard binary within ~ 0.5 –1 Gyr through dynamical friction against the stellar and dark matter background. The timescale is substantially longer than in more massive, denser systems, a direct consequence of Leo I's comparatively low stellar and dark matter densities, which are themselves a legacy of repeated tidal stripping by the Milky Way Pacucci et al. (2023).
- (ii) **Hardening and the final parsec problem.** Following binary formation, three-body stellar encounters drive the binary separation from ~ 300 pc to sub-parsec scales. The $q = 0.025$ model reaches the final parsec somewhat faster than the $q = 0.01$ model, but both ultimately stall at separations of order ~ 0.4 –1 pc. The inferred hardening rates are $s_{\text{NB}} \approx 0.31 \text{ kpc}^{-1} \text{ Myr}^{-1}$ for $q = 0.01$ and $s_{\text{NB}} \approx 0.076 \text{ kpc}^{-1} \text{ Myr}^{-1}$ for $q = 0.025$, among the lowest reported in binary evolution studies, reflecting the depleted loss cone in this low-density environment.

- (iii) **Gravitational-wave evolution.** Extrapolating beyond the simulated period using Peters' equations Peters (1964), neither model merges within a Hubble time. Even for the extreme case of $e = 0.99$, the binary only contracts to ~ 0.4 pc over 13.8 Gyr, falling far short of the Schwarzschild radius of the primary ($r_s = 3.16 \times 10^{-7}$ pc).
- (iv) **Eccentricity growth.** Orbital eccentricity grows progressively during the hardening phase, reaching $e \sim 0.85$ for $q = 0.01$ and $e \sim 0.95$ for $q = 0.025$. Although higher eccentricity enhances gravitational-wave energy loss, this enhancement is insufficient to overcome the stalled separation in the absence of an efficient loss-cone replenishment mechanism.
- (v) **Inner morphological evolution.** Three-body ejections from the binary heat and scatter the surrounding stellar and dark matter distributions, driving the inner morphology toward a more spherical configuration (c/a increases from ~ 0.8 to ~ 0.85) and reducing the population of centrophilic orbits available for loss-cone refilling. Simultaneously, the central density drops by roughly an order of magnitude within 50 pc, and the inner power-law slope γ shows model-dependent behaviour: it remains near 0.8 for the less massive secondary but first rises then falls below 0.5 for the more massive secondary, indicating the onset of core formation.

Taken together, these results demonstrate that while a SMBH–IMBH binary can assemble dynamically in a Leo I-type dwarf spheroidal, the final parsec problem persists robustly. The low-density, tidally stripped environment of Leo I is insufficient to drive the binary to coalescence through stellar dynamics alone, even when dark matter is included. Additional physical processes – such as gas inflows, residual triaxiality, or the presence of a nuclear star cluster – would be needed to overcome this barrier.

4.2 Discussion

The final parsec problem has long been recognised as a potential bottleneck in the merger history of massive black holes Sesana & Khan (2015), and our results provide a concrete illustration of how acutely it manifests in the low-density environment of a dwarf spheroidal. In more massive, gas-rich systems, AGN feedback and nuclear gas flows can replenish the loss cone on relatively short timescales. In Leo I, however, the stellar mass is comparatively low




Figure 4. Evolution of mass and density profiles over 2.4 Gyrs. Top panels for q_1 and bottom panels for q_2 model.

and largely tidally depleted [Pacucci et al. \(2023\)](#), and no significant gas reservoir is expected, leaving three-body stellar scattering as the dominant hardening channel – one that ultimately proves insufficient.

The question of whether the Leo I black hole is a descendant of a heavy seed formation event has been raised by recent theoretical work. [Scoggins et al. \(2025\)](#) argue that a black hole of $\sim 3 \times 10^6 M_\odot$ residing in a system with Leo I’s properties is broadly consistent with the expected descendants of heavy black hole seeds, motivating the interpretation of Leo I as a surviving relic of an early seed formation site. If this picture is correct, the binary studied here can be thought

of as the dynamical encounter between a heavy seed survivor and an infalling IMBH. We note, however, that the existence and mass of the Leo I black hole itself remains debated; some studies argue that the observed kinematic signal may be reproduced by an unusually high dark matter concentration in the nucleus, without requiring a black hole at all [Scoggins et al. \(2025\)](#). The conclusions drawn here are therefore conditional on the black hole being real.

A related astrophysical channel involves Leo I itself being accreted by the Milky Way. Dwarf galaxies like Leo I are expected to lose orbital energy to dynamical friction and eventually sink toward the centres of their more massive hosts. Should this occur, the Leo I




Figure 5. Mass and density evolution in the inner region. Profiles show central depletion due to BBH interactions, limiting further evolution.

black hole could be delivered to the Galactic nucleus and pair with the Milky Way's central black hole, Sgr A*, forming what is termed a heavy IMRI – an intermediate-mass ratio inspiral between a super-massive and an intermediate-mass black hole [Askar et al. \(2024\)](#). Dense stellar clusters carrying IMBHs can undergo a similar infall process, reaching galactic nuclei on timescales that depend on their birth radii and host properties [Askar et al. \(2024\)](#). Within Leo I itself, however, our simulations suggest that the SMBH–IMBH binary formed internally will not reach the gravitational-wave band within a Hubble time. This places Leo I-type dSphs in a different category from dwarf galaxies with nuclear star clusters, where higher central densities can sustain loss-cone refilling and drive binaries toward coalescence on more favourable timescales ([Khan & Holley-Bockelmann 2021](#)).

More broadly, these results highlight the sensitivity of MBHB evolution to the host galaxy environment. Dwarf spheroidals occupy the low end of the galaxy mass function [Mezcua \(2017\)](#), and their black hole populations remain poorly characterised observationally. Understanding whether and how binaries in such systems can merge is important for constructing realistic LISA source populations and for interpreting the nanohertz gravitational wave background detected by pulsar timing arrays [Amaro-Seoane et al. \(2017\)](#). Our work suggests that isolated, gas-poor dwarf spheroidals are unlikely to be significant contributors to either signal, but that the same black holes may become relevant IMRI sources if and when their host galaxies are absorbed into more massive systems.

ACKNOWLEDGMENTS

The authors gratefully acknowledge Vanderbilt University's Advanced Computing Center for Research and Education (ACCRE) and the National Center of GIS and Space Application (NCGSA)'s Space and Astrophysics Research Lab (SARL) and Modeling and

Simulation Lab for providing the computational resources used for this research.

DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Abe Y., Toma T., Tsumura K., 2020, *Journal of High Energy Physics*
- Abramowski A., et al., 2015, *Physical Review Letters*
- Amaro-Seoane P., et al., 2017, arXiv e-prints, p. [arXiv:1702.00786](#)
- Askar A., Baldassare V. F., Mezcua M., 2024, Intermediate-Mass Black Holes in Star Clusters and Dwarf Galaxies ([arXiv:2311.12118](#)), <https://arxiv.org/abs/2311.12118>
- Aslanyan G., Price L. C., Adams J., Bringmann T., Clark H. A., Easter R., Lewis G. F., Scott P., 2016, *Physical Review Letters*
- Bartelmann M., Schneider P., 2001, *Physics Reports*
- Begelman M. C., Blandford R. D., Rees M. J., 1980, *Nature*, 287, 307
- Bellovary J., et al., 2019, Where are the Intermediate Mass Black Holes?, [doi:10.48550/ARXIV.1903.08144](#), <https://arxiv.org/abs/1903.08144>
- Berczik P., et al., 2011, in International conference on High Performance Computing. pp 8–18
- Berera A., 2011, *Pramana*
- Bertone G., Hooper D., Silk J., 2005, *Physics Reports*
- Bosler T., Smecker-Hane T., Stetson P. B., 2007, *Monthly Notices of the Royal Astronomical Society*
- Bustamante-Rosell M. J., Noyola E., Gebhardt K., Fabricius M. H., Mazzalay X., Thomas J., Zeimann G., 2021, *The Astrophysical Journal*, 921, 107
- Calore F., 2020, *Physics*
- Dehnen W., 1993, *Monthly Notices of the Royal Astronomical Society*, 265, 250

- Esmaili A., Ibarra A., G. Peres O. L., 2012, [Journal of Cosmology and Astroparticle Physics](#)
- Grebel E. K., Gallagher J. S., Harbeck D., 2003, [The Astronomical Journal](#)
- Gualandris A., Merritt D., 2008, [The Astrophysical Journal](#), 678, 780
- Guo W., Wu Y., 2009, [Physical Review D](#)
- Harfst S., Gualandris A., Merritt D., Spurzem R., Zwart S. P., Berczik P., 2007, [New Astronomy](#), 12, 357
- Khan F. M., Holley-Bockelmann K., 2021, [MNRAS](#), 508, 1174
- Koch A., Wilkinson M. I., Kleyna J., Gilmore G., Grebel E. K., Mackey A. D., Evans N. W., G. Wyse R. F., 2007, [The Astrophysical Journal](#)
- Kronawitter A., Saglia R., Gerhard O., Bender R., 2000, [Astronomy and Astrophysics Supplement Series](#)
- Maness H., Taylor G., Zavala R., Peck A., Pollack L., 2004, [The Astrophysical Journal](#), 602, 123
- Mateo M., Olszewski E. W., Vogt S. S., Keane M., 1998, [The Astronomical Journal](#)
- Mezcua M., 2017, [International Journal of Modern Physics D](#), 26, 1730021
- Okamoto S., Arimoto N., Yamada Y., Onodera M., 2011, [The Astrophysical Journal](#)
- Pacucci F., Ni Y., Loeb A., 2023, [The Astrophysical Journal Letters](#), 956, L37
- Pefianco J., 2022, [Journal of Student Research](#)
- Peters P. C., 1964, [Physical Review](#), 136, B1224
- Pires S., Starck J.-L., Réfrégier A., 2010, [Ieee Signal Processing Magazine](#)
- Quinlan G. D., 1996, [New Astronomy](#), 1, 35
- Read J., Agertz O., Collins M., 2016, [Monthly Notices of the Royal Astronomical Society](#), 459, 2573
- Rodríguez C., Taylor G. B., Zavala R., Peck A., Pollack L., Romani R., 2006, [The Astrophysical Journal](#), 646, 49
- Salucci P., Wilkinson M. I., Walker M. G., Gilmore G., Grebel E. K., Koch A., Martins C. F., G. Wyse R. F., 2012, [Monthly Notices of the Royal Astronomical Society](#)
- Scoggins M. T., Haiman Z., Pacucci F., 2025, Heavy black hole seed survivors in dwarf galaxies: a case study of Leo I ([arXiv:2511.04736](#)), <https://arxiv.org/abs/2511.04736>
- Sesana A., Khan F. M., 2015, [MNRAS](#), 454, L66
- Spergel D. N., Steinhardt P. J., 2000, [Physical Review Letters](#)
- Vasiliev E., 2018, [Monthly Notices of the Royal Astronomical Society](#), 482, 1525–1544
- W. McQuinn K. B., et al., 2015, [The Astrophysical Journal](#)

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.