

Infrared Spectral Gap in a Gluonic Dark Sector as the Origin of the Galactic Acceleration Scale

Gilles Cohen-Tannoudji^{1*}, Jean-Pierre Gazeau^{2,3†}, Hamed Pejhan^{4‡}, and Jean-Pierre Treuil^{5§}

¹*Laboratoire de Recherche sur les Sciences de la Matière (LARSIM),
CEA, Université Paris-Saclay, F-91190 Saint-Aubin, France*

²*Université Paris Cité, CNRS, Astroparticule et Cosmologie, F-75013 Paris, France*

³*Faculty of Mathematics, University of Białystok, 15-245 Białystok, Poland*

⁴*Institute of Mathematics and Informatics, Bulgarian Academy of Sciences,
Acad. G. Bonchev Str. Bl. 8, 1113, Sofia, Bulgaria and*

⁵*Institut de Recherche pour le Développement (IRD), France*

(Dated: April 15, 2026)

The radial acceleration relation reveals a nearly universal acceleration scale of order 10^{-10} ms^{-2} in galactic dynamics, whose origin remains unexplained within conventional cold dark matter scenarios. We propose that this scale arises from an intrinsic infrared spectral property of the dark sector. Specifically, we hypothesize that a long-lived, color-neutral gluonic vacuum component survives the post-inflationary expansion era and, at large distances, develops a spectrally rigid lowest-weight structure. The microscopic seed for this infrared organization is provided by the QCD trace anomaly, which breaks classical scale invariance and, through dimensional transmutation, generates an intrinsic infrared scale in the gluonic sector. Requiring Lorentz covariance together with a positive-energy lowest-weight unitary realization then selects the Anti de Sitter algebra $\mathfrak{so}(2,3)$ as the simplest symmetry admitting a discrete tower of states with a representation-theoretically protected gap. The associated gap introduces a finite correlation length r_c that controls the large-scale coherence of the dark sector. A self-gravitating condensate dominated by the lowest-weight mode generates a characteristic acceleration $g_* = GM_h/r_c^2$, naturally of the same order as the observed galactic acceleration scale, within standard Newtonian gravity. In this framework, the galactic acceleration scale appears as the gravitational imprint of a trace-anomaly-seeded infrared spectral gap in a coherent gluonic dark sector, rather than as a consequence of modified gravity or of galaxy-by-galaxy formation histories.

I. INTRODUCTION

We investigate the hypothesis that the galactic radial acceleration scale arises from an intrinsic infrared spectral gap in a coherent gluonic dark sector. Specifically, we propose that a long-lived, colour-neutral gluonic vacuum component — rooted in the non-perturbative structure of the QCD trace anomaly [1] — survives the post-inflationary expansion era and organizes at large distances into a lowest-weight unitary irreducible representation of $\text{SO}(2,3)$. The associated discrete spectrum introduces a dynamically protected infrared correlation length, whose macroscopic gravitational imprint generates the observed galactic acceleration scale without modifying gravitational dynamics.

In this framework, the radial acceleration relation (RAR) is interpreted not as an emergent by-product of baryon-halo assembly histories, but as the large-scale manifestation of a representation-theoretically protected gap in the dark sector. The remainder of this Introduction provides the empirical motivation and theoretical justification for this hypothesis and outlines the theoret-

ical framework used throughout the paper.

The present analysis extends earlier studies [2–5] and further develops the idea that the infrared gluonic sector may organize into a lowest-weight spectral structure whose associated correlation length governs the macroscopic behaviour of the dark component.

A. Empirical Universality of the RAR

The RAR exhibits a remarkably tight correlation between the observed gravitational acceleration in rotationally supported galaxies and that inferred from the baryonic mass distribution. The relation is characterized by a single acceleration scale of order 10^{-10} ms^{-2} and a very small intrinsic scatter across systems spanning wide ranges of mass, morphology, and evolutionary history [6–8].

Although Lambda cold dark matter (ΛCDM) simulations that include baryonic feedback are able to reproduce similar trends [9–11], within the standard framework, the RAR is not determined by fundamental parameters of the dark sector. Rather, it arises statistically from the correlated assembly of baryons and halos, and its apparent regularity reflects the outcome of complex formation histories [11–13].

If the observed near-universality of the acceleration scale — despite recent claims questioning its statistical significance [14] — is confirmed by future data, it be-

*gilles.cohentannoudji@gmail.com

†gazeau@apc.in2p3.fr and j.gazeau@uwb.edu.pl

‡pejhan@math.bas.bg

§treuil.jean-pierre@orange.fr

comes natural to ask whether such a small system-to-system variation can be entirely attributed to baryon-halo coupling. A parameter exhibiting this level of stability across galaxies instead suggests the presence of an intrinsic large-scale property of the dark sector itself. From this perspective, the RAR may be viewed as the macroscopic manifestation of a fundamental infrared scale governing the dynamics of DM.

B. Infrared Rigidity and the Necessity of a Lowest-Weight Structure

The RAR acceleration scale emerges at kiloparsec distances and exhibits striking insensitivity to microscopic particle parameters and detailed halo assembly histories. Its origin must therefore be sought in the large-scale collective dynamics of the dark sector rather than in ultraviolet particle physics. Any successful explanation must account not only for the presence of a characteristic scale but also for its universality and small intrinsic scatter across systems with diverse formation histories.

If the dark sector were infrared scale-free, it would contain no dynamically selected macroscopic length. Halo structure would then be shaped primarily by environmental factors — non-linear collapse, mergers, and baryonic feedback — and significant system-to-system variability would generically arise. The empirical robustness of a single acceleration scale instead suggests the existence of an intrinsic infrared regulator that constrains macroscopic structure.

In quantum field theory and many-body systems, long-distance behaviour is governed by the lowest excitations of the spectrum. A gapless infrared spectrum yields algebraic decay of correlations and selects no preferred length scale, whereas a spectrum bounded from below by a non-zero gap produces exponential decay and dynamically introduces a finite correlation length [15]. The presence of such a gap, therefore, provides a natural mechanism for limiting large-distance coherence and selecting a characteristic structural scale.

Nevertheless, the mere existence of a gap does not uniquely determine the rigidity of that scale under non-linear evolution. If the spectrum above threshold is continuous, long-distance behaviour is controlled by near-threshold states and may depend on details of the spectral density, leaving room for environmental sensitivity. By contrast, if the infrared dynamics organizes into a lowest-weight discrete structure, the ground state is isolated and spectrally protected, so that macroscopic coherence is governed by a single dominant mode intrinsic to the dark sector. While discreteness is not logically required, it offers a particularly natural framework for understanding the observed universality and small intrinsic scatter of the galactic acceleration scale.

In this view, self-gravitating equilibria inherit — rather than dynamically generate — a characteristic structural scale, and the RAR may be interpreted as the large-

distance gravitational imprint of this intrinsic infrared organization.

C. Cosmological Background; the $\beta \rightarrow \psi$ Epoch

We focus on the cosmological interval extending from the onset of the standard post-inflationary expansion era (point β) to the epoch at which the total active gravitational mass density vanishes (point ψ), i.e., the $\beta \rightarrow \psi$ phase in Fig. 1. This interval encompasses radiation domination, matter domination, and essentially the entire epoch of non-linear structure formation during which galactic halos assemble and relax. Any mechanism proposed to account for the galactic acceleration scale must therefore satisfy two basic requirements:

1. It must persist dynamically throughout this cosmological window.
2. It must remain stable under the environmental diversity and non-linear gravitational evolution characteristic of structure formation.

We hypothesize that a long-lived gluonic vacuum sector survives throughout the $\beta \rightarrow \psi$ epoch and becomes effectively decoupled from the thermal plasma after confinement. The relevant late-time degrees of freedom are taken to be colour-singlet bosonic excitations, which we model as scalar di-gluonic bound states, as naturally expected in a confining non-Abelian gauge theory. This assumption is conservative at the level of field content; no new fundamental particles are introduced. Instead, we posit that a coherent component of the gluonic vacuum persists as a dynamically distinct infrared sector whose long-wavelength organization differs from that of ordinary hadronic matter. Concretely, “decoupled” means that number-changing processes and couplings to the Standard Model become inefficient after confinement, so that the comoving abundance (or coherent fraction) of the colour-singlet sector is effectively conserved on cosmological timescales and not re-thermalized by the baryonic plasma.

A crucial microscopic input is provided by the QCD trace anomaly. Classical Yang-Mills theory is scale invariant and contains no intrinsic mass parameter. At the classical level, one has:

$$T^\mu{}_\mu = 0.$$

Thus, no preferred intrinsic infrared scale set by gauge dynamics is selected. Quantum effects, however, break this scale invariance. Renormalization induces a non-vanishing vacuum expectation value of the energy-momentum trace dominated by gluonic contributions [16]:

$$\langle T^\mu{}_\mu \rangle \neq 0,$$




FIG. 1: Schematic representation of the three main stages of the Λ CDM cosmological evolution, showing the Hubble radius $L = H^{-1}$ as a function of the scale factor $a(t)$ (or $R(t)$), both plotted on logarithmic scales. The primordial inflationary phase extends from point α to β , followed by the standard expansion era from β to ψ , and finally by the late-time accelerated expansion (late inflation) from ψ to ω . The labeled points correspond to the following physical events:

- α : past informational event horizon;
- β : Big-Bang onset;
- γ : electroweak symmetry breaking;
- δ : color confinement (DM birth?);
- ϵ : transition from radiation to matter domination;
- ψ : vanishing of the total active gravitational mass;
- ω : future informational event horizon.

During primordial inflation, quantum fluctuations of cosmological relevance exit the comoving Hubble horizon; they subsequently re-enter it during the standard expansion era and finally exit again during the late accelerated phase. Only fluctuations with comoving wavelengths in the finite interval $\lambda_- \leq \lambda \leq \lambda_+$ ever cross the comoving Hubble horizon. Modes with $\lambda < \lambda_-$ remain always sub-horizon, while modes with $\lambda > \lambda_+$ remain always super-horizon. The bounds λ_- and λ_+ define respectively the past and future informational horizons represented by the points α and ω . The evolution and horizon crossing of cosmological perturbation modes follow the standard Mukhanov-Sasaki formalism [17–19], and admit a model-independent interpretation within the effective field theory of inflation [20]. For more details see the pedagogical [21].

thereby implementing dimensional transmutation. The renormalization-group flow replaces the classically scale-free Yang-Mills sector by a quantum theory characterized by a renormalization-group-invariant scale (e.g., Λ_{QCD}). Even in the absence of explicit mass parameters in the Lagrangian, the quantum gluonic vacuum therefore necessarily contains a dynamically generated dimensionful infrared input; the long-wavelength effective dynamics of the gluonic sector cannot remain strictly scale invariant.

This observation is conceptually essential as in a strictly scale-invariant pure gauge sector, any intrinsic infrared length attributable to the gauge dynamics would have to be imposed externally (e.g., via boundary con-

ditions or environmental scales), whereas dimensional transmutation guarantees that a dimensionful infrared input emerges from the internal non-perturbative dynamics itself.

It is equally important, however, to distinguish the existence of a dimensionful scale from the existence of a spectral gap. Dimensional transmutation ensures that the theory is not scale-free, but it does not determine the structure of the infrared excitation spectrum. A theory may possess an intrinsic scale while still supporting gapless modes or continuous spectra in certain channels. The anomaly, therefore, provides a necessary microscopic seed for infrared structure, but it does not guarantee dis-

creteness, boundedness from below, or spectral rigidity.

If the empirical universality of the RAR is to be attributed to a dynamically selected macroscopic length, an additional structural ingredient is required. The relevant colour-singlet sector must contain an isolated lowest excitation above the vacuum, and this lowest scale must remain stable under environmental perturbations and non-linear gravitational evolution. In other words, dimensional transmutation explains why an infrared scale can exist; one must still explain why that scale becomes a protected, macroscopically controlling correlation length rather than remaining sensitive to near-threshold continua or model-dependent spectral densities.

Our central hypothesis, therefore, is that the late-time infrared gluonic sector organizes into a spectrally rigid lowest-weight structure. Confinement motivates the expectation of isolated colour-singlet bound states; we elevate this expectation to a structural principle by postulating that the infrared sector admits a lowest-weight unitary realization. Representation theory then provides a natural mechanism for protecting the gap; the presence of a positive-energy generator and a lowest-weight state isolates the ground level and organizes excitations into a discrete tower anchored at that state. Throughout, “energy” refers to the eigenvalue of the compact $\mathfrak{so}(2)$ generator in this effective infrared algebraic description, not to the global cosmological energy of the Friedmann-Robertson-Walker background.

In four spacetime dimensions, the requirement of Lorentz covariance together with a positive-energy lowest-weight unitary structure strongly restricts the admissible non-compact symmetry algebras. A particularly economical and well-studied choice admitting positive-energy lowest-weight unitary irreducible representations is $\mathfrak{so}(2,3)$, which is the isometry algebra of four-dimensional Anti-de Sitter (AdS) spacetime [22–25]. Its maximal compact subalgebra:

$$\mathfrak{so}(2) \oplus \mathfrak{so}(3)$$

contains a compact $\mathfrak{so}(2)$ generator that defines a Hamiltonian bounded from below. We therefore postulate that the infrared gluonic dynamics organizes into a lowest-weight unitary irreducible representation of $\mathrm{SO}(2,3)$. In such a realization, the quadratic Casimir fixes a discrete spectrum of $\mathfrak{so}(2)$ eigenvalues E_n with a lowest level $E_0 > 0$ and characteristic spacing set by a single infrared scale $\varkappa$ [22–25]. For the class of lowest-weight modules considered here, this may be represented schematically as [22–25]:

$$E_n = E_0 + n\varkappa, \quad n = 0, 1, 2, \dots, \quad E_0 > 0,$$

where $\varkappa$ sets both the gap above the ground level and the typical level spacing.¹ The associated infrared corre-

lation length scales as:

$$r_c \sim \varkappa^{-1},$$

since, in the effective infrared theory, $\varkappa$ plays the role of an infrared mass parameter controlling the exponential falloff of colour-singlet correlators (up to factors determined by the effective dispersion) [15]. Because the gap arises from the lowest-weight unitary structure, it is representation-theoretically protected; it cannot be continuously removed without leaving the lowest-weight unitary class. In this way, a dimensionful input generated by dimensional transmutation is promoted to a spectrally rigid and dynamically stable infrared length.

The appearance of “AdS” symmetry is therefore strictly representation-theoretic. We do not assume that cosmological spacetime is globally AdS. The background Universe remains described by a Friedmann-Robertson-Walker geometry that is observationally close to spatially flat. The $\mathrm{SO}(2,3)$ structure characterizes only the algebraic organization of long-wavelength modes within the gluonic dark sector.

This subsection thus establishes the logical program pursued in the remainder of the paper:

1. Construct an effective infrared description of the colour-singlet gluonic sector.
2. Demonstrate how a lowest-weight $\mathrm{SO}(2,3)$ organization yields a protected gap and correlation length r_c .
3. Derive the gravitational imprint of this correlation length that sets the galactic acceleration scale.

D. From Spectral Gap to Galactic Acceleration

At sufficiently large occupation number, Bose statistics favor a macroscopic population of the isolated lowest-weight state. It is well established in ultracold atomic physics that Bose-Einstein condensation does not require strong interparticle binding; the phase transition already occurs in an ideal Bose gas and is driven primarily by Bose statistics and spectral discreteness, rather than by dynamical attraction. In laboratory systems, this discreteness is typically produced by a finite volume or an external trapping potential. In the present framework, however, no external trap is introduced; the discrete spectrum arises intrinsically from the lowest-weight $\mathrm{SO}(2,3)$ structure itself, through the compact $\mathfrak{so}(2)$ generator that organizes the infrared modes into a tower bounded from below. Condensation in the decoupled

discrete tower of $\mathfrak{so}(2)$ eigenvalues anchored at a lowest weight; the form above captures the existence of a single controlling spacing scale.

¹ More generally, lowest-weight $\mathrm{SO}(2,3)$ representations yield a

colour-singlet sector thus follows as a statistical consequence of its representation-theoretic spectral organization, rather than from additional binding dynamics. The resulting ground-state configuration defines a finite-mass self-gravitating structure with characteristic size r_c .

Coupling this condensate to Newtonian gravity produces a halo of mass M_h with characteristic acceleration:

$$g_\star = \frac{GM_h}{r_c^2}.$$

The galactic acceleration scale, therefore, emerges as the macroscopic imprint of the representation-theoretic parameter $\varkappa$. No modification of gravitational dynamics is required.

In this framework, the RAR is interpreted as the gravitational manifestation of an infrared spectral gap in a gluonic dark sector that persists throughout the $\beta \rightarrow \psi$ cosmological epoch.

We now follow this program in detail, showing how a lowest-weight $\text{SO}(2, 3)$ realization of the infrared gluonic sector yields a spectrally protected length scale whose gravitational imprint accounts for the observed universality of the RAR.

II. GLUONIC BOSE-EINSTEIN CONDENSATE IN AN EFFECTIVE ADS BACKGROUND

In an effective AdS description characterized by a negative curvature scale, it is convenient to introduce the curvature parameter [24, 25]:

$$\varkappa = \sqrt{\frac{|\Lambda_{\text{AdS}}^{\text{eff}}|}{3}}, \quad (1)$$

where $\Lambda_{\text{AdS}}^{\text{eff}} < 0$ denotes the effective cosmological constant defining the curvature radius $\varkappa^{-1}$ of the auxiliary AdS background. Restoring physical dimensions by reintroducing the speed of light c and Planck's constant $\hbar$, it is natural to define the corresponding frequency scale:

$$\omega_{\text{AdS}} = \varkappa c. \quad (2)$$

In such an effective AdS background, the rest energy of a massive excitation with (effective) spin s is given by [24, 25]:

$$E_{\text{AdS}}^{\text{rest}} = \left[m^2 c^4 + \hbar^2 \omega_{\text{AdS}}^2 \left(s - \frac{1}{2} \right)^2 \right]^{1/2} + \frac{3}{2} \hbar \omega_{\text{AdS}}. \quad (3)$$

We consider an assembly of $N_G \gg 1$ non-interacting, colour-neutral scalar di-gluonic excitations with discrete energy levels:

$$E_n = E_{\text{AdS}}^{\text{rest}} + n \hbar \omega_{\text{AdS}}, \quad n = 0, 1, 2, \dots, \quad (4)$$

and degeneracy:

$$G_n = \frac{(n+1)(n+3)}{2}. \quad (5)$$

Note that, introducing the dimensionless parameter:

$$\nu_0 = \frac{E_{\text{AdS}}^{\text{rest}}}{\hbar \omega_{\text{AdS}}}, \quad (6)$$

the spectrum may equivalently be written as:

$$E_n = \hbar \omega_{\text{AdS}} (n + \nu_0). \quad (7)$$

These excitations form an ensemble equivalent to isotropic harmonic oscillators in three spatial dimensions and may therefore be treated as an ideal Bose gas in the grand-canonical ensemble. At temperature T , the chemical potential μ is determined by the condition that the total occupation number equals the total number of bosons:

$$N_G = \sum_{n=0}^{\infty} \frac{G_n}{\exp \left[\frac{E_n - \mu}{k_B T} \right] - 1}. \quad (8)$$

For sufficiently large N_G , the system undergoes Bose-Einstein condensation at the critical temperature:

$$T_{\text{critical}} \sim \frac{\hbar \omega_{\text{AdS}}}{k_B} \left(\frac{N_G}{\zeta(3)} \right)^{1/3}, \quad \zeta(3) \approx 1.202. \quad (9)$$

This critical temperature coincides with the standard result for an ideal Bose gas in an isotropic three-dimensional harmonic potential. In the present framework, however, no external trapping potential is imposed. The harmonic structure of the spectrum arises intrinsically from the effective AdS geometry, which fixes the kinematical organization of the energy levels and thereby plays the role normally attributed to an external confining potential.

The eigenstates underlying this construction are solutions of the second-order wave equation [22–25]:

$$\left(Q_{\text{AdS}}^{(1)} - \zeta(\zeta - 3) \right) \varphi(x) = 0, \quad (10)$$

where $Q_{\text{AdS}}^{(1)}$ denotes the quadratic Casimir operator of the AdS isometry group $\text{SO}(2, 3)$. In the present work, we restrict attention to the scalar sector, so that the Casimir acts on a scalar field $\varphi(x)$. This choice is physically motivated by the expectation that the lightest colour-singlet gluonic excitation in QCD is the scalar glueball 0^{++} (see, for instance, Ref. [26]), which naturally dominates the infrared spectrum of the gluonic sector. Higher-spin glueball states correspond to heavier excitations and do not modify the lowest-weight spectral structure that governs the large-distance condensate dynamics considered here.

The parameter ζ fixes the representation and determines the lowest weight of the corresponding unitary irreducible module. Below, we make this operator realization explicit and review the associated normalized eigenfunctions that form the discrete lowest-weight tower.

A. Embedding, Signature, and Fixed-Casimir Equation

We realize four-dimensional AdS as a hyperboloid embedded in the five-dimensional ambient space $\mathbb{R}^{2,3}$ with coordinates X^A ($A = 0, 1, 2, 3, 5$)² equipped with metric of signature $(+, -, -, -, +)$:

$$\eta_{AB} dX^A dX^B = (dX^0)^2 - (d\mathbf{X})^2 + (dX^5)^2, \quad (11)$$

where $\mathbf{X} = (X^1, X^2, X^3)$ and $(d\mathbf{X})^2 = (dX^1)^2 + (dX^2)^2 + (dX^3)^2$. Four-dimensional AdS spacetime is identified with the hyperboloid defined by:

$$(X)^2 = X \cdot X = \eta_{AB} X^A X^B \\ = (X^0)^2 - \|\mathbf{X}\|^2 + (X^5)^2 = \kappa^{-2}. \quad (12)$$

The constant κ^{-1} therefore plays the role of the AdS curvature radius, fixing the scalar curvature and setting the intrinsic infrared length scale that controls the representation-theoretic spectrum discussed above.

The generators of $\text{SO}(2, 3)$ in the scalar (orbital) representation are realized as first-order differential operators on the ambient space:

$$L_{AB} = X_A \partial_B - X_B \partial_A, \quad (13)$$

where $\partial_A = \partial/\partial X^A$. These satisfy the $\mathfrak{so}(2, 3)$ commutation relations with indices raised and lowered by the ambient metric η_{AB} . The associated quadratic Casimir operator is given by:

$$Q_{\text{AdS}}^{(1)} = -\frac{1}{2} L_{AB} L^{AB}. \quad (14)$$

In terms of the flat five-dimensional d'Alembertian in the ambient space:

$$\square_5 = \eta^{AB} \partial_A \partial_B = \partial_0^2 - \nabla_{\mathbf{X}}^2 + \partial_5^2, \quad (15)$$

one obtains the useful identity (see, for instance, Ref. [22]):

$$Q_{\text{AdS}}^{(1)} = -(X \cdot X) \square_5 + (X \cdot \partial)^2 + 3 X \cdot \partial, \quad (16)$$

where $X \cdot \partial = X^A \partial_A$. Upon restriction to the hyperboloid $X^2 = \kappa^{-2}$, the Casimir eigenvalue equation (10) becomes an explicit second-order partial differential equation on AdS:

$$\left(-\kappa^{-2} \square_5 + (X \cdot \partial)^2 + 3 X \cdot \partial - \zeta(\zeta - 3) \right) \varphi(X) = 0. \quad (17)$$

Different metric sign conventions modify the overall sign of $\square_5$ and consequently of $Q_{\text{AdS}}^{(1)}$; however, the eigenvalue equation (10) is invariant up to this global sign. In particular, the spectrum determined by $\zeta(\zeta - 3)$ remains unchanged.

B. Global AdS Coordinates and Wave Operator

A convenient representation of the AdS metric in global static coordinates $x = (t, \underbrace{r, \theta, \phi}_{\mathbf{x}})$ is:

$$ds^2 = (1 + (\kappa r)^2) c^2 dt^2 - \frac{dr^2}{1 + (\kappa r)^2} - r^2 d\Omega_2^2, \quad (18)$$

where $t \in \mathbb{R}$ denotes the global AdS time, $r \geq 0$ is the radial coordinate on the spatial sections, and $d\Omega_2^2 = d\theta^2 + \sin^2 \theta d\phi^2$ is the metric on the unit two-sphere $\mathbb{S}^2$. The induced spatial line element reads:

$$d\ell^2 = \frac{dr^2}{(1 + (\kappa r)^2)} + r^2 d\Omega_2^2. \quad (19)$$

The non-vanishing components of the metric tensor, together with the corresponding components of its inverse, take the form:

$$g_{tt} = (1 + (\kappa r)^2) c^2, \quad g^{tt} = \frac{1}{(1 + (\kappa r)^2) c^2}, \\ g_{rr} = -\frac{1}{1 + (\kappa r)^2}, \quad g^{rr} = -(1 + (\kappa r)^2), \\ g_{\theta\theta} = -r^2, \quad g^{\theta\theta} = -\frac{1}{r^2}, \\ g_{\phi\phi} = -r^2 \sin^2 \theta, \quad g^{\phi\phi} = -\frac{1}{r^2 \sin^2 \theta}. \quad (20)$$

The determinant of the metric tensor is:

$$\det g = -c^2 r^4 \sin^2 \theta. \quad (21)$$

The covariant d'Alembertian on AdS in these coordinates is given by:

$$\square_{\text{AdS}} \varphi = \frac{1}{\sqrt{-g}} \partial_\mu \left(\sqrt{-g} g^{\mu\nu} \partial_\nu \varphi \right), \quad (22)$$

which, after a straightforward computation, yields:

$$\square_{\text{AdS}} \varphi = \frac{1}{(1 + (\kappa r)^2) c^2} \partial_t^2 \varphi - (1 + (\kappa r)^2) \partial_r^2 \varphi \\ - \left(\frac{2}{r} + 4\kappa^2 r \right) \partial_r \varphi - \frac{1}{r^2} \Delta_{\mathbb{S}^2} \varphi, \quad (23)$$

where:

$$\Delta_{\mathbb{S}^2} = \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\phi^2 \quad (24)$$

denotes the Laplace-Beltrami operator on the unit two-sphere.

For a scalar field belonging to a unitary irreducible representation labeled by ζ , the fixed-Casimir equation (10) can be recast, when expressed in global AdS coordinates,

² The label '4' is intentionally omitted, leaving room for a possible extension to conformal representations in four spacetime dimensions [27].

as a Klein-Gordon-type equation [24, 25]:³

$$(\square_{\text{AdS}} + \varkappa^2 \zeta(\zeta - 3)) \varphi(x) = 0. \quad (25)$$

The quantity $\varkappa^2 \zeta(\zeta - 3)$ therefore plays the role of an effective squared mass term. In particular, its value is constrained by the Breitenlohner-Freedman bound, ensuring the stability of scalar fields in AdS spacetime.

Consistent with the representation-theoretic interpretation, the rest energy (3) of a free excitation in AdS is determined by the curvature scale $\varkappa = \sqrt{\frac{|\Lambda_{\text{AdS}}^{\text{eff}}|}{3}}$ [24, 25]:

$$E_{\text{AdS}}^{\text{rest}} = \hbar \varkappa \zeta. \quad (26)$$

Here ζ is precisely the lowest eigenvalue of the time-translation generator L_{05} , which plays the role of the Hamiltonian in the corresponding unitary irreducible representation of the AdS group.

III. ADS MODES AND THEIR NORMALIZATION

A. General Expression

The normal-mode solutions of Eq. (25) were derived by Fronsdal in Ref. [22]. For $\zeta > 3/2$, they are given by:

$$\begin{aligned} \varphi_{nlm}(x) &= C_{nl} Y_{lm}(\theta, \phi) e^{-i(\zeta+2n+l)\omega_{\text{AdS}}t} \\ &\times (\varkappa r)^{2n+l} (1 + (\varkappa r)^2)^{-\frac{1}{2}(\zeta+2n+l)} \\ &\times {}_2F_1\left(-n, -n-l-\frac{1}{2}; \zeta-\frac{1}{2}; -\frac{1}{(\varkappa r)^2}\right), \end{aligned} \quad (27)$$

with:

$$\begin{aligned} C_{nl} &= \sqrt{\frac{c}{\pi}} \frac{\varkappa^2}{\Gamma(\zeta - \frac{1}{2})} \\ &\times \left[\frac{(\zeta - \frac{3}{2}) \Gamma(\zeta + n + l) \Gamma(\zeta + n - \frac{1}{2})}{\Gamma(n + l + \frac{3}{2}) \Gamma(n + 1)} \right]^{1/2}. \end{aligned} \quad (28)$$

The periodic time variable $t \in [0, 2\pi/\omega_{\text{AdS}}]$, with $\omega_{\text{AdS}} = \varkappa c$, is defined by [24]:

$$e^{-i\omega_{\text{AdS}}t} = \frac{X^5 - iX^0}{\sqrt{(X^5)^2 + (X^0)^2}} = \frac{X^5 - iX^0}{\sqrt{\varkappa^{-2} + r^2}}. \quad (29)$$

³ For scalar fields, the quadratic Casimir of $\text{SO}(2,3)$ restricted to AdS manifold coincides (up to the conventional factor $-\varkappa^{-2}$) with the Laplace-Beltrami operator on AdS [25]:

$$\mathcal{Q}_{\text{AdS}}^{(1)} = -\varkappa^{-2} \square_{\text{AdS}}.$$

In the ambient-space realization (see Eq. (16)), this relation follows after restricting the field to the AdS hyperboloid $(X)^2 = \varkappa^{-2}$ and choosing a homogeneous extension away from the surface (for instance, of degree zero, $X \cdot \partial \varphi = 0$), so that the Casimir operator reduces to the intrinsic AdS d'Alembertian.

The spherical harmonics Y_{lm} are normalized to unity on the two-sphere, and the quantum numbers satisfy $n, l \in \mathbb{N} = \{0, 1, 2, \dots\}$ and $-l \leq m \leq l$. Each mode φ_{nlm} is a simultaneous eigenfunction of the AdS energy generator L_{05} , of the $\mathfrak{so}(3)$ Casimir $\mathbf{L}^2$, and of L_{12} . The corresponding eigenvalues are $\zeta + l + 2n$, $l(l+1)$, and m , respectively. The function ${}_2F_1(a, b; c; r)$ denotes the Gaussian hypergeometric function, defined by the convergent series:

$$\begin{aligned} {}_2F_1(a, b; c; r) &= \frac{\Gamma(c)}{\Gamma(a)\Gamma(b)} \sum_{p=0}^{\infty} \frac{\Gamma(a+p)\Gamma(b+p)}{\Gamma(c+p)} \frac{r^p}{p!} \\ &= 1 + \frac{a}{c} \frac{b}{1!} r + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{r^2}{2!} + \dots, \end{aligned} \quad (30)$$

for $c \neq 0, -1, -2, \dots$. This series possesses the following properties:

1. It converges in the open unit disk $|r| < 1$.
2. Its behaviour on the circle of convergence $|r| = 1$ is given by:
 - (a) divergent for $\text{Re}(a+b-c) \geq 1$,
 - (b) absolutely convergent for $\text{Re}(a+b-c) < 0$,
 - (c) conditionally convergent for $0 \leq \text{Re}(a+b-c) < 1$, the point $r = 1$ being excluded.
3. It reduces to a polynomial of degree p in r , when a or b is a negative integer $-p$ ($p = 0, 1, 2, 3, \dots$).

The natural invariant measure on AdS may be written as:

$$\begin{aligned} d\mu(x) &= \sqrt{-g} d^4x \\ &= r^2 \sin \theta dt dr d\theta d\phi \\ &= 2\varkappa^{-1} \delta(X^2 - \varkappa^{-2}) d^5X, \end{aligned} \quad (31)$$

where the last expression is the ambient-space form of the induced measure on the AdS hyperboloid $(X)^2 = \varkappa^{-2}$. With the normalization constant C_{nl} chosen as in (28), the modes form an orthonormal basis with respect to this inner product:

$$\int_{\text{AdS}} \overline{\varphi_{nlm}(x)} \varphi_{n'l'm'}(x) d\mu(x) = \delta_{nn'} \delta_{ll'} \delta_{mm'}. \quad (32)$$

Here, the integration domain is the AdS hyperboloid itself (not its universal covering), so the global time coordinate is taken to be periodic, $t \in [0, 2\pi/\omega_{\text{AdS}}]$.

B. The Low- ζ Sector, Boundary Behaviour, and Conformal Scalar Structure

a. Breakdown of the L^2 realization and its remedy. It is important to emphasize that, with respect to the standard inner product (32), the interval:

$$\frac{1}{2} \leq \zeta \leq \frac{3}{2} \quad (33)$$

contains modes that fail to be square-integrable in the ordinary L^2 sense. This loss of square integrability does not signal any pathology of the corresponding representations; rather, it indicates that the naive Hilbert-space realization is no longer adequate in this regime.

A consistent framework is nevertheless available. The singleton (Rac) case $\zeta = \frac{1}{2}$, as well as the intermediate range $\frac{1}{2} < \zeta \leq \frac{3}{2}$, can be treated by replacing the standard L^2 norm with a Klein-Gordon-type inner product, suitably renormalized at the AdS boundary, and by defining the physical state space as a quotient by null states (see Ref. [23]).

At the level of the radial integration, this change is reflected in a modification of the effective measure; in the range $\frac{1}{2} < \zeta \leq \frac{3}{2}$, the standard weight $x^2 dx$ is replaced by:

$$x^2(1+x^2)^{-1} dx, \quad (34)$$

which compensates for the slower decay of the modes near the boundary. The endpoint $\zeta = \frac{1}{2}$ (Rac) requires a separate and more delicate treatment. This distinction reflects the different boundary behaviour of the modes and is fully consistent with the renormalized Klein-Gordon framework appropriate to the low- ζ sector.

b. AdS/CFT interpretation and double quantization. This structure is precisely reflected in the AdS/CFT correspondence. For scalar modes in four-dimensional AdS, the near-boundary behaviour involves two independent fall-offs:

$$\begin{aligned} \phi(r) &\sim r^{-\Delta_-}, \quad \Delta_- = \zeta, \\ \phi(r) &\sim r^{-\Delta_+}, \quad \Delta_+ = 3 - \zeta. \end{aligned} \quad (35)$$

The interval $\frac{1}{2} < \zeta \leq \frac{3}{2}$ is exactly the regime in which both branches are admissible with respect to a suitably renormalized Klein-Gordon inner product.

The choice of Hilbert-space structure then selects a quantization; the standard L^2 construction retains the faster-decaying branch $\Delta_+ = 3 - \zeta$ (standard quantization), while the renormalized framework allows one to include the slower-decaying branch $\Delta_- = \zeta$ consistently (alternate quantization). The endpoint $\zeta = \frac{1}{2}$ gives $(\Delta_-, \Delta_+) = (\frac{1}{2}, \frac{5}{2})$ and reproduces the Rac singleton.

c. Conformal scalar and the special role of $\zeta = 2$. A particularly important situation arises for the conformally coupled scalar in four-dimensional AdS. The complementary values $\zeta = 2$ and $\zeta = 1$ yield the same Casimir eigenvalue:

$$\zeta(\zeta - 3) = -2, \quad (36)$$

and therefore correspond to the same conformally invariant wave equation. They should thus be viewed as two realizations of a single underlying conformal structure, associated with the reducible $SO(2, 3)$ module $D(2, 0) \oplus D(1, 0)$, which is naturally mixed by the action of the full conformal group.

Their distinction lies in their boundary behaviour; the $\zeta = 2$ modes correspond to the faster fall-off (standard quantization), whereas the $\zeta = 1$ modes realize the slower branch (alternate quantization). In modern terms, this system lies in the Breitenlohner-Freedman window and admits a one-parameter family of self-adjoint extensions, equivalently Robin boundary conditions, with the pure $\zeta = 2$ and $\zeta = 1$ realizations recovered as limiting cases.

In the present work, we particularly focus on the $\zeta = 2$ branch, as it is the smallest scalar lowest-weight representation compatible with the standard normalizable choice and, within our halo construction, leads to a finite total mass. The $\zeta = 1$ branch nevertheless remains conceptually essential as the complementary conformal partner in the full boundary-condition problem.

d. Energy density and selection by finite mass. This choice can be justified more concretely by examining the asymptotic behaviour of the static energy density. For a scalar mode with asymptotics $\phi(r) \sim r^{-\zeta}$, the dominant contribution to the energy density T^0_0 behaves, up to metric factors, as:

$$T^0_0(r) \sim |\partial_r \phi|^2 + m^2 |\phi|^2 \sim r^{-2\zeta-2} + r^{-2\zeta} \sim r^{-2\zeta}. \quad (37)$$

Taking into account the spatial volume element $\sim r^2 dr$, the total mass integral behaves at large r as:

$$M \sim \int^\infty r^2 T^0_0(r) dr \sim \int^\infty r^{2-2\zeta} dr. \quad (38)$$

This integral converges if and only if:

$$2 - 2\zeta < -1 \iff \zeta > \frac{3}{2}. \quad (39)$$

Thus, within the double-quantization window $\frac{1}{2} < \zeta \leq \frac{3}{2}$, the total mass necessarily diverges, reflecting the slow decay of the corresponding modes. The value $\zeta = 2$ is the smallest admissible choice for which the mass becomes finite, making it the natural candidate for a physically acceptable halo profile. This provides a direct dynamical justification for selecting the $\zeta = 2$ branch in our construction.

C. Ground State and First Excited Angular Modes

For $n = 0$, the hypergeometric function in (27) reduces to 1, and the modes simplify to:

$$\begin{aligned} \varphi_{0lm}(x) &= \varkappa^2 \sqrt{\frac{c}{\pi}} \left[B\left(l + \frac{3}{2}, \zeta - \frac{3}{2}\right) \right]^{-1/2} \\ &\times Y_{lm}(\theta, \phi) e^{-i(\zeta+l)\omega_{\text{AdS}} t} (\varkappa r)^l \frac{1}{(1 + (\varkappa r)^2)^{\frac{1}{2}(\zeta+l)}}, \end{aligned} \quad (40)$$

where $B(p, q) = \Gamma(p)\Gamma(q)/\Gamma(p+q)$ denotes the Euler (complete) beta function.

For $l = 0$, this gives us the expression for the ground state:

$$\varphi_{000}(x) = \frac{\varkappa^2 \sqrt{c}}{2\pi} \left[B\left(\frac{3}{2}, \zeta - \frac{3}{2}\right) \right]^{-1/2} \frac{e^{-i\zeta\omega_{\text{AdS}}t}}{(1 + (\varkappa r)^2)^{\frac{\zeta}{2}}}. \quad (41)$$

Note that $Y_{00} = 1/\sqrt{4\pi}$. The first angularly excited states ($n = 0, l = 1, m = -1, 0, 1$) are:

$$\begin{aligned} \varphi_{01m}(x) = & \varkappa^2 \sqrt{\frac{c}{\pi}} \left[B\left(\frac{5}{2}, \zeta - \frac{3}{2}\right) \right]^{-1/2} \\ & \times Y_{1m}(\theta, \phi) e^{-i(\zeta+1)\omega_{\text{AdS}}t} \frac{\varkappa r}{(1 + (\varkappa r)^2)^{\frac{1}{2}(\zeta+1)}}. \end{aligned} \quad (42)$$

D. Higher Radial Excitations $n \geq 1$

For $n \geq 1$, the hypergeometric function appearing in (27) truncates to a polynomial of degree p in $1/(\varkappa r)^2$. Consequently, the radial dependence can be written as a finite sum of terms of the form:

$$\frac{(\varkappa r)^{2n+l-2p}}{(1 + (\varkappa r)^2)^{\frac{1}{2}(\zeta+2n+l)}}, \quad (43)$$

with $p = 0, \dots, n-1$. Since our analysis focuses primarily on the ground state and the lowest excited levels, we do not pursue the general structure of these higher-order polynomials further here.

IV. INTERFACE BETWEEN DM BOSE-EINSTEIN CONDENSATION AND BARYONIC MATTER

In the present framework, the CDM component is modeled as a Bose-Einstein condensate of colour-singlet gluonic degrees of freedom whose long-wavelength modes are organized by an effective AdS spectral structure. The condensate ground state (41) corresponds to the lowest-energy configuration of this sector. It is isotropic and spherically symmetric, and its spatial extent is set by the curvature scale of the effective AdS environment. This introduces a characteristic halo radius:

$$r_c \sim \varkappa^{-1} = \sqrt{\frac{3}{|\Lambda_{\text{AdS}}^{\text{eff}}|}}. \quad (44)$$

For definiteness, and to anchor the discussion in a phenomenologically relevant regime, we shall frequently use the fiducial benchmark:

$$r_c = 1 \text{ kpc}, \quad M_h = 10^9 M_\odot, \quad \zeta = 2, \quad (45)$$

which is representative of a typical dwarf-galaxy halo. As was previously asserted, the choice $\zeta = 2$ corresponds to

the smallest normalizable scalar lowest-weight representation and leads to a finite total halo mass in the present construction.

In the absence of baryons, the condensate forms an approximately spherical halo dominated by the ground-state density profile. Baryonic structures such as stellar cores, bulges, or localized baryonic clumps generate additional gravitational potentials that act as perturbations on this background configuration. These perturbations generally break the spherical symmetry of the condensate and induce transitions from the ground state to excited AdS modes. In a multipole decomposition of the baryonic potential, the dominant response is controlled by the lowest angular excitations, leading to multipolar distortions of the DM density in the baryon-halo interface region.

A. Dipolar Deformation; $n = 0, l = 1$

The triplet of modes (42), with $(n, l) = (0, 1)$ and $m = -1, 0, 1$, has a dipolar angular structure:

$$\varphi_{01m}(x) \propto Y_{1m}(\theta, \phi) \frac{\varkappa r}{(1 + (\varkappa r)^2)^{\frac{1}{2}(\zeta+1)}}. \quad (46)$$

Any departure from exact spherical symmetry in the baryonic sector may be decomposed into spherical multipoles. The leading contribution is the dipolar component of the baryonic gravitational potential. In the presence of such a perturbation, the condensate wave-function can be written schematically as:

$$\Psi_{\text{DM}} \sim a_{000}\varphi_{000} + \sum_{m=-1}^1 a_{01m}\varphi_{01m} + \dots, \quad (47)$$

where the coefficients a_{01m} are generated by matrix elements of the perturbing potential between the ground state and the $(0, 1)$ triplet. To leading order, the response of the condensate is therefore dominated by these lowest angular excitations.

The occupation of these first angularly excited levels controls several physically important effects:

1. The polarisation of the condensate around baryonic mass concentrations. In analogy with the polarisation of a dielectric medium placed in an external electric field, the dipolar modes generate a spatial displacement of the condensate density toward the baryonic potential minimum.
2. The directional response of the DM distribution to an external gravitational field. Because the Y_{1m} harmonics encode a preferred direction, the excitation of the $(0, 1)$ modes produces anisotropic distortions of the halo density aligned with the baryonic mass distribution.
3. The formation of asymmetric DM wakes generated by moving baryonic objects. Such wakes arise from

the coherent excitation of dipolar modes in the condensate and provide a natural microscopic mechanism for dynamical friction acting on stars, stellar clusters, or satellite galaxies moving through the halo.

The $(n, l) = (0, 1)$ triplet carries the lowest non-zero angular momentum and therefore provides the first channel through which departures from spherical symmetry can be encoded. Its radial structure is also particularly suited to mediating the baryon-halo coupling. While its asymptotic decay matches that of the ground state, the additional factor $\varkappa r$ suppresses the mode at the origin and shifts its maximal radial weight toward radii of order the condensate scale $r \sim r_c = \varkappa^{-1}$. Consequently, these modes are naturally concentrated in the intermediate region where the halo profile changes from the inner core to the outer regime, which observationally corresponds to the radial domain where the baryonic and DM contributions to the gravitational potential become comparable. In this transition zone, the dipolar excitations efficiently transmit anisotropic perturbations from the baryonic sector to the condensate, generating the leading multipolar deformations of the halo density.

B. Breathing Mode; $n = 1, l = 0$

The first radial excitation corresponds to $(n, l) = (1, 0)$. As discussed above, the radial dependence of modes with $n \geq 1$ is governed by the truncated polynomial structure given in Eq. (43). In the present case, the hypergeometric function reduces to a polynomial of first degree in $1/(\varkappa r)^2$, and the radial profile takes the form:

$$\varphi_{100}(r) \propto \left((\varkappa r)^2 - \frac{3}{2\zeta - 1} \right) \frac{1}{(1 + (\varkappa r)^2)^{\frac{1}{2}(\zeta + 2)}}. \quad (48)$$

This function changes sign at the radial node:

$$r_{\text{node}} = \varkappa^{-1} \sqrt{\frac{3}{2\zeta - 1}}, \quad (49)$$

which separates an inner region ($r < r_{\text{node}}$) from an outer region ($r > r_{\text{node}}$) with opposite phase of the wavefunction.

When this mode is superposed with the ground state, the interference term in the density produces opposite perturbations inside and outside the node, corresponding to a redistribution of mass between the inner and outer halo. The resulting collective motion is spherically symmetric and corresponds to a “breathing” oscillation of the condensate, in which the halo alternately contracts and expands while preserving its overall symmetry.

For the phenomenologically relevant choice $\zeta = 2$, the node occurs at:

$$r_{\text{node}} = \varkappa^{-1} = r_c, \quad (50)$$

so that the breathing mode naturally redistributes mass across the characteristic halo scale.

In the presence of baryons, transitions between the ground state and this breathing mode — driven, for instance, by slow variations of the baryonic potential or by non-linear couplings — can lead to:

1. Slow radial pulsations of the DM core.
2. Episodic energy exchange between the condensate and the baryonic environment.
3. Modulation of the inner density profile through periodic redistribution of mass within the halo core.

C. Order-of-Magnitude Estimates

The characteristic scales of the condensate are set by the curvature parameter $\varkappa$ and by the effective di-gluonic mass m_{eff} as determined by Eqs. (3) (see also Eq. (26)):

$$m_{\text{eff}} = \frac{E_{\text{AdS}}^{\text{rest}}}{c^2} = \frac{\hbar}{r_c c} \zeta. \quad (51)$$

This quantity characterizes the lowest-weight representation of the AdS symmetry algebra and should not be confused with the proper mass m appearing in the underlying field theory.

a. Core radius. The ground-state density profile:

$$\rho_{\text{DM}}(r) \propto (1 + (\varkappa r)^2)^{-\zeta} \quad (52)$$

is approximately constant for $r \ll r_c \simeq \varkappa^{-1}$ and decreases as $r^{-2\zeta}$ for $r \gg r_c$. The profile, therefore, naturally exhibits a finite-density core followed by a power-law decay at large radii. Identifying r_c with the observed core radii of dwarf and low-surface-brightness galaxies, typically $r_c \sim 0.6$ -5 kpc, one obtains:

$$|\Lambda_{\text{AdS}}^{\text{eff}}| \simeq \frac{3}{r_c^2} \simeq 10^{-40} - 10^{-38} \text{ m}^{-2}. \quad (53)$$

This effective curvature scale is many orders of magnitude larger than the observed cosmological constant, $\Lambda_{\text{obs}} \sim 10^{-52} \text{ m}^{-2}$. This hierarchy simply reflects the fact that $\Lambda_{\text{AdS}}^{\text{eff}}$ is not a cosmological constant of space-time, but an effective parameter controlling the spectral organization of the gluonic dark sector on galactic scales, while the background Universe remains well described by an approximately spatially flat Friedmann-Robertson-Walker geometry.

b. Characteristic frequency. The AdS frequency ω_{AdS} is given by:

$$\omega_{\text{AdS}} = \varkappa c \simeq \frac{c}{r_c} \simeq 10^{-12} - 10^{-11} \text{ s}^{-1}, \quad (54)$$

for core radii $r_c \sim 0.6$ -5 kpc. This corresponds to oscillation periods:

$$T_{\text{AdS}} \sim 4 \times 10^{11} - 3 \times 10^{12} \text{ s}, \quad (55)$$

i.e., of order 10^4 – 10^5 years. These timescales are short compared to typical galactic dynamical times ($\simeq 10^8$ – 10^9 years), allowing the condensate to adjust quasi-adiabatically to slowly evolving baryonic potentials while still supporting internal collective oscillations (breathing and dipole modes).

c. Geometric scale from the Fronsdaal spectrum [22]. In an AdS background, global spatial translational invariance is absent, so spatial momentum is not a good quantum number for labeling field excitations. Instead, the physically relevant scales are determined by the discrete normal-mode spectrum of the covariant wave operator. As discussed above, for a scalar field on AdS, the Fronsdaal equation (see Eq. (27)) leads to a discrete family of global modes labeled by a radial quantum number $n \in \mathbb{N}$ and angular momentum l , with frequencies of the form:

$$\omega_{nl} = \underbrace{\kappa c}_{\omega_{\text{AdS}}} (l + 2n + \zeta) = \frac{c}{r_c} (l + 2n + \zeta), \quad (56)$$

where r_c is taken as the AdS curvature radius, and ζ is the parameter of quantum nature used throughout this work. This parameter is related to the AdS rest energy through Eq. (3) (see also Eq. (26)) and may be expressed in terms of an effective mass parameter m_{eff} (51).

d. Spectral structure and degeneracy. Equation (56) shows that the discrete normal-mode spectrum depends only on the combination $k = l + 2n$, so that:

$$\omega_k = \omega_{\text{AdS}}(k + \zeta), \quad k = 0, 1, 2, \dots \quad (57)$$

The spectrum is therefore evenly spaced, with uniform separation:

$$\Delta\omega = \omega_{\text{AdS}} = \frac{c}{r_c}, \quad (58)$$

or equivalently $\Delta E = \hbar\omega_{\text{AdS}} = \hbar c/r_c$ at the level of energies. For each fixed value of k , the corresponding level is degenerate; its multiplicity is the number of non-negative integer pairs (n, l) satisfying $l + 2n = k$, namely $\lfloor k/2 \rfloor + 1$, not including the usual $(2l + 1)$ degeneracy associated with the magnetic quantum number m . This degeneracy reflects the underlying $\text{SO}(2, 3)$ representation structure and expresses the fact that different radial and angular excitations may belong to the same energy level. Although the spacing ΔE is extremely small in absolute units for galactic values of r_c , this is simply the quantum manifestation of the macroscopic correlation length of the condensate. The relevant physics is therefore not that of isolated microscopic transitions, but of a coherent configuration whose large-scale structure is governed by the ordering of these modes. In this perspective, baryonic gravitational perturbations may be viewed as inducing controlled admixtures of degenerate or nearby levels, thereby deforming the ground-state halo profile. A detailed dynamical treatment of this mode mixing is left for future work.

The lowest mode ($n = 0 = l$) carries a characteristic inverse length scale of order $1/r_c$, and its spatial profile extends over a region of size $\mathcal{O}(r_c)$. The condensate is thus macroscopic in the precise sense that it occupies the lowest AdS normal modes whose spatial support is controlled by the curvature scale r_c , rather than by any microscopic wavelength.

The constituent mass m enters primarily through the rest energy and the scaling dimension ζ , while the spatial extension of the ground state is governed geometrically by $\Lambda_{\text{AdS}}^{\text{eff}}$ via r_c . Any flat-space de Broglie interpretation should thus be regarded only as a heuristic rephrasing of this spectral statement, not as a dynamical momentum estimate. This characterization is fully intrinsic to AdS geometry and does not rely on any notion of local particle momentum.

V. DM MASS PROFILES, ROTATION CURVES AND ACCELERATIONS

In this section, we examine the main properties of the corresponding DM halo structure, including the mass profile, circular velocity, and radial acceleration. For clarity, we first restrict attention to a purely spherically symmetric configuration dominated by the ground state. Within the AdS-Bose-Einstein condensate framework described above, the DM density profile associated with the ground state of the gluonic condensate is given by Eq. (52), or equivalently by:

$$\rho_{\text{DM}}(x) = \rho_c (1 + x^2)^{-\zeta}, \quad x = \kappa r = \frac{r}{r_c}. \quad (59)$$

We then illustrate the phenomenological implications of the model using a representative dwarf-galaxy halo and investigate how including the first radial excitation modifies the resulting profiles. The effective AdS curvature parameter is chosen so as to produce a core radius consistent with observationally inferred values for dwarf galaxies.

a. On the role of AdS coordinates and the physical measure. It is important to emphasize that, in the present construction, the effective AdS geometry is not interpreted as the physical spacetime geometry of the galactic halo. Rather, it provides an auxiliary representation-theoretic framework that organizes the infrared spectrum and the associated normal modes of the dark sector. The resulting ground-state profile is then mapped to physical space by identifying the dimensionless radial variable as $x = \kappa r = r/r_c$, where r denotes the usual radial coordinate in the weak-field spatial geometry relevant for galactic dynamics. Accordingly, all physical quantities considered in this section — such as the enclosed mass, circular velocity, and radial acceleration — are computed using the standard three-dimensional Euclidean measure, $d^3\mathbf{x} = 4\pi r^2 dr$, appropriate to the Newtonian regime. In this sense, the AdS structure encodes the infrared spectral organization of the condensate, whereas the halo phenomenology is formulated in

the ordinary physical space relevant for galactic dynamics.

A. Mass Profile Properties

For a fixed parameter pair $(\varkappa, \zeta)$, the DM mass enclosed within a sphere of radius r follows from integrating the density profile (59) over the corresponding spherical volume. Introducing the dimensionless radius $x = \varkappa r = r/r_c$, one obtains:

$$M_{\text{DM}}(< x, \zeta) = 4\pi\rho_c r_c^3 F(x, \zeta), \quad (60)$$

where the dimensionless mass function $F(x, \zeta)$ is defined by:

$$F(x, \zeta) = \int_0^x f(u, \zeta) du, \quad f(u, \zeta) = u^2(1+u^2)^{-\zeta}. \quad (61)$$

a. Relation to the incomplete beta function. The function $F(x, \zeta)$ can be expressed in closed form by introducing the variable:

$$t = \frac{u^2}{1+u^2}, \quad (62)$$

for which:

$$u^2 = \frac{t}{1-t}, \quad du = \frac{dt}{2t^{1/2}(1-t)^{3/2}}. \quad (63)$$

Substituting into Eq. (61), one finds:

$$\begin{aligned} F(x, \zeta) &= \int_0^x u^2(1+u^2)^{-\zeta} du \\ &= \frac{1}{2} \int_0^{\frac{x^2}{1+x^2}} t^{1/2}(1-t)^{\zeta-\frac{5}{2}} dt. \end{aligned} \quad (64)$$

This is precisely the incomplete beta function:

$$B_z(a, b) = \int_0^z t^{a-1}(1-t)^{b-1} dt, \quad (65)$$

with:

$$a = \frac{3}{2}, \quad b = \zeta - \frac{3}{2}, \quad z = \frac{x^2}{1+x^2}. \quad (66)$$

Therefore:

$$\begin{aligned} F(x, \zeta) &= \frac{1}{2} B_{\frac{x^2}{1+x^2}}\left(\frac{3}{2}, \zeta - \frac{3}{2}\right) \\ &= \frac{1}{3} \left(\frac{x^2}{1+x^2}\right)^{3/2} {}_2F_1\left(\frac{3}{2}, \frac{5}{2} - \zeta; \frac{5}{2}; \frac{x^2}{1+x^2}\right), \end{aligned} \quad (67)$$

where in the second line we used the standard hypergeometric representation of the incomplete beta function:

$$B_z(a, b) = \frac{z^a}{a} {}_2F_1(a, 1-b; a+1; z). \quad (68)$$

For $\zeta > \frac{3}{2}$, $F(x, \zeta)$ converges as $x \rightarrow \infty$ to the finite limit:

$$F(\infty, \zeta) = \frac{1}{2} B\left(\frac{3}{2}, \zeta - \frac{3}{2}\right). \quad (69)$$

Consequently, the total mass of the AdS-gluonic-condensate halo remains finite and is given by:

$$M_h = M_{\text{DM}}(< \infty, \zeta) = 4\pi\rho_c r_c^3 \frac{1}{2} B\left(\frac{3}{2}, \zeta - \frac{3}{2}\right). \quad (70)$$

This property is particularly noteworthy: although the density distribution extends formally to infinite radius, the total halo mass remains finite. This contrasts with several commonly used phenomenological halo profiles (see, for instance, Refs. [28–32]), whose enclosed mass diverges logarithmically or linearly at large radii unless an explicit truncation radius is imposed.

b. Regularized form. It is convenient to express the mass profile in a normalized (regularized) form:

$$\frac{M_{\text{DM}}(< x, \zeta)}{M_h} = \frac{F(x, \zeta)}{F(\infty, \zeta)} = \frac{B_{\frac{x^2}{1+x^2}}\left(\frac{3}{2}, \zeta - \frac{3}{2}\right)}{B\left(\frac{3}{2}, \zeta - \frac{3}{2}\right)}. \quad (71)$$

c. Central and global behaviour. Near the halo center ($x \ll 1$), using $F(x, \zeta) \sim x^3/3$, one recovers a constant-density core:

$$M_{\text{DM}}(< x, \zeta) \sim \frac{4}{3} \pi \rho_c r_c^3 x^3. \quad (72)$$

d. Mass concentration and ζ -dependence of the halo profile. For $\zeta > 1$, the function $f(x, \zeta) = x^2(1+x^2)^{-\zeta}$ reaches its maximum at:

$$x_{\text{max}} = (\zeta - 1)^{-1/2}, \quad (73)$$

as follows from the condition $d(\ln f)/dx = 0$, equivalent to $f'(x, \zeta) = 0$ for $x > 0$. Since $F'(x, \zeta) = f(x, \zeta)$, this point coincides with the inflection point of $F(x, \zeta)$. The maximal value is:

$$f_{\text{max}} = \frac{(\zeta - 1)^{\zeta-1}}{\zeta^\zeta}. \quad (74)$$

Note that the limiting case $\zeta = 1$, discussed in the literature, corresponds to non-spherical (disk-like) configurations rather than genuine spherical halos.

These properties show that increasing ζ leads simultaneously to a smaller total halo mass and to a stronger central concentration. Indeed, for the density profile $(1+x^2)^{-\zeta}$ (see Eq. (59)), larger values of ζ produce a progressively steeper fall-off at large radii:

$$(1+x^2)^{-\zeta} \sim x^{-2\zeta}, \quad \text{for } x \gg 1. \quad (75)$$

Consequently, the outer halo contributes less to the mass integral, and a larger fraction of the total mass becomes confined within the inner region $r \lesssim r_c$.

Figure 2 shows the normalized enclosed mass $M_{\text{DM}}(< x, \zeta)/M_h$ as a function of $x = r/r_c$ for $\zeta = 2, 5/2, 3, 4$. Figure 3 compares the normalized enclosed mass profile of the AdS-gluonic-condensate halo (for $\zeta = 2$) with those of the Navarro-Frenk-White (NFW) and Burkert profiles.

A key structural distinction emerges, summarized in Table I. In widely used phenomenological halo models,

the enclosed mass diverges at large radii and a finite halo mass can only be defined by introducing an arbitrary truncation scale (typically R_{200}). In contrast, the AdS-gluonic-condensate profile yields a finite total mass for $\zeta > \frac{3}{2}$ despite its formally infinite spatial support. The halo mass, therefore, arises intrinsically from the profile itself, rather than from an externally imposed cutoff.




FIG. 2: Normalized enclosed mass $M_{\text{DM}}(< x, \zeta)/M_h$ as a function of $x = r/r_c$ for $\zeta = 2, 5/2, 3, 4$. Vertical dashed lines indicate the inflection radii $x_{\text{inf}} = (\zeta - 1)^{-1/2}$. Larger values of ζ correspond to a stronger central concentration and a faster saturation of the total halo mass.




FIG. 3: Comparison of normalized enclosed mass profiles. The AdS-gluonic-condensate halo (here $\zeta = 2$) is compared with the Navarro-Frenk-White (NFW) and Burkert profiles, all normalized at the same outer radius $x_{\text{max}} = 6$. $M(< x) = M_{\text{DM}}(< x, \zeta)$ for the AdS-gluonic-condensate halo. Unlike phenomenological profiles, the present model naturally leads to a finite total mass without requiring an explicit cutoff.

Profile	$\rho(r \rightarrow 0)$	$\rho(r \rightarrow \infty)$	Core / Cusp	$M(< r \rightarrow \infty)$	Reference
Pseudo-isothermal	finite	$\sim r^{-2}$	Core	$\propto r$	Begeman et al. (1991)
Burkert	finite	$\sim r^{-3}$	Core	$\sim \ln r$	Burkert (1995)
NFW	$\sim r^{-1}$	$\sim r^{-3}$	Cusp	$\sim \ln r$	Navarro et al. (1997)
coreNFW	finite	$\sim r^{-3}$	Core	$\sim \ln r$	Read et al. (2016)
This work (Gluonic condensate)	finite	$\sim r^{-2\zeta}$	Core	finite for $\zeta > \frac{3}{2}$	Present work

TABLE I: Comparison of commonly used DM halo density profiles. Here, $\rho(r)$ denotes the standard three-dimensional mass density, so that the enclosed mass is $M(< r) = 4\pi \int_0^r \rho(r') r'^2 dr'$. The asymptotic behaviors quoted in the table refer to this volumetric density. Profiles such as pseudo-isothermal, Burkert, and NFW either lead to a linearly or logarithmically divergent total mass as $r \rightarrow \infty$, unless an explicit outer cutoff radius is introduced. By contrast, the AdS gluonic-condensate profile considered in this work, $\rho(r) \sim r^{-2\zeta}$, yields a finite total mass for $\zeta > \frac{3}{2}$ without requiring any *ad hoc* truncation, thereby providing a self-consistent and intrinsically regular halo model.

B. Rotation Curves

The circular velocity induced by the DM halo in its ground state follows from the usual relation for a spherically symmetric mass distribution:

$$v^2(x, \zeta) = \frac{GM_{\text{DM}}(< x, \zeta)}{r_c x} = 4\pi G \rho_c r_c^2 \frac{F(x, \zeta)}{x}, \quad (76)$$

where $x = r/r_c$ and $F(x, \zeta)$ is the dimensionless enclosed-mass function introduced in Sec. V A. It is convenient to write the velocity profile explicitly as:

$$v(x, \zeta) = \sqrt{4\pi G \rho_c r_c^2 \frac{F(x, \zeta)}{x}}. \quad (77)$$

Differentiating with respect to the dimensionless radius x yields:

$$\frac{dv(x, \zeta)}{dx} = \frac{4\pi G \rho_c r_c^2}{2v(x, \zeta)} \left(-\frac{F(x, \zeta)}{x^2} + \frac{f(x, \zeta)}{x} \right), \quad (78)$$

where $f(x, \zeta) = x^2(1+x^2)^{-\zeta}$ is the integrand appearing in the definition of $F(x, \zeta)$.

a. Central behaviour. Near the halo center ($x \ll 1$), using $F(x, \zeta) \sim x^3/3$, one finds:

$$v(x, \zeta) \sim \sqrt{\frac{4\pi G}{3} \rho_c} \underbrace{r_c x}_r. \quad (79)$$

Thus, the velocity grows linearly with radius, $v \propto r$, which is the characteristic behaviour of a solid-body rotation curve.

b. Asymptotic behaviour. For $x \gg 1$ and $\zeta > \frac{3}{2}$, the function $F(x, \zeta)$ approaches its asymptotic value $F(\infty, \zeta)$ (69). The circular velocity, therefore, becomes (see Eq. (70)):

$$v(x, \zeta) \sim \sqrt{\frac{GM_h(\zeta)}{r_c x}}, \quad (80)$$

which corresponds to a slowly declining Keplerian behaviour $v \propto r^{-1/2}$. For fixed (ρ_c, r_c) , increasing ζ leads to a smaller total halo mass $M_h(\zeta)$, and consequently to systematically lower asymptotic velocities.

c. Maximum of the rotation curve and comparisons with NFW [29] and Burkert [30] profiles. As illustrated in Figs. 4 and 5 (see Tables II and III for the explicit formulas), the rotation velocity reaches a maximum at a dimensionless radius that we denote by $x_{v;\text{max}}$. From Eq. (78), the extremum condition $dv/dx = 0$ immediately yields:

$$\frac{F(x_{v;\text{max}}, \zeta)}{x_{v;\text{max}}} = f(x_{v;\text{max}}, \zeta). \quad (81)$$

This relation admits a simple geometric interpretation. The quantity $F(x, \zeta)/x$ represents the slope of the straight line joining the origin to the point $(x, F(x, \zeta))$, while $f(x, \zeta) = F'(x, \zeta)$ is the slope of the tangent to the curve $F(x, \zeta)$. Equation (81) therefore states that, at $x = x_{v;\text{max}}$, the tangent to the curve $F(x, \zeta)$ passes through the origin.

The velocity maximum occurs at a radius $x_{v;\text{max}}$ larger than the inflection point x_{max} (73) of $F(x, \zeta)$, which is defined by $f'(x, \zeta) = 0$. Indeed, for $0 < x \leq x_{\text{max}}$ the function $f(x, \zeta)$ is strictly increasing, so that for any $x_o < x$ one has $f(x_o, \zeta) < f(x, \zeta)$. It follows that:

$$F(x, \zeta) = \int_0^x f(x_o, \zeta) dx_o < \int_0^x f(x, \zeta) dt = x f(x, \zeta), \quad (82)$$

and therefore:

$$\frac{F(x, \zeta)}{x} < f(x, \zeta), \quad \text{for all } 0 < x \leq x_{\text{max}}. \quad (83)$$

On the other hand, the maximum of the rotation curve is determined by the condition (81). Since this equality cannot hold for $x \leq x_{\text{max}}$, it follows that:

$$x_{v;\text{max}} > x_{\text{max}}. \quad (84)$$

Because Eq. (81) has no closed-form solution, the order-

ing is verified numerically; for example:

$$\begin{aligned}
 \zeta = 2 : \quad x_{\max} &= 1, \quad x_{v;\max} \approx 1.8253, \\
 \zeta = \frac{5}{2} : \quad x_{\max} &= \sqrt{\frac{2}{3}}, \quad x_{v;\max} = \sqrt{2}, \\
 \zeta = 3 : \quad x_{\max} &= \frac{1}{\sqrt{2}} \approx 0.7071, \quad x_{v;\max} \approx 1.1892, \\
 \zeta = 4 : \quad x_{\max} &= \frac{1}{\sqrt{3}} \approx 0.5773, \quad x_{v;\max} \approx 0.9403.
 \end{aligned}
 \tag{85}$$

In all cases one finds $x_{v;\max} > x_{\max}$.

Importantly, the dimensionless location $x_{v;\max}$ depends only on the shape parameter ζ and is therefore universal within the model, while the corresponding physical radius is simply:

$$r_{v;\max} = x_{v;\max} r_c. \tag{86}$$




FIG. 4: On the top, dimensionless rotation curves $v(x, \zeta)/\sqrt{4\pi G\rho_c r_c^2}$ as a function of $x = r/r_c$ for $\zeta = 2, 5/2, 3, 4$. The inner linear rise reflects the constant-density core, while the outer Keplerian decline follows from the finiteness of the total halo mass. Larger values of ζ lead to lower and more centrally concentrated velocity profiles. On the bottom, shape comparison of circular-velocity profiles $v_c(r)$. The AdS-gluonic-condensate halo (shown for $\zeta = 2, 3, 4$) is compared with the NFW and Burkert profiles. Each curve is normalized to its maximum value on the plotted interval in order to emphasize differences in shape independently of the amplitude set by (ρ_s, r_s) or (ρ_0, r_0) .




FIG. 5: Calibrated comparison of circular-velocity profiles. All models are expressed in terms of the same dimensionless radius $x = r/r_0$ (common scale radius) and are normalized such that $v_c(1)$ is the same for all curves. The AdS-gluonic-condensate model is shown for $\zeta = 2, 5/2, 3, 4$ and compared with the NFW and Burkert profiles. The dashed vertical line marks $x = 1$.

Model	$\rho(x)$	$M(< x)$
Gluonic condensate	$\rho_c(1+x^2)^{-\zeta}$	$4\pi\rho_c r_0^3 F(x, \zeta)$
NFW	$\frac{\rho_s}{x(1+x)^2}$	$4\pi\rho_s r_0^3 \left(\ln(1+x) - \frac{x}{1+x} \right)$
Burkert	$\frac{\rho_0}{(1+x)(1+x^2)}$	$\pi\rho_0 r_0^3 (\ln(1+x^2) + 2\ln(1+x) - 2\arctan x)$

TABLE II: Summary of density profiles and enclosed masses. Here, $x = r/r_0$ denotes the dimensionless radius, where r_0 is the scale radius of each model ($r_0 = r_c$ for the gluonic condensate, $r_0 = r_s$ for NFW, and $r_0 = r_B$ for Burkert). For the AdS-gluonic-condensate model, $F(x, \zeta) = \int_0^x u^2(1+u^2)^{-\zeta} du$.

Model	$\rho(x)$	$v_c^2(x) = GM(< x)/x$
Gluonic condensate	$\rho_c(1+x^2)^{-\zeta}$	$4\pi G\rho_c r_0^2 F(x, \zeta)/x$
NFW	$\frac{\rho_s}{x(1+x)^2}$	$4\pi G\rho_s r_0^2 \frac{\ln(1+x) - x/(1+x)}{x}$
Burkert	$\frac{\rho_0}{(1+x)(1+x^2)}$	$\pi G\rho_0 r_0^2 \frac{\ln(1+x^2) + 2\ln(1+x) - 2\arctan x}{x}$

TABLE III: Summary of density profiles and circular-speed laws in the same conditions as in Table II.

d. A remark on asymptotic falloff and the choice of measure. It is worth noting that both the NFW and Burkert profiles share the same large-radius decay, $\rho(x) \sim x^{-3}$. This is suggestive when compared with the gluonic-condensate profile $\rho(x) \propto (1+x^2)^{-\zeta}$, whose asymptotic behaviour is $\rho(x) \sim x^{-2\zeta}$. Formally, the exponent -3 corresponds to $\zeta = \frac{3}{2}$, with the standard mea-

sure $x^2 dx$ (see below Eq. (32)). By contrast, in the low- ζ sector discussed below Eq. (32), the limiting case $\zeta = \frac{1}{2}$ is associated with the modified measure $x^2(1+x^2)^{-1} dx$, so that the same asymptotic behaviour may also be viewed, in that sense, as suggestive of $\zeta = \frac{1}{2}$. This analogy should not be overinterpreted. In the present framework, the distinction between $\zeta = \frac{3}{2}$ and $\zeta = \frac{1}{2}$ is governed not by

the density falloff alone, but by the functional setting, in particular by the measure entering the norm and normalization problem. For phenomenological profiles such as NFW and Burkert, it is therefore not obvious a priori which of these two measures provides the more relevant comparison. A careful clarification of this point would require a more systematic analysis, which we leave for future work.

C. Radial Accelerations

The radial gravitational acceleration generated by the DM halo in its ground state is defined as:

$$g(r, \zeta) = \frac{v^2(r, \zeta)}{r} = 4\pi G \rho_c r_c^3 \frac{F\left(\frac{r}{r_c}, \zeta\right)}{r^2}. \quad (87)$$

In terms of the dimensionless radius $x = r/r_c$, this reads:

$$g(x, \zeta) = 4\pi G \rho_c r_c \frac{F(x, \zeta)}{x^2}, \quad (88)$$

with radial derivative:

$$\frac{dg(x, \zeta)}{dx} = 4\pi G \rho_c r_c \left(-2 \frac{F(x, \zeta)}{x^3} + \frac{f(x, \zeta)}{x^2} \right), \quad (89)$$

where, again, $f(x, \zeta) = x^2(1+x^2)^{-\zeta}$.

a. Central behaviour. As $x \ll 1$, using $F(x, \zeta) \sim x^3/3$, one finds:

$$g(x, \zeta) \sim \frac{4\pi G}{3} \rho_c \underbrace{r_c x}_r, \quad (90)$$

corresponding to a linear growth of the acceleration. This behaviour is independent of ζ and reflects once again the solid-body character of the constant-density core.

b. Asymptotic behaviour. For $x \gg 1$ and $\zeta > \frac{3}{2}$, the enclosed mass approaches a constant value, $F(x, \zeta) \rightarrow F(\infty, \zeta)$, so that the acceleration becomes:

$$g(x, \zeta) \sim \frac{GM_h(\zeta)}{r_c^2 x^2}, \quad (91)$$

which corresponds to the standard Newtonian inverse-square fall-off produced by a finite total halo mass.

c. Maximum of the radial acceleration. The radial acceleration reaches a maximum at the radius $x_{g;\max}$ determined by the condition:

$$\frac{F(x_{g;\max}, \zeta)}{x_{g;\max}} = \frac{1}{2} f(x_{g;\max}, \zeta). \quad (92)$$

This relation determines the position of the maximum of the radial acceleration. For the values of ζ considered in this work, one obtains:

$$\begin{aligned} \zeta = 2 : \quad x_{g;\max} &\approx 0.9682, \\ \zeta = \frac{5}{2} : \quad x_{g;\max} &= 1, \\ \zeta = 3 : \quad x_{g;\max} &\approx 0.8260, \\ \zeta = 4 : \quad x_{g;\max} &\approx 0.6766, \end{aligned} \quad (93)$$

which satisfy the ordering (see Eq. (85)):

$$x_{g;\max} < x_{v;\max}, \quad (94)$$

showing that the maximum of the radial acceleration is reached at a smaller radius than the maximum of the circular velocity.

As in the case of the mass and velocity profiles, the dimensionless position $x_{g;\max}$ depends only on the shape parameter ζ and is therefore universal, i.e., independent of the particular properties of a given galaxy. The corresponding physical radius is simply:

$$r_{g;\max} = x_{g;\max} r_c. \quad (95)$$

D. On the Core Radius

The previous analysis shows that the notion of a “core radius” is not unique and may be defined in several physically meaningful ways. Within the present framework, at least four natural definitions can be identified:

1. The value of x beyond which the central approximation $M_{\text{DM}}(< x, \zeta) \sim \frac{4}{3}\pi\rho_c r_c^3 x^3$ (72) is no longer valid, signaling the breakdown of the constant-density regime.
2. The value of $x = x_{\max}$ (73) corresponding to the inflection point of the enclosed-mass profile, where the curvature of $M_{\text{DM}}(< x, \zeta)$ changes sign and the mass growth begins to slow.
3. The value of $x = x_{v;\max}$ (81) (see also (85)) at which the circular velocity $v(x, \zeta)$ reaches its maximum.
4. The value of $x = x_{g;\max}$ (92) at which the radial acceleration $g(x, \zeta)$ attains its maximum.

Although these definitions are not identical, they all select radii of the same order, thereby identifying a characteristic scale of the halo set by r_c . This reflects the existence of an intermediate radial region separating the inner constant-density core from the outer domain in which the dynamics approaches the asymptotically Newtonian regime.

As will become clear in the following sections, this transition region plays a special physical role. It is in this radial range that the relative importance of baryonic and dark components typically becomes comparable, and where additional effects — such as the excitation of higher condensate modes or the response of the halo to baryonic perturbations — are expected to be most significant.

E. A Representative Example; $\zeta = 2$ in the Context of Dwarf Galaxies

The choice $\zeta = 2$ is the smallest admissible integer or half-integer value of the AdS representation (confor-

mal) parameter for which the scalar lowest-weight unitary irreducible representation is normalizable. Within the present family of halo profiles, and for fixed (ρ_c, r_c) , this case corresponds to the least centrally concentrated halo, since the density profile $(1 + x^2)^{-\zeta}$ then has the slowest large-radius decay among the admissible values of ζ (see the discussion following Eqs. (73)-(75)). Consequently, it also yields the largest total halo mass within this class. Moreover, the corresponding dimensionless velocity profile varies only weakly over the intermediate range $x = \mathcal{O}(1-5)$, producing an extended quasi-flat portion of the rotation curve (see the discussion following Eqs. (81)-(86)). This makes the case $\zeta = 2$ particularly relevant phenomenologically. For this value, the function $F(x, \zeta)$ (64) takes the closed form:

$$F(x, 2) = \frac{1}{2} \left(\arctan x - \frac{x}{1 + x^2} \right). \quad (96)$$

Accordingly, the total halo mass M_h (70) becomes:

$$M_h = \pi^2 r_c^3 \rho_c, \quad \text{or equivalently} \quad \rho_c = \frac{M_h}{\pi^2 r_c^3}. \quad (97)$$

The enclosed-mass profile $F(x, 2)$ has an inflection point $x_{\max} = 1$ (see Eq. (73)), corresponding to the physical radius $r \sim r_c$. At this point:

$$F(1, 2) = \frac{1}{8}(\pi - 2), \quad f(1, 2) = \frac{1}{4}. \quad (98)$$

The latter identity follows directly from Eq. (74). The rotation curve is still rising at this radius, since its maximum occurs at $x_{v;\max} > x_{\max} = 1$ (see Eq. (85)). By contrast, the radial acceleration reaches its maximum at a slightly smaller radius $x_{g;\max} < x_{\max} = 1$ (see Eq. (93)).

To illustrate the relevance of this configuration for dwarf galaxies, let us adopt the fiducial values (45).

Equation (97) then yields:

$$\rho_c \sim 0.10 M_\odot \text{ pc}^{-3}, \quad (99)$$

which lies within the range of central densities inferred from rotation-curve fits of dwarf and low-surface-brightness galaxies (see, for instance, Refs. [30–33]).

In this setting:

1. At the inflection radius $x = 1$ ($r \sim r_c$), the enclosed mass fraction is:

$$\frac{F(1, 2)}{F(\infty, 2)} \sim 0.18, \quad (100)$$

and the circular velocity (76) is:

$$v(1, 2) = \sqrt{\frac{GM_{\text{DM}}(< r_c)}{r_c}} \sim 28 \text{ kms}^{-1}. \quad (101)$$

2. At a few core radii, say $x \simeq 3$ ($r \simeq 3r_c$), the enclosed mass fraction increases to:

$$\frac{F(3, 2)}{F(\infty, 2)} \sim 0.6, \quad (102)$$

while the circular velocity (76) reaches:

$$v(3, 2) = \sqrt{\frac{GM_{\text{DM}}(< 3r_c)}{3r_c}} \sim 30 \text{ kms}^{-1}. \quad (103)$$

These values are characteristic of observed dwarf-galaxy rotation curves. In the following, illustrative Figs. 6 and 7 (in arbitrary units with $\rho_c = r_c = 1$) display the corresponding density and mass profiles, as well as the circular-velocity and radial-acceleration curves.




FIG. 6: Illustrative density (left) and enclosed-mass (right) profiles for $\zeta = 2$, in arbitrary units ($\rho_c = r_c = 1$).




FIG. 7: Illustrative circular-velocity (left) and radial-acceleration (right) profiles for $\zeta = 2$, in arbitrary units ($\rho_c = r_c = 1$).

F. Role of the First Radial Excitation

Including a small admixture of the first radial excitation $(n, l) = (1, 0)$:

$$\rho_{\text{DM}}(r) \sim \mathcal{N}(|\varphi_{000}(r)|^2 + \epsilon |\varphi_{100}(r)|^2), \quad \epsilon \ll 1, \quad (104)$$

provides a controlled mechanism for modifying the transition region between the inner solid-body rise and the quasi-flat portion of the rotation curve. Because the radial wave function φ_{100} has a node, its contribution produces a mild redistribution of mass across the characteristic scale $r \sim r_c$ (see Eq. (50)). The resulting profile can slightly flatten the rotation curve or generate a modest overshoot in the intermediate region, while leaving

the constant-density core essentially unchanged at very small radii.

In a more detailed phenomenological treatment, the parameter ϵ may be used to account for the weak curvature features often visible in high-quality dwarf-galaxy rotation curves. This benchmark, therefore, shows that, even within a minimal parameterization based on (r_c, ρ_c, ζ) , the inclusion of a tiny first-radial-excitation component can already produce rotation curves of realistic amplitude and shape, while preserving a direct connection with the underlying AdS-Bose-Einstein-condensate microphysics.

G. Superposition Near Baryons

In the absence of baryons, the condensate is well described by the spherically symmetric ground state (41) discussed above. In the vicinity of a baryonic structure, however, the gravitational potential generated by the baryons acts as a perturbation that couples the ground state to the lowest excited levels of the AdS spectrum. The DM wave-function is therefore naturally described as a superposition of the lowest-lying modes:

$$\begin{aligned} \Psi_{\text{DM}}(t, \mathbf{x}) &\sim a_{000} \varphi_{000}(t, \mathbf{x}) \\ &+ \sum_{m=-1}^1 a_{01m} \varphi_{01m}(t, \mathbf{x}) \\ &+ a_{100} \varphi_{100}(t, \mathbf{x}) + \dots \end{aligned} \quad (105)$$

This superposition becomes particularly relevant in the intermediate radial region $r \sim r_c$, set by the intrinsic condensate scale. For the halo profiles considered here, this is the radius at which the ground-state density profile begins to depart from its inner nearly constant-density behaviour and cross over to the outer decaying regime (see Sect. VA). It therefore marks the onset of the region where the halo structure is no longer controlled solely by the unperturbed core, but becomes increasingly sensitive to external gravitational perturbations. At the same time, the lowest excited modes have their largest radial support precisely in this domain (see Sects. IV A and IV B), so that the response of the condensate to the baryonic potential is expected to be maximal there. In the present framework, it is therefore natural to identify $r \sim r_c$ with the baryon-halo interface region, namely the radial domain in which baryonic gravity begins to produce an appreciable deformation of the condensate away from its spherically symmetric ground-state configuration. In particular, the condensate:

1. Responds to the tidal gravitational field generated by the baryonic mass distribution.
2. Develops multipolar deformations encoded by modes with $l \geq 1$; see Sect. IV A.
3. Undergoes breathing and higher radial oscillations associated with modes with $n \geq 1$; see Sect. IV B.
4. Exerts a gravitational backreaction on the baryonic component, potentially affecting rotation curves, dynamical friction, and the stability of stellar systems.

The first excited levels thus play a central physical role. They encode the leading deviations from spherical symmetry and quantify the dynamical susceptibility of the condensate to baryonic perturbations. In this picture, baryons do not simply move in a fixed DM potential; rather, they locally deform a coherent medium whose response is controlled by the lowest modes of the AdS

spectrum. This provides a natural framework for understanding how baryons can reshape the effective DM distribution in the interface region without destroying the global coherence of the condensate.

VI. DISCUSSION: RAR AND EMERGENT ACCELERATION SCALE

Having established the structure of the AdS-gluonic-condensate halo and its associated mass, velocity, and acceleration profiles, we now confront the model with one of the most stringent empirical constraints on galactic dynamics, namely the RAR.

As already discussed in the Introduction, one of the central results of the SPARC analysis [6–8] is the existence of a remarkably tight correlation between the observed radial acceleration:

$$g_{\text{obs}}(r) = \frac{v_{\text{obs}}^2(r)}{r}, \quad (106)$$

and the Newtonian acceleration generated by the baryonic mass distribution:

$$g_{\text{bar}}(r) = \frac{G M_{\text{bar}}(< r)}{r^2}. \quad (107)$$

This correlation takes the form of an approximately galaxy-independent functional relation:

$$g_{\text{obs}} = g_{\text{obs}}(g_{\text{bar}}), \quad (108)$$

and exhibits an unexpectedly small intrinsic scatter across galaxies spanning a broad range of masses, morphologies, and evolutionary histories.

The relation is controlled by a single characteristic acceleration scale:

$$g_{\dagger} \sim 10^{-10} \text{ ms}^{-2}, \quad (109)$$

which delineates the transition between the baryon-dominated and DM-dominated regimes. Its apparent universality, combined with the exceptional tightness of the observed correlation, makes the RAR one of the strongest observational constraints on models of galaxy dynamics.

Although the RAR is often discussed in the context of modified-gravity phenomenology, it may equally be interpreted as an empirical constraint on the coupling between baryons and DM within the framework of standard gravity [11–13]. In the present work, we adopt this latter viewpoint.

Within our framework, the law of gravity itself is not modified. The observed radial acceleration is assumed to follow from the standard superposition of baryonic and DM contributions, so that the decomposition:

$$g_{\text{obs}}(r) = g_{\text{bar}}(r) + g_{\text{DM}}(r) \quad (110)$$

remains valid at all radii. The novelty of the model, therefore, does not lie in a change of the gravitational

force law, nor in the introduction of an ad hoc interpolation formula, but in the physical nature of the dark component and in its dynamical response to the baryonic gravitational field.

In the scenario developed above, the DM halo is described as a coherent condensate whose long-wavelength dynamics are governed by an effective AdS spectral structure, dominated by its lowest-weight state. The presence of baryons perturbs this ground-state configuration and excites the lowest modes of the spectrum, so that the dark component reacts to the baryonic potential in a controlled and essentially universal manner. We now investigate whether, and under which conditions, this response can naturally generate an effective acceleration scale of order $g_{\dagger} \sim 10^{-10} \text{ ms}^{-2}$ without galaxy-by-galaxy fine tuning, thereby reproducing the observed RAR within an otherwise standard Newtonian framework.

A. Phenomenology of the RAR

As established above, the radial acceleration produced by the DM halo in its ground state can be written as (see Eqs. (88), (60), and (71)):

$$\begin{aligned} g_{\text{DM}}(x, \zeta) (= g(x, \zeta)) &= 4\pi G \rho_c r_c \frac{F(x, \zeta)}{x^2} \\ &= \frac{GM_{\text{DM}}(< x, \zeta)}{r_c^2 x^2} \\ &= g_{\star} \frac{1}{x^2} \frac{F(x, \zeta)}{F(\infty, \zeta)}, \end{aligned} \quad (111)$$

where:

$$g_{\star} = \frac{GM_h}{r_c^2} \quad (112)$$

defines an intrinsic acceleration scale associated with the halo mass M_h and its geometric core radius r_c .

For a fixed value of the shape parameter ζ , the DM contribution thus factorizes into a universal dimensionless shape function and a single acceleration scale $g_{\star}$. In particular, for $\zeta = 2$ and $x = x_{\text{max}} = 1$, corresponding both to the geometric core radius and to the inflection point of the mass profile (see Sect. VE), one finds:

$$g_{\text{DM}}(1, 2) (= g(1, 2)) \sim 0.18 g_{\star}, \quad (113)$$

a value slightly below the maximal radial acceleration, because the latter is attained at a slightly smaller radius, $x_{\text{g;max}} < x_{\text{max}} = 1$ (see Eq. (93)). At large radii, the same intrinsic scale governs the asymptotic decay as:

$$g_{\text{DM}}(x) (= g(x, \zeta)) \sim \frac{1}{x^2} g_{\star}. \quad (114)$$

Observationally, the SPARC RAR, with both g_{obs} and g_{bar} evaluated at the same galactocentric radius, can be written in the convenient form [6, 8]:

$$g_{\text{obs}} = g_{\text{bar}} \nu \left(\frac{g_{\text{bar}}}{g_{\dagger}} \right), \quad \nu(y) = \frac{1}{1 - e^{-\sqrt{y}}}, \quad (115)$$

where $g_{\dagger}$ is the characteristic acceleration scale entering the empirical relation. The interpolation function $\nu(y)$ satisfies the asymptotic behaviours:

$$\begin{aligned} \nu(y) &\longrightarrow 1, \quad \text{when } y \gg 1, \\ \nu(y) &\longrightarrow y^{-1/2}, \quad \text{when } y \ll 1. \end{aligned} \quad (116)$$

On the other hand, using the exact decomposition (110), the RAR (115) can be rewritten in terms of the DM acceleration alone as:

$$g_{\text{DM}} = g_{\text{bar}} \mu \left(\frac{g_{\text{bar}}}{g_{\dagger}} \right), \quad \mu(y) (= \nu(y) - 1) = \frac{1}{e^{\sqrt{y}} - 1}, \quad (117)$$

with the asymptotic behaviour:

$$\begin{aligned} \mu(y) &\longrightarrow 0, \quad \text{when } y \gg 1, \\ \mu(y) &\longrightarrow y^{-1/2}, \quad \text{when } y \ll 1. \end{aligned} \quad (118)$$

Then, in the low-acceleration regime ($g_{\text{bar}} \ll g_{\dagger}$, that is, $y \ll 1$), one obtains:

$$g_{\text{obs}} \sim g_{\text{DM}} \sim \sqrt{g_{\dagger} g_{\text{bar}}}, \quad (119)$$

so that the observed acceleration is dominated by the DM contribution and follows the characteristic square-root scaling of the RAR. In the opposite, high-acceleration regime ($g_{\text{bar}} \gg g_{\dagger}$, that is, $y \gg 1$), one finds:

$$g_{\text{obs}} \sim g_{\text{bar}}, \quad (120)$$

while the DM contribution becomes negligible:

$$g_{\text{DM}} \sim 0, \quad (121)$$

corresponding to a baryon-dominated dynamics.

Although the functions $g_{\text{obs}}(g_{\text{bar}})$ and $g_{\text{DM}}(g_{\text{bar}})$ are universal, individual galaxies populate only finite regions of the RAR plane. Each galaxy samples the relation over the range of radii accessible to its baryonic mass distribution, so that the allowed values of g_{bar} are bounded from above by a galaxy-dependent maximum, typically reached at small galactocentric radii [6–8].

SPARC data show that more massive galaxies attain larger values of g_{bar} , often exceeding 10^{-9} ms^{-2} , whereas low-mass dwarf galaxies remain confined to the low-acceleration regime and never reach $g_{\text{bar}} \gtrsim g_{\dagger}$ [8, 34].

A key structural property of the RAR is that the dimensionless function:

$$\frac{g_{\text{DM}}}{g_{\dagger}} = \frac{g_{\text{bar}}}{g_{\dagger}} \mu \left(\frac{g_{\text{bar}}}{g_{\dagger}} \right) \quad (122)$$

is non-monotonic. As g_{bar} increases, the DM acceleration g_{DM} reaches a maximum value $g_{\text{DM}}^{\text{max}}$ at a characteristic baryonic acceleration $g_{\text{bar}}^{\star}$. Since both quantities are determined entirely by the single scale $g_{\dagger}$, the values of $g_{\text{DM}}^{\text{max}}$ and $g_{\text{bar}}^{\star}$ are universal. This behaviour is illustrated in Fig. 8.




FIG. 8: Left: the function $\mu(y)$ entering the RAR decomposition $g_{\text{DM}} = g_{\text{bar}}\mu(g_{\text{bar}}/g_{\text{t}})$. Right: the product $y\mu(y) = g_{\text{DM}}/g_{\text{t}}$, exhibiting a universal maximum of the DM acceleration. Both axes are logarithmic.

The existence of a universal upper bound on g_{DM} has an immediate consequence: the intrinsic halo scale $g_{\star} = GM_h/r_c^2$ must itself be bounded. For a given core radius r_c , this translates into an upper bound on the halo mass.

Accordingly, galaxy profiles in the $[g_{\text{obs}}, g_{\text{bar}}]$ (or equivalently, $[g_{\text{DM}}, g_{\text{bar}}]$) plane naturally separate into two broad categories, depending on whether the maximum baryonic acceleration reached within a given galaxy, denoted $g_{\text{bar}}^{\text{max}}$, is larger or smaller than the universal scale $g_{\text{bar}}^{\star}$ at which the RAR predicts the maximum DM acceleration.

In the first category, the DM acceleration decreases as the baryonic acceleration increases, whereas in the second, it grows with it. As an illustration, Ref. [34] discusses two extreme members of the SPARC sample that belong to these two categories:

1. The massive spiral galaxy NGC 2841, with flat rotation velocities of order 300kms^{-1} , reaches $g_{\text{obs}} \sim g_{\text{bar}} \simeq 10^{-9}\text{ms}^{-2}$ within a few kiloparsecs of the center, where the DM contribution is negligible. At large radii ($\sim 70\text{kpc}$), g_{obs} decreases to $\sim 10^{-10.5}\text{ms}^{-2}$, while $g_{\text{bar}} \lesssim 10^{-11}\text{ms}^{-2}$, allowing the DM contribution to become significant.
2. The gas-dominated dwarf galaxy IC 2574, with rotation velocities below 80kms^{-1} , reaches a maximal observed acceleration of $\sim 10^{-10.7}\text{ms}^{-2}$ at radii of 8-10kpc, while the baryonic acceleration there is an order of magnitude smaller, indicating a regime dominated by DM.

B. From the Intrinsic DM Halo Scale to the RAR Scale

The AdS-Bose-Einstein condensation framework naturally suggests a line of reasoning that starts from the DM distribution itself. Within this perspective, the empirical correspondence:

$$M_{\text{DM}}(< r) \longrightarrow g_{\text{DM}}(r) \xrightarrow{\text{RAR}} g_{\text{bar}}(r) \longrightarrow M_{\text{bar}}(< r)$$

may be viewed as the manifestation of an underlying dynamical chain. The gravitational field generated by the condensate — determined primarily by its ground state together with the lowest excited modes — sets the large-scale gravitational environment in which baryons evolve (see Sect. V G). In this interpretation, the baryonic distribution is not statistically independent of the dark sector, but instead adjusts to the potential produced by a coherent halo whose structure is governed by a small number of intrinsic parameters.

From this viewpoint, the RAR is not regarded as a fundamental law of dynamics. Rather, it appears as an emergent phenomenological relation encoding the response of baryonic matter to a coherent DM background whose properties are fixed by the condensate spectrum. The observed one-parameter universality of the RAR then reflects the fact that the halo structure is largely controlled by a single intrinsic scale, so that different galaxies probe different regions of the same underlying gravitational configuration.

Within this picture, the emergence of a single effective acceleration scale follows directly from the condensate structure and from its response to the presence of baryons. The physical content of the SPARC RAR may therefore be translated into a constraint on the allowed correlations between the intrinsic condensate parameters

(M_h, r_c) and the baryonic mass distribution, once the halo shape parameter ζ is fixed.

The halo is characterized by a distinguished length scale r_c , together with a discrete excitation spectrum separated from the ground state by a gap of order:

$$\Delta E \sim \frac{\hbar c}{r_c}. \quad (123)$$

In the regime where the baryonic gravitational field acts as a weak perturbation of the condensate, the response of the system is expected to be dominated by the lowest excitations of this spectrum. To leading order, the condensate wave-function may therefore be described by a truncated expansion over the lowest AdS modes, parametrized by a small set of amplitudes: a_{000} for the ground state, together with a_{01m} and a_{100} for the first angular and radial excitations (see again Sect. V G). These coefficients encode the linear response of the condensate to the external baryonic potential and determine the corresponding modification of the halo density profile in the baryon-halo interface region.

A first-principles determination of these amplitudes would require a linear-response analysis formulated in the AdS mode basis, in which the baryonic gravitational potential induces mixing between the ground state and the lowest excited modes of the condensate. Such a calculation is expected to generate an effective acceleration scale $g_{\dagger} = g_{\dagger}(\mathcal{N}, \zeta)$ whose value is controlled primarily by the intrinsic condensate parameters and only weakly affected by galaxy-dependent details of the baryonic distribution. Carrying out this program would require a systematic treatment of the condensate response beyond the present phenomenological level, and is therefore left for future work.

Nevertheless, even at the phenomenological level, one may already extract a robust scaling relation. Evaluating the DM acceleration at radii of order the core scale, $r \sim r_c$, Eqs. (111) and (112) give:

$$g_{\text{DM}}(r \sim r_c) \sim \alpha(\zeta) g_{\star}, \quad (124)$$

where the coefficient $\alpha(\zeta)$ is a dimensionless shape factor obtained by evaluating the dimensionless halo-acceleration profile at radii of order r_c . In the simplest spherical ground-state description, it is determined entirely by the halo profile. In particular, for the phenomenologically relevant case $\zeta = 2$, one finds $\alpha(2) \sim 0.18$ at $r = r_c$ (see Eq. (113)); in this case r_c lies slightly outside the radius at which $g_{\text{DM}}(r)$ reaches its maximum (see Eq. (93)).

This estimate provides the natural bridge between the intrinsic halo scale $g_{\star}$ and the empirical RAR scale $g_{\dagger}$. Observationally, $g_{\dagger} \sim 10^{-10} \text{ ms}^{-2}$ characterizes the transition between baryon-dominated and DM-dominated dynamics. In the RAR, this transition corresponds to the regime in which the baryonic and dark contributions to the total acceleration become comparable. Indeed,

from Eqs. (115) and (117), if:

$$\frac{g_{\text{bar}}}{g_{\dagger}} = \mathcal{O}(1), \quad (125)$$

then the interpolation functions are themselves of order unity, and we have:

$$g_{\text{bar}} \sim g_{\text{DM}} \sim g_{\dagger}, \quad (126)$$

up to numerical coefficients of order one.

In the framework developed here, the halo-side radial domain most relevant for comparison with the RAR is the baryon-halo interface region, where the condensate is most sensitive to baryonic perturbations and where the lowest excited modes attain their largest radial support. As discussed in Sect. V G, this region is expected to lie at radii of order $r \sim r_c$. It is therefore natural to regard $g_{\text{DM}}(r \sim r_c)$ as the characteristic halo-side acceleration entering the phenomenological comparison. This should not, however, be interpreted as implying that $g_{\text{DM}}(r_c)$ must coincide with $g_{\dagger}$ in every galaxy. The condition that baryonic and dark contributions be comparable characterizes only the crossover portion of the universal RAR, and an individual galaxy need not probe that regime near $r \sim r_c$, or even within the observed radial range at all. The more robust statement is therefore that the intrinsic halo scale $g_{\star}$ is naturally of the same order as the empirical scale $g_{\dagger}$, whereas the local interface value $g_{\text{DM}}(r \sim r_c) \sim \alpha(\zeta) g_{\star}$ is determined by the halo profile and need not itself equal $g_{\dagger}$. Only in galaxies for which the baryon-DM transition occurs near $r \sim r_c$ does one expect the DM acceleration in that radial domain to be of order $g_{\dagger}$. In systems sampling a different portion of the universal RAR curve, the local relation is less direct, even though the same intrinsic halo scale $g_{\star}$ continues to control the overall order of magnitude.

a. *Numerical sanity check.* For reference:

$$g_{\star} = \frac{GM_h}{r_c^2} \sim 1.4 \times 10^{-10} \left(\frac{M_h}{10^9 M_{\odot}} \right) \left(\frac{1 \text{ kpc}}{r_c} \right)^2 \text{ ms}^{-2}, \quad (127)$$

which is naturally of the same order as the empirical SPARC value:

$$g_{\dagger} \sim (1-2) \times 10^{-10} \text{ ms}^{-2}. \quad (128)$$

This agreement should be interpreted as a consistency check of the scaling picture rather than as a precision fit. In particular, the locally sampled acceleration near $r \sim r_c$ is smaller than $g_{\star}$ by the profile factor $\alpha(\zeta)$, and may be further shifted by baryonic geometry and by small admixtures of excited condensate modes. The robust statement is therefore not that $g_{\text{DM}}(r_c)$ universally equals $g_{\dagger}$, but rather that the intrinsic halo scale $g_{\star}$ is naturally of the same order as the empirical RAR scale.

b. *Why dwarf galaxies naturally satisfy $g_{\star} \sim g_{\dagger}$.* The near coincidence between $g_{\star}$ and $g_{\dagger}$ is particularly

natural in dwarf galaxies. In these systems, baryons remain dynamically subdominant over most of the radial range, including around $r \sim r_c$, where one typically has $g_{\text{bar}}(r_c) \ll g_{\dagger}$. The point $r \sim r_c$ therefore already lies on the low-acceleration, DM-dominated branch of the RAR, rather than at the baryon-DM crossover itself. In this regime, Eq. (119) implies

$$g_{\text{DM}}(r_c) \sim \sqrt{g_{\dagger} g_{\text{bar}}(r_c)}. \quad (129)$$

On the other hand, within the AdS-Bose-Einstein condensate halo model, $g_{\text{DM}}(r_c) \sim \alpha(\zeta) g_{\star}$, with $\alpha(\zeta) \sim 0.18$ for $\zeta = 2$. Equating the two estimates gives:

$$g_{\star} \sim \frac{1}{\alpha(\zeta)} \sqrt{g_{\dagger} g_{\text{bar}}(r_c)}. \quad (130)$$

This shows why dwarf galaxies can naturally satisfy $g_{\star} \sim g_{\dagger}$ without requiring $g_{\text{DM}}(r_c)$ itself to be equal to $g_{\dagger}$. If $g_{\text{bar}}(r_c)$ lies one to two orders of magnitude below $g_{\dagger}$, then the geometric mean $\sqrt{g_{\dagger} g_{\text{bar}}(r_c)}$ lies only a factor of a few below $g_{\dagger}$; dividing by $\alpha(\zeta) \sim 0.18$ brings the inferred $g_{\star}$ back to the same order of magnitude as $g_{\dagger}$. Thus, dwarf galaxies do not generally satisfy $g_{\text{DM}}(r_c) \sim g_{\dagger}$; rather, they support the weaker but more robust conclusion that the intrinsic halo scale $g_{\star}$ is naturally of the same order as the universal empirical scale $g_{\dagger}$, without fine-tuning of the halo parameters.

In this sense, the robust identification is between the intrinsic halo scale $g_{\star}$ and the empirical RAR scale $g_{\dagger}$, whereas the local quantity $g_{\text{DM}}(r_c)$ depends on how a given galaxy samples the universal RAR curve.

C. Scaling Condition for RAR Universality

This subsection emphasizes that the existence of a universal RAR places non-trivial constraints on any halo model with a finite total mass, i.e., such that $F(\infty, \zeta) < \infty$. The argument developed below, therefore, does not rely on the specific AdS-Bose-Einstein condensate realization, but follows more generally from the combination of a cored density profile, a finite halo mass, and the empirical universality of the RAR. As a consequence, the conclusions apply to a broad class of physically admissible halo models rather than to a particular microscopic construction.

For a given value of ζ , the circular velocity generated by the condensate can be written as (see Eqs. (76) and (71)):

$$v^2(r) = \frac{GM_h}{r_c} \frac{F(x, \zeta)}{x F(\infty, \zeta)}, \quad x = \frac{r}{r_c}. \quad (131)$$

In the quasi-flat regime, typically corresponding to $x = \mathcal{O}(1-5)$, the velocity varies only weakly with radius, and one may write (see Eq. (80)):

$$v_f^2 \sim \eta(\zeta) \frac{GM_h}{r_c}, \quad (132)$$

where $\eta(\zeta)$ is a dimensionless coefficient determined solely by the halo shape.

Empirically, the SPARC RAR implies the scaling [6, 7]:

$$g_{\dagger} \sim \frac{v_f^4}{GM_{\text{bar}}}, \quad (133)$$

which is closely related to the baryonic Tully-Fisher relation. Combining Eqs. (132) and (133), we obtain:

$$g_{\dagger} \sim \eta(\zeta)^2 G \frac{M_h^2}{r_c^2 M_{\text{bar}}}. \quad (134)$$

Equation (134) shows that reproducing a universal value of $g_{\dagger}$ across galaxies requires a correlated scaling between halo and baryonic parameters. In the present variables, the condition for RAR universality takes the form:

$$\frac{M_h^2}{r_c^2 M_{\text{bar}}} \approx \text{const.} \quad \Longleftrightarrow \quad M_h \propto r_c \sqrt{M_{\text{bar}}}. \quad (135)$$

This relation should not be interpreted as a fine-tuning requirement, but rather as a consistency condition expressing the fact that halo structure and baryonic content cannot be statistically independent if a universal RAR is to emerge. Within the AdS-Bose-Einstein condensation framework, such a correlation is naturally expected to arise from the response of the condensate to baryonic matter, mediated by the lowest excited modes and controlled by the intrinsic scale r_c .

D. Synthesis and Perspective

The analysis developed in Sects. V and VI establishes a coherent link between the internal structure of a finite-mass DM halo and the empirical RAR observed in galaxies. Starting from a minimal AdS-Bose-Einstein condensation description of the halo, characterized by a core scale r_c , a total mass M_h , and a shape parameter ζ , we have derived the corresponding mass, velocity, and acceleration profiles within a fully Newtonian framework. These profiles naturally introduce an intrinsic halo acceleration scale $g_{\star} = GM_h/r_c^2$, which governs both the inner behaviour of the halo and its asymptotic Newtonian regime.

When baryons are included, the condensate no longer remains in its spherically symmetric ground state but responds through the excitation of its lowest AdS modes. This response induces an effective coupling between baryonic and DM accelerations, leading to the appearance of an emergent acceleration scale $g_{\dagger}$ that controls the RAR. In dwarf galaxies, which predominantly probe the low-acceleration regime, the intrinsic halo scale $g_{\star}$ is directly reflected in the observed relation, explaining why these systems naturally populate the DM-dominated branch of the RAR.

The empirical universality of the RAR further implies a correlated scaling between halo and baryonic parameters.

In the present framework, this requirement takes the form $M_h \propto r_c \sqrt{M_{\text{bar}}}$ (see Eq. (135)), which expresses the fact that halo structure and baryonic content cannot be statistically independent if a galaxy-independent acceleration scale is to emerge. Within the AdS-Bose-Einstein condensation scenario, this relation is not interpreted as a fine-tuning, but rather as the natural consequence of the dynamical response of a coherent condensate to the gravitational field generated by baryonic matter.

Taken together, these results support the interpretation of the RAR as an emergent phenomenon arising from the interaction between a coherent dark-sector background and baryonic structure, rather than as an indication of a modification of the gravitational law. A first-principles derivation of the condensate's linear response to baryonic perturbations, which lies beyond the scope of the present work, would provide a decisive test of this interpretation and determine the acceleration scale $g_{\dagger}$ directly from the microscopic parameters of the model.

The discussion of Sects. V-VI has been carried out using a Newtonian description of galactic dynamics, while the microscopic organization of the dark sector was formulated in terms of an effective AdS geometry. It is therefore necessary to clarify the consistency between these two levels of description, and to show that the use of Newtonian dynamics for rotation curves remains justified despite the AdS-based spectral structure underlying the condensate.

VII. IS A NEWTONIAN TREATMENT LEGITIMATE, ALTHOUGH THE CONDENSATE LIVES IN AN ADS GEOMETRY?

At first sight, it may appear inconsistent to describe the condensate using Newtonian dynamics while its microscopic modes are defined in an effective AdS background. In reality, the two descriptions correspond to different levels of approximation, and their compatibility follows from taking the appropriate non-relativistic and weak-field limits of the underlying relativistic theory.

a. Non-relativistic/weak-field limit. Galaxy rotation curves probe orbital velocities satisfying $v \ll c$ and dynamical time scales that are much longer than the intrinsic microscopic time scales associated with the condensate modes. In this regime, the motion of baryonic tracers (stars, gas, and stellar clusters) is governed by the slow-motion, weak-field limit of the relativistic gravitational dynamics. In this limit, the equations of motion reduce to Newtonian mechanics, provided that the gravitational potential Φ satisfies:

$$\frac{|\Phi|}{c^2} \ll 1. \quad (136)$$

This condition is well satisfied on galactic scales, where typical potentials are of order $|\Phi|/c^2 \sim 10^{-6}$ or smaller. Consequently, the use of Newtonian dynamics to relate

the enclosed mass to the circular velocity remains fully justified, even if the microscopic description of the dark sector involves a relativistic field theory defined in an effective AdS geometry.

b. AdS as an effective background for the dark sector. In the present scenario, the AdS structure is introduced only to organize the normal-mode spectrum of the dark sector and to fix the condensate scale $r_c = \kappa^{-1}$ through the discrete Fronsda spectrum. It does not imply that the galactic gravitational field itself is AdS. Rather, AdS acts as an effective kinematical background governing the long-wavelength excitations of the gluonic condensate, while the spacetime relevant for galactic dynamics remains in the weak-field regime.

The motion of baryonic matter is therefore determined by the usual Newtonian potential sourced by:

$$\rho_{\text{tot}} = \rho_{\text{bar}} + \rho_{\text{DM}}, \quad (137)$$

for which $|\Phi|/c^2 \ll 1$ on galactic scales. In this sense, the AdS structure fixes the internal spectrum and correlation length of the dark component without modifying the external gravitational field experienced by baryonic tracers.

c. Newton-Hooke correspondence and the effective harmonic structure. A useful consistency check is obtained by considering the non-relativistic limit of the static AdS metric. In the weak-field approximation, the timelike component of the metric may be written as:

$$g_{tt} \sim \left(1 + \frac{2\Phi}{c^2}\right) c^2, \quad (138)$$

where Φ is the effective Newtonian potential. Comparing this expression with the AdS metric component:

$$g_{tt} = (1 + \kappa^2 r^2) c^2 \quad (\text{see Eq. (20)}), \quad (139)$$

one obtains — to leading order in the weak-field limit:

$$\Phi_{\text{AdS}}(r) \sim \frac{c^2}{2} \kappa^2 r^2. \quad (140)$$

The corresponding non-relativistic equation of motion then reads:

$$\ddot{\mathbf{r}} = -\nabla \Phi_{\text{AdS}} = -\kappa^2 c^2 \mathbf{r}, \quad (141)$$

which corresponds to a harmonic-type contribution controlled by the AdS curvature scale. This behaviour is consistent with the Newton-Hooke limit of AdS symmetry, in which the non-relativistic dynamics acquires a quadratic term reflecting the presence of a negative cosmological constant.

It is important to emphasize that this effective harmonic term should not be interpreted as a physical Newtonian trapping potential acting on baryonic matter. In the present framework, the discrete and oscillator-like structure of the spectrum discussed in Sects. II-III originates from the global geometry of AdS itself, in particular from the existence of a timelike boundary and the

associated normal-mode quantization conditions. The Newton-Hooke correspondence, therefore, provides only a macroscopic analogy for the role of curvature in the non-relativistic limit. The actual spectral discreteness responsible for the condensate structure is a genuinely relativistic and geometric effect, and does not arise from a classical harmonic potential in ordinary Newtonian gravity.

d. Newtonian gravity for rotation curves as a macroscopic closure. The comparison with observed rotation curves ultimately requires a relation between the enclosed mass and the circular velocity. In the weak-field, slow-motion regime, the orbital velocity of a test body in a quasi-spherical gravitational field satisfies:

$$v^2(r) \sim \frac{GM_{\text{tot}}(r)}{r}, \quad (142)$$

which follows from the Newtonian limit of the relativistic equations of motion. Within the present framework, the AdS structure does not alter this macroscopic relation. Its role is to determine the internal organization of the dark sector, and in particular the DM density profile $\rho_{\text{DM}}(r)$ obtained from the lowest AdS modes and their baryon-induced admixtures. Once the mass distribution is specified, Eq. (142) provides a consistent closure for the dynamics of baryonic tracers at galactic scales.

This description remains valid as long as the system stays in the weak-field, non-relativistic regime, namely, provided that:

1. The orbital velocities satisfy $v \ll c$.
2. The gravitational potential obeys $|\Phi|/c^2 \ll 1$.
3. One does not attempt to describe genuinely relativistic phenomena such as strong lensing or strong-field dynamics.

These conditions are well satisfied in galaxies, ensuring that the Newtonian relation (142) can be safely used even though the microscopic description of the dark sector involves an effective AdS geometry.

e. A more systematic formulation. A fully explicit derivation may be obtained by starting from a relativistic scalar field propagating on the effective curved background and performing a controlled non-relativistic expansion. Writing the field in the standard form

$$\varphi(t, \mathbf{x}) = e^{-im_{\text{eff}}c^2t/\hbar}\psi(t, \mathbf{x}), \quad (143)$$

and assuming that the residual field ψ varies slowly compared with the Compton time scale, the relativistic field equation reduces, at leading order in $1/c$, to a Schrödinger- or Gross-Pitaevskii-type equation defined on the spatial slices, whose metric is inherited from the effective background geometry.

The weak gravitational field generated by baryons and by the condensate then enters through a Newtonian potential satisfying a Poisson equation sourced by the total

mass density. In this formulation, the use of Newtonian dynamics for galactic rotation curves is not an ad hoc assumption, but the natural macroscopic limit of the underlying relativistic field theory describing the dark condensate coupled to gravity.

In summary, the use of Newtonian dynamics for the description of galactic rotation curves is fully consistent with an AdS-based microscopic description of the condensate. In the present framework, the role of the AdS geometry is to determine the mode structure of the dark sector and to fix the characteristic core scale of the halo, while the motion of baryonic tracers takes place in the weak-field, non-relativistic regime relevant for galactic dynamics.

Possible curvature-induced contributions may arise in the non-relativistic limit as quadratic corrections of the Newton-Hooke type. However, such terms remain parametrically small on kiloparsec scales and therefore do not affect the standard relation between the enclosed mass and the circular velocity. As a result, the usual Newtonian closure (142), which connects the density profile to the observed rotation curve, remains fully applicable even though the microscopic organization of the dark component is governed by an effective AdS geometry.

VIII. CONCLUSION AND OUTLOOK

The existence of a nearly universal acceleration scale in rotationally supported galaxies, encoded in the RAR, suggests that the structure of DM halos is governed by an intrinsic infrared property of the dark sector rather than by galaxy-by-galaxy details of formation history. In this work, we have explored the possibility that this scale originates from a coherent gluonic dark sector, in which the QCD trace anomaly provides the microscopic seed of an infrared scale, and the large-distance dynamics organizes into a positive-energy lowest-weight unitary structure. Requiring Lorentz covariance then selects the group $\text{SO}(2, 3)$, whose discrete spectrum carries a representation-theoretically protected infrared gap and an associated correlation length. This correlation length sets the macroscopic size of galactic halos and provides the basis for interpreting the observed acceleration scale as the gravitational imprint of an intrinsic infrared organization of the dark sector.

Within this framework, the infrared gluonic sector admits an effective AdS description whose lowest scalar modes form a Bose-Einstein condensate. The ground state yields a regular cored density profile with finite total mass, while the lowest excited modes provide the leading channels through which the condensate responds to baryonic perturbations. The corresponding halo mass, rotation curve, and radial acceleration follow analytically from the condensate structure and introduce the intrinsic acceleration scale $g_\star$ defined in Eq. (112).

We have shown that this scale is naturally of the same order as the empirical acceleration entering the RAR,

$g_\star \sim g_\dagger$. In this perspective, the RAR can be interpreted as the macroscopic manifestation of a coherent dark sector characterized by a spectrally protected infrared scale, rather than as evidence for a modification of gravity or as the outcome of statistical correlations between baryons and halos. The observed universality of the relation then reflects the fact that halo structure is controlled, to leading order, by a small number of intrinsic parameters fixed by the infrared spectrum.

The present work should be regarded as a structural and phenomenological proposal. A decisive step will be a first-principles calculation of the condensate response to baryonic gravity, formulated as a linear-response problem in the AdS mode basis. Such a calculation is expected to determine explicitly how the empirical scale $g_\dagger$ emerges from the intrinsic parameters ($\varkappa, \zeta$) and from the mixing between the ground state and the lowest excited modes, and would therefore provide a quantitative test of the mechanism proposed here.

Further progress also requires clarifying the relation between the microscopic gluonic sector and its effective infrared description, as well as the cosmological conditions under which a coherent colour-singlet component can survive from the confinement epoch to galactic times. On the phenomenological side, the model should be confronted systematically with high-quality rotation curves, the SPARC sample, and dwarf-galaxy observations, including possible signatures of dipolar response, breathing modes, and other condensate-specific environmental effects.

More generally, the present scenario suggests that the dark component may behave as a coherent medium whose large-scale properties are fixed by an underlying infrared spectral organization. In such a picture, cored halos, finite-mass profiles, and the emergence of a nearly universal acceleration scale are not independent empirical facts, but different manifestations of the same underlying structure. Whether this interpretation can be elevated to a predictive theory will depend on the success of the linear-response program outlined above, but the results obtained here indicate that the observed galactic acceleration scale may indeed be tracing a fundamental infrared property of the dark sector.

Viewed from this perspective, it is instructive to compare the present framework with other DM scenarios in which large-scale coherence also emerges as a fundamental ingredient.

a. Comparison with ultralight-boson scenarios. Models based on axions or other ultralight bosons, often referred to as fuzzy or wave DM [35–37], provide a natural point of comparison with the present framework. In those approaches, coherence arises from an extremely small particle mass, leading to macroscopic de Broglie wavelengths and dynamics governed by the Schrödinger-Poisson system. In the present scenario, by contrast, coherence is not tied to the introduction of an ultralight elementary particle but to the existence of a spectrally protected infrared structure in a colour-singlet gluonic

sector. The characteristic halo scale is therefore set by the correlation length associated with the lowest-weight spectrum, rather than by a de Broglie wavelength.

Although both classes of models can produce cored density profiles and wave-like phenomena, they attribute these features to different physical mechanisms. In particular, the connection to the RAR is indirect in ultralight-boson models, whereas in the present framework, the appearance of a characteristic acceleration scale follows directly from the intrinsic parameters of the condensate and from its response to baryonic gravity. Distinguishing between these possibilities will likely require going beyond static rotation curves and studying time-dependent, environmental, and strongly non-spherical effects in the dark component.

b. Large-radius regime and weak-lensing extensions of the RAR. It is also useful to comment on recent weak-lensing analyses (see, for instance, Refs. [38, 39]) that extend the radial acceleration relation down to accelerations as low as $g \sim 10^{-15} \text{ ms}^{-2}$, corresponding to projected distances of order Mpc from galaxy centres. These studies report a smooth continuation of the SPARC relation into the very low-acceleration regime and have been interpreted as favouring an effective outer behaviour close to that produced by a density profile of the form $\rho(r) \propto r^{-2}$.

At first sight, this may seem in tension with the class of finite-mass halo profiles considered here. In particular, for the phenomenologically relevant case $\zeta = 2$, the density falls off as $\rho(r) \sim r^{-4}$ at large radii. However, the weak-lensing regime probes distances at which the notion of a strictly isolated halo becomes less clear. In the present framework, each halo formally extends to infinity while still carrying a finite total mass, so that at sufficiently large distances the local DM density should be viewed not as the asymptotic tail of a single halo, but as the superposition of contributions from many halos and from the surrounding large-scale environment.

From this viewpoint, the effective behaviour inferred from weak lensing need not coincide with the intrinsic outer profile of an individual condensate halo. Rather, it may reflect a collective background generated by the overlap of many finite-mass halos. In particular, if halo centres are distributed throughout space, the cumulative contribution of distant halos can be substantially shallower than the r^{-4} tail of each individual profile, thereby producing an effective large-radius behaviour closer to that inferred from weak-lensing analyses. The weak-lensing extension of the RAR would then probe the onset of this collective regime, rather than the genuine asymptotic profile of a single isolated halo.

In this sense, the apparent tension is not necessarily a contradiction, but may instead indicate that the very low-acceleration, megaparsec-scale regime is sensitive to environmental superposition effects that lie beyond the single-halo description developed in the present work. A more complete treatment of this regime would require embedding the finite-mass condensate halo in a popu-

lation of neighbouring halos and in the corresponding large-scale dark-sector background.

c. Cosmological origin of the infrared scale. Beyond these large-scale environmental effects, a further open question concerns the origin of the infrared scale itself. In the present analysis, the correlation length r_c (equivalently, the frequency scale $\omega_{\text{AdS}} = c/r_c$) has been treated as an intrinsic property of the dark sector, inferred from galactic phenomenology and tied to the lowest-weight structure of the underlying $\text{SO}(2,3)$ representation. It is natural, however, to ask whether this scale may originate from an earlier cosmological epoch. In particular, one may envisage that it emerges from a primordial phase of the dark sector, possibly associated with a high-energy transition such as the QCD epoch, and is subsequently transported to late times through cosmic expansion. In such a picture, a microscopic frequency scale at formation

could be redshifted into the present-day value of ω_{AdS} , thereby linking the macroscopic size of galactic halos to a primordial infrared imprint. While simple scaling arguments suggest that relations between the characteristic energy, temperature, and size of the condensate may arise along these lines, a consistent dynamical treatment of the formation and evolution of the condensate lies beyond the scope of the present work and will be addressed elsewhere.

Acknowledgements

Hamed Pejhan is supported by the National Science Fund, Ministry of Education and Science of Bulgaria, under contract KP-06-N92/2.

-
- [1] S. L. Adler, *Einstein gravity as a symmetry-breaking effect in quantum field theory*, Rev. Mod. Phys., **54**, 729-766 (1982)
 - [2] G. Cohen-Tannoudji, *Lambda, the fifth foundational constant considered by Einstein*, Metrologia, **55**, 486 (2018)
 - [3] J.-P. Gazeau, *Mass in de Sitter and Anti-de Sitter Universes with Regard to Dark Matter*, Universe, **6**, 66 (2020).
 - [4] G. Cohen-Tannoudji and J.-P. Gazeau, *Cold DM: A Gluonic Bose-Einstein Condensate in Anti-de Sitter Space Time*, Universe, **7**, 402 (2021).
 - [5] G. Cohen-Tannoudji and J.-P. Gazeau, *DM as a QCD effect in an Anti-de Sitter Geometry: Cosmogonic Implications of de Sitter, Anti-de Sitter and Poincaré Symmetries*, SciPost Phys. Proc., **14**, 004 (2023).
 - [6] S. S. McGaugh, F. Lelli, and J. M. Schombert, *Radial Acceleration Relation in Rotationally Supported Galaxies*, Phys. Rev. Lett., **117**(20), 201101 (2016).
 - [7] F. Lelli, S. S. McGaugh, J. M. Schombert, *SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves*, Astron. J., **152**, 157 (2016).
 - [8] F. Lelli, S.S. McGaugh, J. M. Schombert, and M. S. Pawlowski, *One Law to Rule Them All: the Radial Acceleration Relation of Galaxies*, Astrophys. J., **836**(2), 152 (2017).
 - [9] A. D. Ludlow, et al., *Mass-Discrepancy Acceleration Relation: a Natural Outcome of Galaxy Formation in Cold Dark Matter Halos*, Phys. Rev. Lett., **118**(16), 161103 (2017).
 - [10] B. W. Keller and J. W. Wadsley, *Λ CDM is Consistent with SPARC Radial Acceleration Relation*, Astrophys. J. Lett., **835**(1), L17 (2017).
 - [11] J. F. Navarro, et al., *The Origin of the Mass Discrepancy-Acceleration Relation in Λ CDM*, Mon. Not. R. Astron. Soc., **471**(2), 1841-1848 (2017).
 - [12] H. Desmond, *The Scatter, Residual Correlations and Curvature of the SPARC Baryonic Tully-Fisher Relation*, Mon. Not. R. Astron. Soc., **472**, L35-L39 (2017).
 - [13] B. Famaey and S. S. McGaugh, *Modified Newtonian Dynamics (MOND): Observational Phenomenology and Relativistic Extensions*, Liv. Rev. Relativ., **15**, 10 (2012).
 - [14] M. P. Júlio, J. I. Read, M. S. Pawlowski, P. Li, D. Vaz, J. Brinchmann, M. P. Rey, O. Agertz, and T. Holmes, *The Radial Acceleration Relation at the EDGE of Galaxy Formation; Testing its Universality in Low-Mass Dwarf Galaxies* A&A, **704**, A330 (2026).
 - [15] M.B. Hastings and T. Koma, *Spectral gap and exponential decay of correlations*, Commun. Math. Phys., **265**(3), 781-804 (2006).
 - [16] R. Pasechnik and M. Šumbera, *Phenomenological Review on Quark-Gluon Plasma: Concepts vs. Observations*, Universe, **3**(1), 7 (2017).
 - [17] V. F. Mukhanov, *Gravitational Instability of the Universe Filled with a Scalar Field*, JETP Lett., **41**, 493-496 (1985).
 - [18] M. Sasaki, *Large Scale Quantum Fluctuations in the Inflationary Universe*, Prog. Theor. Phys., **76**, 1036-1046, (1986).
 - [19] S. Weinberg, *Cosmology*, Oxford University Press, (2008).
 - [20] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan, and L. Senatore, *The Effective Field Theory of Inflation*, JHEP, **03**, 014, (2008).
 - [21] D. Baumann, *TASI Lectures on Inflation*, arXiv:0907.5424v2 [hep-th] (2011).
 - [22] C. Fronsda, *Elementary Particles in a Curved Space. II*, Phys. Rev. D, **10**, 589 (1974).
 - [23] C. Fronsda, *Elementary Particles in a Curved Space. IV. Massless Particles*, Phys. Rev. D, **12**(12), 3819 (1975).
 - [24] M. Enayati, J.-P. Gazeau, M. A. del Olmo, H. Pejhan, *Anti-de Sitterian “massive” elementary systems and their Minkowskian and Newton-Hooke contraction limits*, J. Math. Phys., **66**, 053501 (2025).
 - [25] M. Enayati, J.-P. Gazeau, H. Pejhan, and A. Wang, *The de Sitter (dS) Group and its Representations* (2nd edition), Springer, Cham, Switzerland (2024).
 - [26] C. J. Morningstar and M. Peardon, *Glueball Spectrum from an Anisotropic Lattice Study*, Phys. Rev. D, **60**(3), 034509 (1999).
 - [27] J.-P. Gazeau, H. Pejhan, and I. Todorov, *Massless Representations in Conformal Space and Their de Sitter Re-*

- strictions*, arXiv:2601.18433 (2026).
- [28] K. G. Begeman, A. H. Broeils, and R. H. Sanders, *Extended Rotation Curves of Spiral Galaxies: Dark Haloes and Modified Dynamics*, Mon. Not. R. Astron. Soc., **249**, 523-537 (1991).
 - [29] J. F. Navarro, C. S. Frenk, and S. D. M. White, *A Universal Density Profile From Hierarchical Hlustering*, Astrophys. J., **490**, 493-508 (1997).
 - [30] A. Burkert, *The Structure of DM Halos in Dwarf Galaxies*, Astrophys. J. Lett., **447**, L25-L28 (1995).
 - [31] J. I. Read, O. Agertz, and M. L. M. Collins, *DM Cores Sll the Way Down*, Mon. Not. R. Astron. Soc., **459**, 2573-2590 (2016).
 - [32] P. Li, F. Lelli, S. S. McGaugh, and J. M. Schombert, *A Comprehensive Catalog of Dark Matter Halo Models for SPARC Galaxies*, Astrophys. J. Suppl. Ser., **247**, 31 (2020).
 - [33] W. J. G. de Blok, *The Core-Cusp Problem*, Adv. Astron., **2010**, 789293 (2010).
 - [34] F. Lelli, S. S. McGaugh, and J. M. Schombert, *The Baryonic Tully-Fisher Relation and the Limits of Galaxy Scaling Laws*, Astron. Astrophys., **615**, A3 (2018).
 - [35] L. Hui, J. P. Ostriker, S. Tremaine, E. Witten, *Ultralight Scalars as Cosmological Dark Matter*, Phys. Rev. D, **95**, 043541 (2017).
 - [36] W. Hu, R. Barkana, and A. Gruzinov, *Fuzzy Cold Dark Matter: The Wave Properties of Ultralight Particles*, Phys. Rev. Lett., **85**, 1158-1161 (2000).
 - [37] H.-Y. Schive, T. Chiueh, and T. Broadhurst, *Cosmic Structure as the Quantum Interference of a Coherent Dark Wave*, Nature Phys., **10**, 496-499 (2014).
 - [38] M. M. Brouwer, et al. *The Weak Lensing Radial Acceleration Relation: Constraining Modified Gravity and Cold Dark Matter Theories with KiDS-1000*, Astronomy & Astrophysics, **650**, A113 (2021).
 - [39] T. Mistele, et al. *Radial Acceleration Relation of Galaxies with Joint Kinematic and Weak-Lensing Data*, JCAP, **2024.04**, 020 (2024).