

Galaxy Zoo: Star Formation Histories in the COSMOS Survey

1★ 2 2,3 3
1' 2 3 ,

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Key words: keyword1 – keyword2 – keyword3

1 INTRODUCTION

Becky’s group environment paper [Smethurst et al. \(2017\)](#).

2 DATA

2.1 Multi-wavelength data

In this study we used the K_s -selected catalog of the COSMOS/UltraVISTA field from [Muzzin et al. \(2013\)](#). The catalog contains PSF-matched photometry in 30 photometric bands covering the wavelength range $0.15\mu\text{m} \rightarrow 24\mu\text{m}$ and includes the available *GALEX* ([Martin et al. 2005](#)), CFHT/Subaru ([Capak et al. 2007](#)), UltraVISTA ([McCracken et al. 2012](#)), S-COSMOS ([Sanders et al. 2007](#)), and zCOSMOS ([Lilly et al. 2009](#)) datasets.

We required the rest-frame U - V and V - J colours which were calculated using the EAZY code ([Brammer et al. 2008](#)). EAZY determines rest-frame colors by integrating the best-fit SED through the redshifted filter curves over the appropriate wavelength range. For the U and V filters the Johnson response curves calculated in [Maíz Apellániz \(2006\)](#) were used. The J filter used the 2MASS response curve from [Skrutskie et al. \(2006\)](#).

2.2 Environment data

A continuous environmental density estimate (δ) for each galaxy was obtained from [Darvish et al. \(2015\)](#). The method used to estimate environment was weighted Voronoi Tessellation (WVT) which is an iteration on the simple Voronoi Tessellation technique ([Ebeling & Wiedenmann 1993](#); [Bernardeau & van de Weygaert 1996](#)). In simple Voronoi Tessellation, galaxies are divided into redshift planes to estimate local density. WVT uses a Monte-Carlo acceptance-rejection process to incorporate a weighted contribution from each galaxy in the density estimate. The

weight assigned to each galaxy corresponds to the likelihood of it residing in the particular redshift slice being examined. Weighting the contributions aims to reduce the impact of redshift uncertainties on the density estimates.

[Darvish et al. \(2015\)](#) showed that WVT outperforms several other density estimation methods when evaluated against known surface density profiles. Moreover, unlike other density estimators (nearest neighbor methods) WVT makes no underlying assumptions about the geometry and morphology of the structures in the density field. (However it is noted that Veronoi Tessellation can not be used to assign density estimates to galaxies near the edge of the field.) Overall [Darvish et al. \(2015\)](#) concludes that WVT is a robust density estimator.

The WVT algorithm and a comparison between several density estimation methods can be found in Sections 4.3 and 6 of [Darvish et al. \(2015\)](#) respectively. In this study galaxies were binned into low ($\log(1 + \delta) \leq -0.25$), intermediate ($-0.25 < \log(1 + \delta) < 0.25$), and high ($\log(1 + \delta) \geq 0.25$) density groups.

2.3 Galaxy Zoo Hubble Morphological classifications

Morphological classifications of galaxies were obtained from the Galaxy Zoo Hubble¹ (GZH) citizen science project ([Willett et al. 2017](#)). GZH allowed several independent visual classifications of each galaxy image by volunteers, the question flowchart for each image is shown in Figure 4 of [Willett et al. \(2017\)](#). Volunteers were shown colour-composite images of real and simulated *Hubble Space Telescope* galaxies to correct for classification redshift bias.

The GZH project provides detailed classifications of 119,849 galaxies from publicly-released *Hubble Space Telescope Legacy* programs conducted with the Advanced Camera for Surveys. In this study we used a subset of the total GZH sample corresponding to 84,954 galaxies found in

★

¹ <https://hubble.galaxyzoo.org>

the COSMOS Survey (Scoville et al. 2007; Koekemoer et al. 2007).

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.

3 METHODS

3.1 Modelling Star Formation History

We used the publically available, Markov Chain Monte Carlo based (Foreman-Mackey et al. 2013), STARPY² code to infer the star formation history (SFH) of each galaxy in our sample. Inputs of redshift and of rest-frame U - V and V - J colours, combined with the stellar population model of Bruzual & Charlot (2003), a solar metallicity, and a Chabrier IMF Chabrier (2003) produces a SFH for each galaxy. These models do not account for intrinsic dust. An explanation of how STARPY works can be found in Section 3

The SFH model used is parameterised by t_q and τ [Gyrs], where t_q is the onset time of quenching and τ characterises the rate of quenching. Increasing τ represents an exponentially slower quench. Assuming that all galaxies formed at $t = 0$ [Gyrs] with an initial burst of constant SFR (SFR_0), we write the SFH over cosmic time ($0 \leq t$ [Gyrs] ≤ 13.8) as:

$$\text{SFR}(t) = \begin{cases} \text{SFR}_0(t_q) & t \leq t_q \\ \text{SFR}_0(t_q) \exp\left[-\frac{(t-t_q)}{\tau}\right] & t > t_q \end{cases} \quad (1)$$

$$P(\theta_k) = \begin{cases} 1 & 0 \leq t_q \text{ [Gyr]} \leq 13.8 \text{ and } 0 \leq \tau \text{ [Gyr]} \leq 4 \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

4 CONCLUSIONS

ACKNOWLEDGEMENTS

REFERENCES

- Bernardeau F., van de Weygaert R., 1996, *Monthly Notices of the Royal Astronomical Society*, 279, 693
- Brammer G. B., van Dokkum P. G., Coppi P., 2008, *ApJ*, 686, 1503
- Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000
- Capak P., et al., 2007, *ApJS*, 172, 99
- Chabrier G., 2003, *PASP*, 115, 763
- Darvish B., Mobasher B., Sobral D., Scoville N., Aragon-Calvo M., 2015, *ApJ*, 805, 121
- Ebeling H., Wiedenmann G., 1993, *Phys. Rev. E*, 47, 704
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
- Koekemoer A. M., et al., 2007, *ApJS*, 172, 196
- Lilly S. J., et al., 2009, *ApJS*, 184, 218
- Maíz Apellániz J., 2006, *AJ*, 131, 1184
- Martin D. C., et al., 2005, *ApJ*, 619, L1
- McCracken H. J., et al., 2012, *A&A*, 544, A156
- Muzzin A., et al., 2013, *ApJS*, 206, 8
- Sanders D. B., et al., 2007, *ApJS*, 172, 86
- Scoville N., et al., 2007, *ApJS*, 172, 150
- Skrutskie M. F., et al., 2006, *AJ*, 131, 1163
- Smethurst R., Lintott C., Bamford S., Hart R., Kruk S., Masters K., Nichol R., Simmons B., 2017, *Monthly Notices of the Royal Astronomical Society*, 469, 3670
- Willett K. W., et al., 2017, *MNRAS*, 464, 4176

² <http://github.com/zooniverse/starpy/>