

Interstellar Extinction Towards OB Stars in the W3 Giant Molecular Cloud

Jessica Campbell
Supervised by Prof. Peter Martin




Astronomy & Astrophysics
UNIVERSITY OF TORONTO

AST1501 Final Presentation
May 17, 2017




Dust in the interstellar medium

causes *dimming* and *reddening* of background sources



Mie Theory of scattered light


can explain this with scattering and absorption off dust¹



* not actually spherical


The Interstellar Extinction Curve

describes the amount of scattering and absorption with λ



The CCM Model

characterizes extinction with a single parameter R_V



“ratio of total-to-selective extinction”


$$R_V = \frac{A_V}{E(B-V)} = \frac{A_V}{A_B - A_V}$$

$$E(B-V) = (B-V) - (B-V)_0$$


observed
colour

intrinsic
colour

Intrinsic Colours of OB Stars ~ 0 ; *can directly measure the line-of-sight extinction curve**



* caveat: IR excesses, emission/absorption lines etc. can bias extinction measurements



What is the empirical relationship between A_V and τ for de-reddening astrophysical observations in the molecular ISM via the W3 giant molecular cloud?

The search for an empirical relation *between extinction (A_V) and dust optical depth (τ)*




molecular cloud

A_V (mag)


τ

The search for an empirical relation *between extinction (A_V) and optical depth (τ)*




The search for an empirical relation

between extinction (A_V) and optical depth (τ)





The search for an empirical relation

between extinction (A_V) and optical depth (τ)



The search for an empirical relation *between extinction (A_V) and optical depth (τ)*







Expanding upon the OB stellar population for improving A_V/τ

STAR FORMATION IN W3 AND W4: DISCOVERY OF 135 POSSIBLY EMBEDDED NEAR-INFRARED STARS

DEBRA MELOY ELMEGREEN

The Mount Wilson and Las Campanas Observatories, Carnegie Institution of Washington


Received 1980 February 19; accepted 1980 March 26




Bright Infrared Stars (BIRS)




2MASS ($J-H$) vs ($H-K$) colour-colour diagram





M2II red supergiant



References

1. Mie, G. 1908, *Annalen der Physik*, 330, 377
2. Hollenbach, D.J., Werner, M.W., & Salpeter, E.E. 1971, *ApJ*, 163, 165
3. Kiminki, M.M., Kim, J.S., Bagley, M.B., Sherry, W.H., & Rieke, G.H. 2015, *ApJ*, 813, 42
4. Draine, B. 2003, *Annual Review of Astronomy & Astrophysics*, 41, 241
5. Cardelli, J.A., Clayton, G.C., & Mathis, J.S. 1989, *ApJ*, 345, 245
6. Bohlin, R.C., Savage, B.D., & Drake, J.F. 1978, *ApJ*, 224, 132