

GENERAL QUALIFYING EXAM SOLUTIONS: APPENDIX

Jessica Campbell, Dunlap Institute for Astronomy & Astrophysics (UofT)

Contents

1	Appendix	2
1.1	Acronyms	2
1.2	Variables	5
1.3	Useful Values	12
1.4	Equations	15
1.5	General Notes	29

1 Appendix

1.1 Acronyms

AGB: asymptotic giant branch
AGN: active galactic nuclei
AU: astronomical unit
BAOs: baryonic acoustic oscillations
BB: Big Bang or blackbody
BCD: blue compact dwarf
BBN: Big Bang nucleosynthesis
bf: bound-free
BH: black hole
BJ: Balmer jump
CCD: charged couple device
CCD: colour-colour diagram
CCM: Cardelli-Clayton-Mathis
CIB: Cosmic Infrared Background
CMB: Cosmic Microwave Background
CMD: colour-magnitude diagram
CNB: Cosmic Neutrino Background
CNM: cold neutral medium
CNO: carbon-nitrogen-oxygen
COBE: COsmic Background Explorer
CDM: cold dark matter
cE: compact elliptical
CIB: cosmic infrared background
CMB: cosmic microwave background
CR: cosmic ray
CRB: cosmic radio background
CGB: cosmic gamma-ray background
CUVOB: cosmic ultraviolet/optical background
CXB: cosmic x-ray background
DDA: discrete dipole approximation
DE: dark energy
dE: dwarf elliptical
DIBS: diffuse interstellar bands
DM: dark matter
DM: distance modulus
dSph: dwarf spheroidal
E: elliptical
E-AGB: early asymptotic giant branch
EM: electromagnetic
EOM: equation of motion
EoR: epoch of reionization
EOS: equation of state
ERE: extended red emission
FWHM: full-width at half-maximum
ff: free-free
FIR: far infrared
GC: galactic center
GC: globular cluster
gE: giant elliptical
GMC: giant molecular cloud
GP: Gunn-Peterson
GUT: grand unified theory
HB: horizontal branch
HR: Hertzsprung-Russel
HRD: Hertzsprung-Russel diagram
HST: Hubble Space Telescope
ICM: intracluster medium

IGM: intergalactic medium
IMF: initial mass function
IR: infrared
Irr: irregular
ISM: interstellar medium
KH: Kelvin-Helmholtz
LAE: Lyman-alpha emitter
LAT: linear adiabatic theory
LBG: Lyman break galaxy
LG: Local Group
LMC: Large Magellanic Cloud
LNAT: linear non-adiabatic theory
LTE: local thermodynamic equilibrium
MACHO: massive compact halo objects
MMR: mass-magnitude relation
MRI: magneto-rotational instability
MW: Milky Way
MWG: Milky Way Galaxy
NLTE: non-local thermodynamic equilibrium
NS: neutron star
ONC: Orion Nebula cluster
PAH: polycyclic aromatic hydrocarbon
pc: parsec
PDMF: present-day mass function
PDF: probability distribution function
PDR: photodissociation region
PGM: pre-galactic medium
PL: period-luminosity
PN: planetary nebula
PSF point-spread function
QFT: quantum field theory
QSO: quasi-stellar object
RGB: red giant branch
RJ: Rayleigh-Jeans
RT: radiative transfer
RTE: radiative transfer equation
S: spiral
SB: barred spiral
SDSS: Sloan Digital Sky Survey
SED: spectral energy distribution
SGB: sub giant branch
SF: star formation
SFR: star formation rate
SFRD: star formation rate density
SGB: subgiant branch
SMC: Small Magellanic Cloud
SMBH: supermassive black hole
SMG: sub-millimeter galaxy
S/N: signal-to-noise
SNO: Sudbury Neutrino Observatory
SNR: signal-to-noise ratio
SNU: solar neutrino unit
STMD: spectral type - magnitude diagram
SZ: Sunyaev-Zeldovich
SZE: Sunyaev-Zeldovich effect
TE: thermal equilibrium
TE: thermodynamic equilibrium
TO: turnoff
TRGB: red giant branch tip
VLBI: Very Long Baseline Interferometry
WD: white dwarf

WIM: warm ionized medium
WIMP: weakly interacting massive particle
WNM: warm neutral medium
ZAMS: zero age main sequence
 Λ CDM: Λ cold dark matter
2dFGRS: 2 degree Field Galactic Redshift Survey

1.2 Variables

Absolute magnitude: M [mag]
Absorption coefficient: α_ν [cm^{-1}]
Absorption cross section: $\sigma_{\ell u}(\nu)$ [cm^2]
Absorption cross section (intrinsic): σ_ν^{intr} [cm^2]
Absorption line profile: ϕ_ν [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
Absorption line profile (intrinsic): ϕ_ν^{intr} [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
Absorption rate: $\left(\frac{dn_u}{dt}\right)_{\ell \rightarrow u}$ [s^{-1}]
Albedo: ω [dimensionless]
Ampere: A [unit]
Angström: \AA [unit]
Angular diameter distance: d_A [pc]
Angular momentum per unit mass: \vec{L} [$\text{m}^2 \text{s}^{-1} \text{kg}^{-1}$]
Angular size: θ [rad]
Angular velocity: Ω [$\text{km s}^{-1} \text{kpc}^{-1}$]
Angular velocity (Solar value): Ω_0 [rad s^{-1}]
Apocenter: r_2 [AU]
Apparent magnitude: m [mag]
Atomic number: Z [dimensionless]
Attenuation: see *extinction*
Auxiliary magnetic field: \vec{H} [A m^{-1}]
Azimuthal angle: ψ [rad]
Azimuthal period: T_ψ [yr]
Balmer jump: BJ [$\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1} \text{sr}^{-1}$]
Baryon fraction: f_b [dimensionless]
Baryon temperature: T_b [K]
Binding energy of deuterium: E_D [MeV]
Binding energy of helium: E_{He} [MeV]
Binding energy of hydrogen: E_{H} [MeV]
Black hole mass: M_\bullet [M_\odot]
Black hole radius of influence: r_{BH} [pc]
Black hole radius of influence (angular): r_{BH} [arcsec]
Blackbody function: $B_\nu(T)$ [$\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1} \text{sr}^{-1}$]
Bohr magneton: μ_B [erg G^{-1}]
Boltzmann constant: k_B [$\text{m}^2 \text{kg s}^{-2} \text{K}^{-1}$]
Bonnor-Ebert mass: M_{BE} [M_\odot]
Broadening parameter: b [km s^{-1}]
Brightness temperature: T_b [K]
Brightness temperature of radio source: T_r [K]
Central temperature: T_c [K]
Chandrasekhar mass: M_C [M_\odot]
Charge: q [C]
CMB power spectrum amplitude: C_ℓ [μK^2]
CMB temperature: T_{CMB} [K]
CMB temperature fluctuations: $\Delta T/T$ [dimensionless]
CNB temperature: T_{CNB} [K]
Coefficient of conduction: k_{rad} [$\text{erg cm}^{-1} \text{K}^{-1} \text{s}^{-1}$]
Collisional coupling for species i : x_c^i [dimensionless]
Collisional excitation rate: C_{10} [s^{-1}]
Colour excess ($B - V$): $E(B - V)$ [mag]
Comoving sound horizon: $r_{\text{H,com}}$ [Mpc]
Compton y -parameter: y [dimensionless]
Cooling rate per unit mass: Λ [$\text{J kg}^{-1} \text{s}^{-1}$]
Cooling functions: Λ [$\text{J cm}^3 \text{s}^{-1}$]
Core density: ρ_c [g cm^{-3}]
Core mass: M_c [M_\odot]
Core radius: R_c [pc]
Correlation function: $C(\theta)$ [dimensionless]
Cosmological constant: Λ [m^{-2}]

Coulomb: C [unit]
 Coupling coefficient of collisions: x_c [dimensionless]
 Coupling coefficient of Ly α scattering: x_α [dimensionless]
 Coupling coefficient (total): x_{tot} [dimensionless]
 Critical density: ρ_c [g cm $^{-3}$]
 Critical density: ρ_{crit} [g cm $^{-3}$]
 Cross section: σ [cm 2]
 Cross section (absorption): C_{abs} [cm 2]
 Cross section (extinction): C_{ext} [cm 2]
 Cross section (radiation pressure): C_{pr} [cm 2]
 Cross section (scattering): C_{sca} [cm 2]
 Crossing timescale: t_{cross} [yr]
 Curvature term: $S_\kappa(r)$ [Mpc]
 Dark energy density: ρ_Λ [g cm $^{-3}$]
 Deceleration parameter: q_0 [dimensionless]
 Degeneracy: *see statistical weight*
 Density: ρ [kg m $^{-3}$ or g cm $^{-3}$]
 Density (Solar value): ρ_\odot [g cm $^{-3}$]
 Density contrast: $\delta(\mathbf{x}, t)$ [dimensionless]
 Density of baryons: ρ_b [m $^{-3}$]
 Density of dark energy: ρ_Λ [kg m $^{-3}$]
 Density of neutrinos: ρ_ν [m $^{-3}$]
 Density of neutrons: ρ_b [m $^{-3}$]
 Density of protons: ρ_b [m $^{-3}$]
 Density of vacuum: ρ_{vac} [eV m $^{-3}$]
 Density parameter: Ω_i [dimensionless]
 Density parameter of baryonic matter: Ω_b [dimensionless]
 Density parameter of curvature: Ω_κ [dimensionless]
 Density parameter of dark energy: Ω_Λ [dimensionless]
 Density parameter of dark matter: Ω_c [dimensionless]
 Density parameter of matter: Ω_m [dimensionless]
 Density parameter of neutrinos: Ω_ν [dimensionless]
 Density parameter of radiation: Ω_r [dimensionless]
 Density perturbation field: $\delta(\mathbf{x}, t)$ [dimensionless]
 Density PDF: $p(\rho)$ [???]
 Deuterium: D [element]
 Diffusion coefficient: D [s $^{-1}$]
 Diffusive flux: \vec{j} [???]
 Dimensionless density: θ [dimensionless]
 Dimensionless radius: ξ [dimensionless]
 Dipole moment: \vec{d} [C m]
 Disk scale length: h_R [Mpc]
 Distance: d [m]
 Distance modulus: $m - M$ [mag]
 Dust abundance (scaled to Solar neighborhood value): Z'_d [???]
 Dust optical depth: τ_d [dimensionless]
 Eccentricity: e [dimensionless]
 Eddington luminosity: L_E [J s $^{-1}$]
 Effective radius of dust grain: a_{eff} [μ m]
 Efficiency factor (absorption): Q_{abs} [cm $^{-2}$]
 Efficiency factor (extinction): Q_{ext} [cm $^{-2}$]
 Efficiency factor (scattering): Q_{sca} [cm $^{-2}$]
 Effective number of neutrino families: N_{eff} [dimensionless]
 Effective radius: R_e [Mpc]
 Einstein A coefficient (spontaneous emission): A_{ul} [s $^{-1}$]
 Einstein B coefficient (spontaneous absorption): B_{lu} [s $^{-1}$]
 Einstein B coefficient (stimulated emission): B_{ul} [s $^{-1}$]
 Electric field: $\vec{E} = \text{N C}^{-1}$
 Electric polarizability: $\alpha = \text{C m}^2 \text{V}^{-1}$
 Electrical conductivity: $\sigma = \text{S m}^{-1}$

Electron: e^- [particle]
Electronvolt: eV [unit]
Electron optical depth: τ_e [dimensionless]
Electroweak age: t_{ew} [s]
Electroweak energy: E_{ew} [TeV]
Electroweak temperature: T_{ew} [K]
Electron radius (classical): $r_{e,0}$ [cm]
Elliptical D_n : D_n [pc]
Ellipticity: ϵ [dimensionless]
Emission coefficient: j_ν [cm^{-1}]
Emissivity index: β [dimensionless]
Energy: E [eV or J]
Energy (lower state): E_ℓ [eV]
Energy (upper state): E_u [eV]
Energy density: *see density parameter*
Energy of CMB photons: E_{CMB} [eV]
Energy of photons: E_γ [eV]
Escape velocity: v_{esc} [km s^{-1}]
Excitation temperature: T_{ex} [K]
Extinction: A_ν or A_λ [mag]
Extinction coefficient: *see extinction*
Extinction coefficient for the MWG: $A_{\text{V,MWG}}$ [mag]
Farad: F [unit]
Flux: F [$\text{erg s}^{-1} \text{cm}^{-2}$]
Flux (conductive): F_{cd} [$\text{erg s}^{-1} \text{cm}^{-2}$]
Flux (convective): F_{con} [$\text{erg s}^{-1} \text{cm}^{-2}$]
Flux (radiative): F_{rad} [$\text{erg s}^{-1} \text{cm}^{-2}$]
Flux (spectral): F_ν [$\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$]
Free-fall time: τ_{ff} [yr]
Free-fall velocity: v_{ff} [km s^{-1}]
Frequency: ν [Hz]
Fourier transform: $P(k)$ [function]
Fractional ionization: x_x [dimensionless]
FUV intensity: χ_{FUV} [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
Gas surface mass density: Σ_{gas} $\text{M}_\odot \text{pc}^{-2}$
Galactic latitude: b [deg]
Galactic longitude: ℓ [deg]
Gas temperature: T_g [K]
Gaunt factor: g [dimensionless]
Gravitational acceleration (norm): g [m s^{-2}]
Gravitational acceleration (vector): \vec{g} [m s^{-2}]
Gravitational potential: $\Phi(r)$ [J kg^{-1}]
Guiding center radius: R_g [AU]
GUT age: t_{GUT} [s]
GUT energy: E_{GUT} [TeV]
GUT temperature: T_{GUT} [K]
Gravitational constant: G [$\text{m}^3 \text{kg}^{-1} \text{s}^{-2}$]
Growth factor: $D_+(t)$ [dimensionless]
Gunn-Peterson optical depth: τ_{GP} [dimensionless]
Gunn-Peterson oscillator strength: f_α [dimensionless]
Hamiltonian per unit mass: H [J kg^{-1}]
Harrison-Zeldovich power spectrum: $P(k)$ [μK^2]
Hayashi temperature: T_H [K]
Heating rate per unit mass: Γ [$\text{J kg}^{-1} \text{s}^{-1}$]
Helium: He [element]
Helium mass fraction: Y [dimensionless]
Henry: H [unit]
Horizon angular size: θ_{hor} [rad]
Horizon angular size (at recombination): $\theta_{\text{hor,rec}}$ [rad]
Horizon distance: d_{hor} [Mpc]
Hubble constant: H_0 [$\text{km s}^{-1} \text{Mpc}$]

Hubble distance: d_H [Mpc]
 Hubble time: t_H [Gyr]
 Hydrogen (atomic): H [element]
 Hydrogen (molecular): H_2 [element]
 IMF: $\xi(m)$ [dimensionless]
 IMF peak mass: M_{peak} [M_\odot]
 Inflationary period: t_{inflate} [s]
 Initial mass doubling time: (M_0/\dot{M}_0) [yr]
 Intensity: I [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
 Isochrone potential: $\Phi(r)$ [J]
 Jeans mass: M_J [M_\odot]
 Jeans length: λ_J [m]
 Jeans number: n_J [dimensionless]
 Jeans wavenumber: κ_J [m^{-1}]
 Joule: J [unit]
 Kelvin-Helmholtz timescale: t_{KH} [yr]
 Kinetic energy: E_{kin} [J]
 Kinetic temperature: T_k [K]
 Lagrangian per unit mass: \mathcal{L} [J]
 Lagrangian per unit mass: \mathcal{L} [J kg^{-1}]
 Line profile: $\phi(\nu)$ [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
 Line profile (intrinsic): ϕ_ν^{intr} [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
 Lithium: Li [element]
 Lorentz line profile: ϕ_ν [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
 Luminosity: L [erg s^{-1}]
 Luminosity (envelope base): L_B [erg s^{-1}]
 Luminosity (Faber-Jackson): L_{FJ} [erg s^{-1}]
 Luminosity (nuclear): L_N [erg s^{-1}]
 Luminosity (surface): L_S [erg s^{-1}]
 Luminosity (Tully-Fisher): L_{TF} [erg s^{-1}]
 Luminosity distance: d_L [Mpc]
 Luminosity of the Sun: L_\odot [W]
 Luminosity of type Ia SN (at peak): L_{Ia} [L_\odot]
 Luminosity distance: d_L [Mpc]
 Lyman- α rest wavelength: λ_α [\AA]
 Lyman- α temperature: T_α [K]
 Magnetic field: \vec{B} [G]
 Magnetic monopole density: n_M [m^{-3}]
 Magnetic monopole mass: m_M [kg]
 Magnetic permeability: μ [H m^{-1}]
 Magnitude (absolute): M [mag]
 Magnitude (apparent): m [mag]
 Mass: m or M [kg]
 Mass (luminous matter): M_{lum} [kg]
 Mass absorption coefficient: κ_ν [$\text{cm}^2 \text{g}^{-1}$]
 Mass accretion rate index: η [dimensionless]
 Mass number: A [dimensionless]
 Mass of the stellar population: M_* [kg]
 Mass of the Sun: M_\odot [kg]
 Mass surface density: Σ [$M_\odot \text{pc}^{-2}$]
 Mean free path: ℓ [cm]
 Mean free path (Solar value): $\ell_{\text{mfp},\odot}$ [cm]
 Mean molecular weight: μ [dimensionless]
 Megaelectronvolt: MeV [unit]
 Metallicity: Z [dimensionless]
 Mixing length: ℓ_m [cm]
 Moment of inertia: I [kg m^2]
 Muller matrix: S_{ij} [μ dimensionless]
 Multipole moment: C_ℓ [μK^2]
 Neutral fraction of hydrogen: x_{HI} [dimensionless]
 Neutron: n [particle]

Neutrino: ν [particle]
Neutrino temperature: T_ν [K]
Neutrino velocity: v_ν [km s⁻¹]
Newton: N [unit]
Number density: n [m⁻³]
Number density of baryons: n_b [m⁻³]
Number density of electrons: n_e [m⁻³]
Number density of Lyman- α forest lines: $N_\alpha(z)$ [dimensionless]
Number density of neutrons: n_n [m⁻³]
Number density of protons: n_p [m⁻³]
Oort constants: A, B [s⁻¹]
Opacity: κ_ν [cm² g⁻¹]
Opacity (conductive): κ_{cd} [cm² g⁻¹]
Opacity (radiative): κ_{rad} [cm² g⁻¹]
Opacity (Planck mean): κ_P [cm² g⁻¹]
Opacity (dust): κ_d [cm² g⁻¹]
Optical depth: τ [dimensionless]
Parallax: p [rad or °]
Parsec: pc [unit]
Pascal: P [unit]
Peculiar velocity: v_{pec} [km s⁻¹]
Pericenter: r_1 [AU]
Period: P [yr]
Permittivity (vacuum): ϵ [F m⁻¹]
Photon: γ [particle]
Photon occupancy number: n_γ [dimensionless]
Photon temperature: T_γ [K]
Photon scattering rate (ionized): Γ [s⁻¹]
Planck constant: h [m² kg s⁻¹]
Planck constant (reduced): \hbar [m² kg s⁻¹]
Planck energy: E_P [eV]
Planck function: $B_\nu(T)$ [erg s⁻¹ cm⁻² Hz⁻¹ sr⁻¹]
Planck length: ℓ_P [m]
Planck mass: m_P [kg]
Planck time: t_P [s]
Polarization fraction: p [dimensionless]
Polarization fraction (max value): p_{max} [dimensionless]
Polarized angle: χ [rad]
Polarized intensity: P [Jy]
Polytropic constant: K [dimensionless]
Polytropic index: n [dimensionless]
Positron: e^+ [particle]
Potential energy: E_{pot} [J]
Power spectrum: $P(k)$ [function]
Precession rate: Ω_p [rad yr⁻¹]
Pressure: P [P]
Pressure (radiation): P_{rad} [P]
Pressure scale height: H_p [cm]
Proper distance: d_p [Mpc]
Proper motion: μ [″ year⁻¹]
Proton: p [particle]
R parameter: R_V [dimensionless]
Radial period: T_r [yr]
Radial unit vector: \hat{e}_r [dimensionless]
Radial velocity: v_r [km s⁻¹]
Radiation density constant: a [erg cm⁻³ K⁻⁴]
Radiation energy density: u [erg cm⁻³]
Radiation energy density (specific): u_ν [erg cm⁻³ Hz⁻¹]
Radiation pressure acceleration: g_{rad} [m s⁻²]
Radiative energy transport: $\frac{\partial T}{\partial r}$ [K m⁻¹]
Radius (norm): R or r [pc]

Radius (Solar value): R_{\odot} [pc]
 Radius (vector): \vec{r} [pc]
 Radius of curvature: R_0 [Mpc]
 Radiative flux: F [erg s⁻¹ m⁻²]
 Rate of grain photoelectric heating: Γ_{PE} [erg s⁻¹]
 Redshift: z [dimensionless]
 Redshift (cosmological): z_{cos} [dimensionless]
 Redshift of last scattering: z_{ls} [dimensionless]
 Redshift of matter- Λ equality: $z_{m\Lambda}$ [dimensionless]
 Redshift of radiation-matter equality: z_{rm} [dimensionless]
 Redshift of thermal coupling: z_t [dimensionless]
 Reduced Hubble constant: h_0 [km s⁻¹ Mpc]
 Reduced Planck constant: \hbar [m² kg s⁻¹]
 Refractive index: $m(\omega)$ [dimensionless]
 Relaxation time: t_{relax} [yr]
 Rest mass of electron: m_e [kg]
 Rest mass of neutron: m_n [kg]
 Rest mass of proton: m_p [kg]
 Rotational energy: E_{rot} [J]
 Rotational velocity (at Solar position): V_0 [km s⁻¹]
 Scalar spectral index: n_s [dimensionless]
 Scale factor: a [dimensionless]
 Scale height: h_z [pc]
 Scattering rate between hydrogen atoms: k_{1-0}^{HH} [s⁻¹]
 Scattering rate between hydrogen atoms and electrons: $k_{1-0}^{e\text{H}}$ [s⁻¹]
 Scattering rate between hydrogen atoms and protons: $k_{1-0}^{p\text{H}}$ [s⁻¹]
 Schwarzschild radius: r_S [pc]
 Semi-major axis: a [Mpc]
 Siemen: S [unit]
 Semi-minor axis: b [Mpc]
 Sign of curvature: κ [dimensionless]
 Silk mass: M_s [kg]
 Size scale: ℓ [pc]
 Solar radius: R_{\odot} [kpc]
 Solid angle: Ω [sr]
 Sonic length: ℓ_s [pc]
 Sonic mass: M_s [M_⊙]
 Sound speed: c_s [m s⁻¹]
 Sound speed (isothermal): c_{s0} [m s⁻¹]
 Spherical harmonics: $Y_m(\theta, \phi)$ [dimensionless]
 Spin temperature: T_s [K]
 Specific intensity: I_{ν} [erg s⁻¹ cm⁻² Hz⁻¹ sr⁻¹]
 Speed of light: c [m s⁻¹]
 Spin de-excitation rate via collisions for species i : k_{10}^i [cm³s⁻¹]
 Spin temperature: T_s [K]
 Strömgren radius: r_s [pc]
 Star formation history: $\psi(z)$ [M_⊙ year⁻¹ Mpc⁻³]
 Star formation rate per unit area: Σ_{SFR} [M_⊙ year⁻¹ pc⁻²]
 Statistical weight: g [dimensionless]
 Stefan-Boltzmann constant: σ [W m⁻² K⁻⁴]
 Steradian: sr [unit]
 Stokes I: Q [Jy]
 Stokes Q: Q [Jy]
 Stokes U: Q [Jy]
 Stress energy tensor: $T_{\alpha\beta}$ [unit]
 Sunyaev-Zeldovich temperature fluctuation: ΔT_{SZ} [K]
 Sum of neutrino masses: Σm_{ν} [eV]
 Surface brightness: μ [mag arcsec⁻²]
 Surface brightness (central): μ_0 [mag arcsec⁻²]
 Surface brightness at effective radius R_e : μ_e [mag arcsec⁻²]

Tangential velocity: v_t [km s^{-1}]
Temperature: T [K]
Temperature gradient: ∇ [K P^{-1}]
Temperature gradient (actual): ∇ or ∇_{act} [K P^{-1}]
Temperature gradient (adiabatic): ∇_{ad} [K P^{-1}]
Temperature gradient (radiative): ∇_{rad} [K P^{-1}]
Temperature of the CMB: T_{CMB} [K]
Temperature of recombination: T_{rec} [K]
Temperature of the Sun: T_{\odot} [K]
Thermal energy: E_{th} [J]
Thermal energy per unit mass: e_{th} [J kg^{-1}]
Thomson cross section: σ_T [m^2]
Time: t [s or yr]
Time in units of the initial mass doubling time: τ [dimensionless]
Transfer function: $T(k)$ [???]
Tritium: T [element]
Two-point correlation function: $\xi(x)$ [dimensionless]
Velocity: v [m s^{-1}]
Velocity FWHM: $(\Delta v)_{\text{FWHM}}$ [m s^{-1}]
Velocity (luminous matter): v_{lum} [m s^{-1}]
Velocity (circular): v_c [m s^{-1}]
Velocity (radial): v_r [m s^{-1}]
Velocity (rotational): v_{rot} [m s^{-1}]
Velocity (tangential): v_t [m s^{-1}]
Velocity dispersion: σ_v [m s^{-1}]
Velocity dispersion on size scale ℓ : $\sigma_v(\ell)$ [m s^{-1}]
Virial parameter: α_{vir} [dimensionless]
Virial temperature: T_{vir} [K]
Visual extinction: A_V [mag]
Voigt profile: $\phi_{\nu}^{\text{Voigt}}$ [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
Volume: V [m^3]
Volume (specific): V [$\text{g}^{-1} \text{cm}^3$]
Wavelength: λ [m]
Wavelength of Balmer jump: λ_{BJ} [m]
Wavelength of Balmer jump (blueward): $\lambda_{\text{BJ,blue}}$ [m]
Wavelength of Balmer jump (redward): $\lambda_{\text{BJ,red}}$ [m]
Wavenumber (comoving): k [m^{-1}]

1.3 Useful Values

Age of the Earth: $t_{\oplus} = 4.5$ [Gyr]

Age of the Solar System: $t_{\text{SS}} = 4.6$ [Gyr]

Age of the Universe: $t_{\text{universe}} = 13.7$ [Gyr]

Angström: $\text{\AA} = 10^{-10}$ [m]

Astronomical Unit: $1 \text{ AU} = 1.5 \times 10^{11}$ [m]

Binding energy of deuterium: $E_{\text{D}} = 1.1$ [MeV]

Binding energy of helium: $E_{\text{He}} = 7$ [MeV]

Binding energy of hydrogen: $E_{\text{H}} = 13.7$ [MeV]

Chandrasekhar mass: $M_{\text{C}} = 1.4$ [M_{\odot}]

CMB peak photon wavelength: $\lambda_{\text{CMB}} = 2$ [mm]

CMB temperature: $T_{\text{CMB}} = 2.725$ [K]

CMB temperature fluctuations: $\Delta T/T \approx 10^{-5}$ [dimensionless]

CMB temperature: $T_{\text{CMB}} = 1.9$ [K]

Cosmological constant: $\Lambda = 1.11 \times 10^{-52}$ [m^{-2}]

Critical density (current): $\rho_{c,0} \approx 10^{-26}$ [kg m^{-3}]

Density (Solar value): $\bar{\rho}_{\odot} = 1.4$ [g cm^{-3}]

Density of dark energy: $\rho_{\Lambda} = 10^{-27}$ [kg m^{-3}]

Density parameter of baryons: $\Omega_b \approx 0.044$ [dimensionless]

Density parameter of CMB: $\Omega_{\text{CMB}} \approx 5 \times 10^{-5}$ [dimensionless]

Density parameter of curvature: $\Omega_{\kappa} \approx 0.0$ [dimensionless]

Density parameter of cold dark matter: $\Omega_c \approx 0.23$ [dimensionless]

Density parameter of dark energy: $\Omega_{\Lambda} \approx 0.73$ [dimensionless]

Density parameter of matter: $\Omega_m \approx 0.27$ [dimensionless]

Density parameter of neutrinos: $\Omega_{\nu} < 0.12$ [dimensionless]

Density parameter of radiation: $\Omega_r \approx 5 \times 10^{-5}$ [dimensionless]

Dipole: $\ell = 1$ [dimensionless]

Electronvolt in Joules: $1 \text{ eV} = 1.6 \times 10^{-19}$ J

Electron radius (classical): $r_{e,0} = 2.82 \times 10^{-13}$ [cm]

Electroweak age: $t_{\text{ew}} \sim 10^{-12}$ [s]

Electroweak energy: $E_{\text{ew}} \sim 1$ [TeV]

Electroweak temperature: $T_{\text{ew}} \sim 10^{16}$ [K]

Energy of CMB photons: $E_{\gamma, \text{CMB}} \sim 6 \times 10^{-4}$ [eV]

Energy of nuclei fission: $E_{\text{fission}} \sim 1$ [MeV]

Energy of nuclei ionization: $E_{\text{ion}} \sim 10$ [eV]

GUT age: $t_{\text{GUT}} \sim 10^{-36}$ [s]

GUT energy: $E_{\text{GUT}} \sim 10^{12} - 10^{13}$ [TeV]

GUT temperature: $T_{\text{GUT}} \sim 10^{28}$ [K]

Helium mass fraction: $Y = 0.25$ [dimensionless]

Hubble constant: $H_0 = 70$ [$\text{km s}^{-1} \text{ Mpc}^{-1}$]

Hubble constant (original): $H_0 \approx 500$ [$\text{km s}^{-1} \text{ Mpc}^{-1}$]

Hubble distance: $d_H \sim 0.2$ [Mpc]

Hubble time: $t_H \sim 14$ [Gyr]

IMF normalization: $\int_{m_L}^{m_U} m \xi(m) dm = 1 M_{\odot}$

Inflationary period: $t_{\text{inflate}} = 10^{-35}$ [s]

Ionization energy of atomic hydrogen: $E_{\text{ion}}^{\text{H}} = 13.6$ [eV]

Ionization energy of atomic helium: $E_{\text{ion}}^{\text{He}} = 24.6$ [eV]

Ionization energy of molecular hydrogen: $E_{\text{ion}}^{\text{H}_2} = 11.26$ [eV]

Large cosmological scale: ~ 100 [Mpc]

Legendre Polynomials: P_l

Line profile normalization: $\int_0^{\infty} \phi(\nu) d\nu = 1$ [$\text{erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$]

Luminosity of the Sun: $L_{\odot} = 3.8 \times 10^6$ [W]

Luminosity of type Ia SN (at peak): $L_{\text{Ia}} = 4 \times 10^9$ [L_{\odot}]

Lyman-alpha rest wavelength: $\lambda_{\alpha} = 1216$ [\AA]

Magnitude (absolute) of the Sun: $M = 26.8$ [dimensionless]

Magnitude (apparent) of the Sun: $m = 4.74$ [dimensionless]

Mass of an electron: $m_e = 9.109 \times 10^{-31}$ [kg]

Mass of atomic hydrogen: $m_{\text{H}} = 1.67 \times 10^{-27}$ [kg]

Mass of a neutron: $m_n = 1.67 \times 10^{-27}$ [kg]
Mass of a proton: $m_p = 1.67 \times 10^{-27}$ [kg]
Mean free path (Solar value): $\ell_{\text{mfp},\odot} = 2$ [cm]
Megaelectronvolt: $\text{MeV} = 10^{10.065}$ [K]
Metallicity of the Sun: $Z \approx 0.02$ [dimensionless]
Milky Way luminosity: $L_{\text{MWG}} = 3.6 \times 10^{10}$ [L_\odot]
Monopole: $\ell = 0$ [dimensionless]
Neutron half life: $t_n \sim 890$ [sec] or ~ 15 [min]
Neutron-proton freezeout: $\frac{n_n}{n_p} \simeq \frac{1}{7}$ [dimensionless]
Neutron-proton rest mass-energy difference: $(m_n - m_p)c^2 = 1.3$ [MeV]
Number density of baryons: $n_{b,0} = 0.22$ [m^{-3}]
Number density of CMB photons: $n_\gamma = 4.11 \times 10^8$ [m^{-3}]
Number density ratio of baryons to CMB photons: $\eta \equiv \frac{n_{b,0}}{n_\gamma} = 5 \times 10^{-10}$ [dimensionless]
Oort constant A: $A = (14.8 \pm 0.8)$ [$\text{km s}^{-1} \text{kpc}^{-1}$]
Oort constant B: $B = (-12.4 \pm 0.6)$ [$\text{km s}^{-1} \text{kpc}^{-1}$]
Parsec: $1 \text{ pc} = 206265 \text{ AU} = 3.1 \times 10^{16} \text{ m}$
Permittivity (vacuum): $\epsilon_0 = 8.85 \times 10^{-12}$ [F m^{-1}]
Photoionization energy of hydrogen: $E_{\text{ion}}^{\text{H}} = 13.6$ [eV]
Planck energy: $E_P = 10^{16}$ [TeV]
Planck length: $\ell_P = 1.6 \times 10^{-35}$ [m]
Planck mass: $M_P = 2.2 \times 10^{-8}$ [kg]
Planck temperature: $T_P = 1.4 \times 10^{32}$ [K]
Planck time: $t_P = 5.4 \times 10^{-44}$ [s]
Proper surface area: $A_p(t_0)$ [Mpc^2]
Radiation density constant: $a = 7.57 \times 10^{-15}$ [$\text{erg cm}^{-3} \text{K}^{-4}$]
Recombination age of Universe: $t_{\text{rec}} \sim 380,000$ [yr]
Recombination redshift: $z_{\text{rec}} \sim 1,100$ [dimensionless]
Reduced Hubble constant: $\frac{H_0}{100} = 0.70$ [$\text{km s}^{-1} \text{Mpc}^{-1}$]
Redshift of radiation-matter equality: $z_{rm} \approx 5399$ [dimensionless]
Redshift of matter- Λ equality: $z_{m\Lambda} \approx 0.39$ [dimensionless]
Rest energy of an deuterium: $m_D c^2 = 2.225$ [MeV]
Rest energy of an electron: $m_e c^2 = 0.511$ [MeV]
Rest energy of a neutron: $m_n c^2 = 939.6$ [MeV]
Rest energy of a proton: $m_p c^2 = 938.3$ [MeV]
Rotational velocity (at Solar position): $V_0 = 220$ [km s^{-1}]
Scalar spectral index: $n_s \approx 1$ [dimensionless]
Scale factor (current): $a_0 = a(t_0) \equiv 1$ [dimensionless]
Scalefactor of radiation-matter equality: $a_{rm} \approx 2.8 \times 10^{-4}$ [dimensionless]
Scalefactor of matter- Λ equality: $a_{m\Lambda} \approx 0.75$ [dimensionless]
Seconds in a year: $1 \text{ yr} = 3.2 \times 10^7$ [s]
Seconds in a gigayear: $1 \text{ Gyr} = 3.2 \times 10^{16}$ [s]
Sign of curvature (closed Universe): $\kappa > 0$ [dimensionless]
Sign of curvature (flat Universe): $\kappa = 0$ [dimensionless]
Sign of curvature (open Universe): $\kappa < 0$ [dimensionless]
Solar absolute magnitude: $M_\odot = 4.74$ [mag]
Solar apparent magnitude: $m_\odot = 26.8$ [mag]
Solar flux: $F_\odot = 1367$ [$\text{erg s}^{-1} \text{m}^{-2}$]
Solar luminosity: $L_\odot = 3.8 \times 10^{26}$ [W]
Solar mass: $M_\odot = 2.0 \times 10^{30}$ [kg]
Solar mean photon energy: $\langle E_\odot \rangle = 1.3$ [eV]
Solar radius: $R_\odot = 8.5$ [kpc]
Solar temperature: $T_\odot = 5800$ [K]
Speed of light: $c = 2.998 \times 10^8$ [m s^{-1}]
Stefan-Boltzmann constant: $\sigma = 5.67 \times 10^{-8}$ [$\text{W m}^{-2} \text{K}^{-4}$]
Supernova peak luminosity: $L_{\text{SN,max}} = 4 \times 10^9 L_\odot$ [W]
Temperature at CMB decoupling: $k_B T_{\text{CMB,dec}} = 1/4$ [eV]
Thomson cross section: $\sigma_T = 0.665 \times 10^{-24}$ [cm^2]
Time of radiation-matter equality: $t_{rm} \approx 4.7 \times 10^4$ [yr]
Time of matter- Λ equality: $t_{m\Lambda} \approx 9.8$ [Gyr]

Vacuum energy density: $\rho_{\text{vac}} = 10^{133} \text{ [eV cm}^{-3}\text{]}$

21 cm energy: $E_{21 \text{ cm}} = 5.9 \times 10^{-6} \text{ [eV]}$

21 cm rest frequency: $\nu_{21 \text{ cm}} = 1420 \text{ [MHz]}$

1.4 Equations

Absolute magnitude:

$$M \equiv -2.5 \log \left(\frac{L}{L_0} \right) \text{ [mag]}$$

$$M \equiv -2.5 \log \left(\frac{L}{78.7 L_\odot} \right) \text{ [mag]}$$

Absorption coefficient: α_ν [cm^{-1}]

$$\alpha_\nu = n\sigma_\nu = \rho\kappa_\nu \text{ [cm}^{-1}\text{]}$$

Absorption cross section (intrinsic):

$$\sigma_\nu^{\text{intr}} = \frac{\pi e^2}{m_e c^2} f_{\ell u} \phi_\nu^{\text{intr}} \text{ [cm}^2\text{]}, \quad \int \phi_\nu^{\text{intr}} d\nu = 1$$

Absorption rate:

$$\left(\frac{dn_u}{dt} \right)_{\ell \rightarrow u} = n_\ell \int \sigma_{\ell u}(\nu) c \frac{u_\nu}{h\nu} d\nu \approx n_\ell u_\nu \frac{c}{h\nu} \int \sigma_{\ell u}(\nu) d\nu \text{ [s}^{-1}\text{]}$$

$$B_{\ell u} = \frac{c}{h\nu} \int \sigma_{\ell u}(\nu) d\nu \text{ [s}^{-1}\text{]}$$

Adiabatic gas law:

$$PV^\gamma = K \text{ [P m}^3\text{]}$$

Albedo:

$$\omega \equiv \frac{C_{\text{sca}}(\lambda)}{C_{\text{abs}}(\lambda) + C_{\text{sca}}} = \frac{C_{\text{sca}}(\lambda)}{C_{\text{ext}}(\lambda)} \text{ [dimensionless]}$$

Angular diameter distance:

$$D_A = D(1+z)^{-1} \text{ [pc]}$$

Angular momentum per unit mass:

$$\vec{L} \equiv \vec{r} \times \frac{d\vec{r}}{dt} \text{ [m}^2 \text{ s}^{-1} \text{ kg}^{-1}\text{]}$$

$$\vec{L} \equiv r^2 \dot{\phi} \text{ [m}^2 \text{ s}^{-1} \text{ kg}^{-1}\text{]}$$

Angular velocity:

$$\Omega \equiv \frac{V(R)}{R} \text{ [km s}^{-1} \text{ kpc}^{-1}\text{]}$$

Apocenter:

$$r_2 = a(1+e) \text{ [AU]}$$

Apparent magnitude:

$$m \equiv -2.5 \log \left(\frac{F}{F_0} \right) \text{ [mag]}$$

$$m \equiv -2.5 \log \left(\frac{F}{2.53 \times 10^{-8} \text{ W m}^{-2}} \right) \text{ [mag]}$$

Apparent-absolute magnitude relation:

$$M = m - 5 \log_{10} \left(\frac{d_L}{10 \text{ pc}} \right) \text{ [mag]}$$

$$= m - 5 \log_{10} \left(\frac{d_L}{1 \text{ Mpc}} \right) - 25 \text{ [mag]}$$

Azimuthal angle (change):

$$\Delta\psi = 2L \int_{r_1}^{r_2} \frac{dr}{r^2 \sqrt{2[E - \Phi(r)] - \frac{L^2}{r^2}}} \text{ [rad]}$$

Azimuthal period:

$$T_\psi = \frac{2\pi}{|\Delta\phi|} T_r \text{ [yr]}$$

Balmer jump:

$$\text{BJ} \equiv I_\lambda(\lambda_{\text{BJ,blue}}) - I_\lambda(\lambda_{\text{BJ,red}}) \text{ [erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}\text{]}$$

Baryon fraction:

$$f_b \equiv \frac{\Omega_b}{\Omega_m} \text{ [dimensionless]}$$

Black hole mass-luminosity relation:

$$M_\bullet = 1.7 \times 10^9 \left(\frac{L_V}{10^{11} L_{V\odot}} \right)^{1.11} [M_\odot]$$

Black hole mass-bulge relation:

$$M_\bullet = 2.9 \times 10^8 \left(\frac{M_{\text{bulge}}}{10^{11} M_\odot} \right)^{1.05} [M_\odot]$$

$$M_\bullet \approx 3 \times 10^{-3} M_{\text{bulge}}$$

Black hole mass-velocity dispersion relation:

$$M_\bullet = 2.1 \times 10^8 \left(\frac{\sigma_v}{200 \text{ km s}^{-1}} \right)^{5.64} [M_\odot]$$

Black hole radius of influence:

$$r_{\text{BH}} = \frac{GM_\bullet}{\sigma_v^2} \sim 0.4 \left(\frac{M_\bullet}{10^6 M_\odot} \right) \left(\frac{\sigma_v}{100 \text{ km s}^{-1}} \right)^{-2} [\text{pc}]$$

Black hole radius of influence (angular):

$$\theta_{\text{BH}} = \frac{r_{\text{BH}}}{d} \sim 0.1 \left(\frac{M_\bullet}{10^6 M_\odot} \right) \left(\frac{\sigma_v}{100 \text{ km s}^{-1}} \right)^{-2} \left(\frac{d}{1 \text{ Mpc}} \right)^{-1} [\text{arcsec}]$$

Blackbody function:

$$B_\nu(T) = \frac{2h\nu^2}{c^2} \frac{1}{\exp(h\nu/k_B T) - 1} [\text{erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}]$$

Blackbody mean photon energy:

$$\langle E_{\text{bb}} \rangle = 2.7 k_B T [\text{eV}]$$

Blackbody peak photon energy:

$$E_{\text{bbpeak}} = 2.82 k_B T [\text{eV}]$$

Bohr magneton:

$$\mu_B \equiv \frac{e\hbar}{2m_e c} [\text{erg G}^{-1}]$$

Boltzmann factor:

$$\frac{n_i}{n_j} = e^{-\Delta mc^2/k_B T} [\text{dimensionless}]$$

Broadening parameter:

$$b \equiv \sqrt{2}\sigma_v [\text{km s}^{-1}]$$

Central temperature (collapsing core):

$$T_c = T_0 \left(\frac{\rho_c}{\rho_{\text{ad}}} \right)^{\gamma-1} [\text{K}]$$

CMB temperature fluctuations:

$$\frac{\delta T}{T}(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi) [\text{dimensionless}]$$

Comoving sound horizon:

$$r_{\text{H,com}}(z) = \int_0^t \frac{cdt}{a(t)} = \int_0^{(1+z)^{-1}} \frac{cda}{a^2 H(a)} [\text{Mpc}]$$

Compton y-parameter:

$$y = \int \frac{k_B T}{m_e c^2} \sigma_T n_e dl [\text{dimensionless}]$$

Cosmological constant:

$$\Lambda = \frac{3H_0^2}{c^2} \Omega_\Lambda [\text{m}^{-2}]$$

Correlation function of CMB temperature anisotropies:

$$C(\theta) = \left\langle \frac{\delta T}{T}(\hat{n}) \frac{\delta T}{T}(\hat{n}') \right\rangle_{\hat{n} \cdot \hat{n}' = \cos \theta} \quad [\text{dimensionless}]$$

$$C(\theta) = \frac{1}{4\pi} \sum_{\ell=0}^{\infty} (2\ell + 1) (C_{\ell}) P_{\ell}(\cos \theta) \quad [\text{dimensionless}]$$

Collisional coupling coefficient:

$$x_c = x_c^{\text{HH}} + x_c^{\text{eH}} + x_c^{\text{pH}}$$

$$= \frac{T_*}{A_{10} T_{\gamma}} \left[k_{1-0}^{\text{HH}}(T_k) n_{\text{H}} + k_{1-0}^{\text{eH}}(T_k) n_e + k_{1-0}^{\text{pH}}(T_k) n_p \right] \quad [\text{dimensionless}]$$

Colour excess-HI column density relation:

$$E(B - V) = 1.7 \left(\frac{N_{\text{H}}}{10^{22} \text{ atoms cm}^{-2}} \right) \quad [\text{mag}]$$

Coupling coefficient (total):

$$x_{\text{tot}} \equiv x_c + x_{\alpha} \quad [\text{dimensionless}]$$

Critical density:

$$\rho_c \equiv \frac{3H_0^2}{8\pi G} \quad [\text{g cm}^{-3}]$$

Critical density:

$$\rho_{\text{crit}} \sim \frac{c_s^2}{G \ell_s \ell} \quad [\text{g cm}^{-3}]$$

Cross section (radiation pressure):

$$C_{\text{pr}}(\lambda) \equiv C_{\text{abs}}(\lambda) + (1 - \langle \cos \theta \rangle) C_{\text{sca}}(\lambda) \quad [\text{cm}^2]$$

Crossing timescale:

$$t_{\text{cross}} = \frac{R}{v} \quad [\text{yr}]$$

Curvature term:

$$S_{\kappa}(r) = \begin{cases} R \sin(r/R), & (\kappa = +1) \\ r, & (\kappa = 0) \\ R \sinh(r/R), & (\kappa = -1) \end{cases}$$

Dark energy density:

$$\rho_{\Lambda} \equiv \frac{\Lambda}{8\pi G} \quad [\text{g cm}^{-3}]$$

Deceleration parameter:

$$q_0 = \Omega_{r,0} + \frac{1}{2} \Omega_{m,0} - \Omega_{\Lambda} \quad [\text{dimensionless}]$$

Density contrast:

$$\delta(\mathbf{x}, t) \equiv \frac{\rho(\mathbf{x}, t) - \langle \rho(t) \rangle}{\langle \rho(t) \rangle} \quad [\text{dimensionless}]$$

Density parameter:

$$\Omega_i \equiv \frac{\rho_i}{\rho_c} \quad [\text{dimensionless}]$$

Density parameter (total):

$$\Omega = \sum \Omega_i \equiv \sum \frac{\rho_i}{\rho_c} \quad [\text{dimensionless}]$$

Density parameter of curvature:

$$\Omega_{\kappa} = 1 - \Omega_0 \quad [\text{dimensionless}]$$

Density parameter of dark energy:

$$\Omega_{\Lambda} \equiv \frac{\rho_{\Lambda}}{\rho_c}$$

$$= \left(\frac{\Lambda}{8\pi G} \right) \left(\frac{8\pi G}{3H_0^2} \right)$$

$$= \frac{\Lambda}{3H_0^2} \quad [\text{dimensionless}]$$

Density parameter of matter:

$$\Omega_m(a) \equiv \frac{\rho_m}{\rho_c} a(t)^{-3} \text{ [dimensionless]}$$

Density parameter of radiation:

$$\Omega_r(a) \equiv \frac{\rho_r}{\rho_c} a(t)^{-4} \text{ [dimensionless]}$$

Density-redshift-scalefactor relation:

$$\bar{\rho}(z) = \bar{\rho}_0(1+z)^3 = \frac{\bar{\rho}_0}{a^3} \text{ [g cm}^{-3}\text{]}$$

Diffusion coefficient:

$$D = \frac{1}{3} v \ell_{\text{mfp}} \text{ [s}^{-1}\text{]}$$

Diffusive flux:

$$\vec{j} = -D \nabla n \text{ [???]}$$

Dimensionless density:

$$\theta = \left(\frac{\rho}{\rho_c} \right)^{1/n} \text{ [dimensionless]}$$

Dimensionless radius:

$$\xi \equiv \frac{r}{a} = \frac{r}{\sqrt{\frac{(n+1)K}{4\pi G} \rho_c^{(1-n)/n}}} \text{ [dimensionless]}$$

Dipole moment:

$$\vec{\mu} = q \vec{d} \text{ [C m]}$$

$$\vec{\mu} = \alpha \vec{E} \text{ [C m]}$$

$$\vec{d} = e \vec{r} \text{ [C m]}$$

Distance (moving cluster):

$$d = \frac{v_r \tan \theta}{\mu} \text{ [pc]}$$

Distance (parallax):

$$d = \left(\frac{p}{1''} \right)^{-1} \text{ [pc]}$$

Distance (photometric):

$$m - M = 5 \log \left(\frac{d}{1 \text{ pc}} \right) - 5 + A \text{ [mag]}$$

Distance (spectroscopic):

$$m_V - A_V - M_V = 5 \log \left(\frac{d}{1 \text{ pc}} \right) - 5 \text{ [mag]}$$

Distance modulus:

$$m - M \approx 43.17 - 5 \log_{10} \left(\frac{H_0}{70 \text{ km s}^{-1} \text{ Mpc}} \right) + 5 \log_{10} z + 1.086(1 - q_0)z \text{ [mag]}$$

Distance relation (low redshift):

$$d_p(t_0) \approx d_L \approx \frac{c}{H_0} z \text{ [Mpc]}$$

Distance-redshift relation (flat Universe):

$$d_L(\kappa = 0) = r(1+z) = d_p(t_0)(1+z) \text{ [Mpc]}$$

Doppler law:

$$\frac{v}{c} = \frac{\Delta \lambda}{\lambda} = \frac{\lambda_{\text{obs}} - \lambda_{\text{em}}}{\lambda_{\text{obs}}} \text{ [dimensionless]}$$

$$\frac{v}{c} = \frac{-\Delta \nu}{\nu} = \frac{\nu_{\text{em}} - \nu_{\text{obs}}}{\nu_{\text{em}}} \text{ [dimensionless]}$$

Eccentricity:

$$e = \sqrt{1 - \frac{b^2}{a^2}} \text{ [dimensionless]}$$

Eddington luminosity:

$$L_E = \frac{4\pi cGM}{\kappa} [\text{J s}^{-1}]$$
$$\frac{L_E}{L_\odot} = 1.3 \times 10^4 \frac{1}{\kappa} \frac{M}{M_\odot}$$

Effective radius of dust grain:

$$a_{\text{eff}} \equiv \left(\frac{3V}{4\pi} \right)^{1/3} [\mu\text{m}]$$

Einstein relations:

$$g_1 B_{lu} = g_2 B_{ul} [\text{s}^{-1}]$$
$$A_{ul} = \frac{2h\nu^3}{c^2} B_{ul} [\text{s}^{-1}]$$

Electric polarizability:

$$\alpha = \frac{\vec{\mu}}{\vec{E}} [\text{C m}^2 \text{V}^{-1}]$$

Electron radius (classical):

$$r_{e,0} = \frac{e^2}{m_e c^2} [\text{cm}]$$

Elliptical $D_n - \sigma$ relation:

$$I_n \left(\frac{D_n}{2} \right)^2 \pi = 2\pi I_e R_e^2 \int_0^{D_n/(2R_e)} [\text{erg s}^{-1} \text{m}^{-2}]$$
$$D_n = 2.05 \times \left(\frac{\sigma_v}{100 \text{ km s}^{-1}} \right) [\text{kpc}]$$

Efficiency factor (absorption):

$$Q_{\text{abs}} \equiv \frac{C_{\text{abs}}}{\pi a_{\text{eff}}^2} [\text{cm}^{-2}]$$

Efficiency factor (extinction):

$$Q_{\text{ext}} \equiv Q_{\text{abs}} + Q_{\text{sca}} [\text{cm}^{-2}]$$

Efficiency factor (scattering):

$$Q_{\text{sca}} \equiv \frac{C_{\text{sca}}}{\pi a_{\text{eff}}^2} [\text{cm}^{-2}]$$

Energy density:

$$\Omega \equiv \frac{\rho}{\rho_c} = nE [\text{dimensionless}]$$

Energy-redshift relation:

$$E = E_0(1+z) [\text{eV}]$$

Escape velocity:

$$v_{\text{esc}} = \sqrt{\frac{2GM}{r}} [\text{km s}^{-1}]$$

Extinction:

$$A_\nu \equiv m - m_0 [\text{mag}]$$
$$= -2.5 \log \left(\frac{I_{\nu,0}}{I_\nu} \right) [\text{mag}]$$
$$= -2.5 \log \left(\frac{F_{\nu,0}}{F_\nu} \right) [\text{mag}]$$
$$= 2.5 \log(e) \tau_\nu [\text{mag}]$$
$$= 1.086 \tau_\nu [\text{mag}]$$

Extinction coefficient for the MWG:

$$A_{V,\text{MWG}} = (3.1 \pm 0.1) E(B-V) [\text{mag}]$$

Extinction coefficient in the Solar neighborhood:

$$A_V \approx \frac{d}{1 \text{ kpc}} [\text{mag}]$$

Extinction cross section:

$$C_{\text{ext}}(\lambda) \equiv C_{\text{abs}}(\lambda) + C_{\text{sca}}(\lambda) \text{ [cm}^2\text{]}$$

Faber-Jackson relation:

$$L_{\text{FJ}} \propto \sigma_v^4 \text{ [erg s}^{-1}\text{]}$$

Farad:

$$1 \text{ F} \equiv \text{s}^4 \text{ A}^2 \text{ m}^{-2} \text{ kg}^{-1}$$

Flux:

$$F = \frac{L}{4\pi S_\kappa(r)^2} (1+z)^{-2} \text{ [erg s}^{-1} \text{ cm}^{-2}\text{]}$$

Flux (conductive):

$$\vec{F}_{\text{cd}} = -k_{\text{cd}} \nabla T \text{ [m}^2 \text{ s}^{-1}\text{]}$$

Flux (radiative):

$$F_r = -\frac{4acT^3}{3\kappa\rho} \frac{dT}{dr} \text{ [erg s}^{-1} \text{ cm}^{-2}\text{]}$$

$$F_r = \frac{L_r}{4\pi r^2} \text{ [erg s}^{-1} \text{ cm}^{-2}\text{]}$$

Flux (spectral):

$$F_\lambda(t) = \int_0^t \text{SFR}(t-t') S_{\lambda,Z(t-t')}(t') dt' \text{ [erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}\text{]}$$

Free-fall time:

$$\tau_{\text{ff}} = \sqrt{\frac{3\pi}{32G\rho_0}} = 4.4 \times 10^4 \left(\frac{n_{\text{H}}}{10^6 \text{ cm}^{-3}} \right)^{-1/2} \text{ [yr]}$$

Friedmann equation:

$$\left(\frac{\dot{a}(t)}{a(t)} \right)^2 = \left(\frac{H}{H_0} \right)^2 = \frac{8\pi G}{3} \rho - \frac{\kappa c^2}{R_0^2 a(t)^2} + \frac{\Lambda}{3}$$
$$\left(\frac{H}{H_0} \right)^2 = \frac{\Omega_{r,0}}{a^4} + \frac{\Omega_{m,0}}{a^3} + \Omega_{\Lambda,0} + \frac{1 - \Omega_0}{a^2}$$

Gravitational acceleration:

$$g(r) = -\frac{d\Phi}{dr} \text{ [m s}^{-2}\text{]}$$

Gravitational potential energy:

$$E = -\frac{GMm}{R} \text{ [J]}$$

Growth factor:

$$D_+(t) \propto \frac{H(a)}{H_0} \int_0^a \frac{da'}{[\Omega_m/a' + \Omega_\Lambda a'^2 - (\Omega_m + \Omega_\Lambda - 1)]^{3/2}} \text{ [dimensionless]}$$

Gunn-Peterson optical depth:

$$\tau_{\text{GP}} = \frac{\pi e^2}{m_e c^2} f_\alpha \lambda_\alpha H^{-1}(z) n_{\text{HI}} \text{ [dimensionless]}$$

$$\tau_{\text{GP}}(z) = 4.9 \times 10^5 \left(\frac{\Omega_m h^2}{0.13} \right)^{-1/2} \left(\frac{\Omega_b h^2}{0.02} \right) \left(\frac{1+z}{7} \right)^{3/2} \left(\frac{n_{\text{HI}}}{n_{\text{H}}} \right) \text{ [dimensionless]}$$

Hamilton's equations:

$$\dot{q}_i \equiv \frac{\partial H}{\partial p_i}; \quad \dot{p} \equiv -\frac{\partial H}{\partial q_i}$$

Harrison-Zeldovich power spectrum:

$$P(k) \propto k^{n_s-1} \text{ [}\mu\text{K}^2\text{]}$$

Helium mass fraction:

$$Y = \frac{4 \times n_n/2}{n_n + n_p} = \frac{2}{1 + n_n/n_p} \text{ [dimensionless]}$$

Henry (unit):

$$1 \text{ H} \equiv 1 \text{ kg m}^2 \text{ s}^{-2} \text{ A}^{-2}$$

Horizon angular size at recombination:

$$\theta_{H,\text{rec}} \approx \sqrt{\frac{\Omega_m}{z_{\text{rec}}}} \sim \frac{\sqrt{\Omega_m}}{30} \sim \sqrt{\Omega_m} 2 \text{ [}^\circ\text{]}$$

Hubble constant:

$$H_0(t) \equiv \frac{\dot{a}_0(t)}{a_0(t)} = \dot{a}_0(t) \text{ [km s}^{-1} \text{ Mpc]}$$

Hubble distance:

$$d_H \equiv \frac{c}{H(t)} \text{ [Mpc]}$$

Hubble time:

$$t_H \equiv ct_H = \frac{1}{H_0(t)} \text{ [Gyr]}$$

Hubble's law:

$$v = H_0 d \text{ [km s}^{-1}\text{]}$$

Hydrostatic equilibrium:

$$\frac{dP}{dr} = -\frac{GM_r \rho}{r^2} = -\rho g \text{ [P m}^{-1}\text{]}, \quad g \equiv \frac{GM_r}{r^2} \text{ [m s}^{-2}\text{]}$$

Ideal gas law:

$$P = \frac{\rho k_B T}{\mu m_H} \text{ [P]}$$

IMF peak mass:

$$M_{\text{peak}} \propto \frac{c_s^3}{\sqrt{G^3 \bar{\rho}}} \text{ [M}_\odot\text{]}$$
$$M_{\text{peak}} \propto \frac{c_s^2 \ell_s}{G} \text{ [M}_\odot\text{]}$$

IMF (Salpeter):

$$\xi(m) \propto m^{-2.35} \text{ [dimensionless]}$$

Isochrone potential:

$$\Phi(r) = -\frac{GM}{b + \sqrt{b^2 + r^2}} \text{ [J]}$$

Jeans growth time:

$$\tau_J = \frac{1}{k_J c_s} = \frac{1}{\sqrt{4\pi G \rho_0}} = 2.3 \times 10^4 \left(\frac{n_H}{10^6 \text{ cm}^{-3}} \right)^{-1/2} \text{ [yr]}$$

Jeans mass:

$$M_J \equiv \frac{4\pi}{3} \rho_0 \left(\frac{\lambda_J}{2} \right)^3$$
$$= \frac{1}{8} \left(\frac{\pi k T}{\mu G} \right)^{3/2} \rho_0^{-1/2}$$
$$= 0.32 \left(\frac{T}{10 \text{ K}} \right)^{3/2} \left(\frac{m_H}{\mu} \right)^{3/2} \left(\frac{10^6 \text{ cm}^{-3}}{n_H} \right)^{1/2} \text{ [M}_\odot\text{]}$$
$$M_J \equiv \frac{\pi^{5/2}}{6} \left(\frac{c_s^2}{G} \right)^{3/2} \frac{1}{\sqrt{\rho}} \text{ [M}_\odot\text{]}$$

Jeans wavelength:

$$\lambda_J \equiv \frac{2\pi}{\kappa_J} = \sqrt{\frac{\pi c_s^2}{G \rho_0}} \text{ [m]}$$

Jeans wavenumber:

$$k_J \equiv \sqrt{\frac{4\pi G \rho_0}{c_s^2}} \text{ [m}^{-1}\text{]}$$

Kepler's third law:

$$P = \sqrt{\frac{4\pi^2}{G(m_1 + m_2)}} a^3 \text{ [yr]}$$

Kramers' Law:

$$\kappa \propto \rho T^{-3.5}$$

Lane-Emden equation:

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d\theta}{d\xi} \right) = -\theta^n$$

Legendre Polynomials:

$$P_0(x) = 1 \text{ [dimensionless]}$$

$$P_1(x) = x \text{ [dimensionless]}$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1) \text{ [dimensionless]}$$

Line profile (intrinsic):

$$\phi_\nu^{\text{intr}} = \frac{m_e c^2}{\pi e^2} \frac{1}{f_{\ell u}} \sigma_\nu^{\text{intr}} \text{ [cm}^2\text{]}, \quad \int \phi_\nu^{\text{intr}} d\nu = 1$$

Lorentz force:

$$\vec{F} = q(\vec{E} + \vec{v} \times \vec{B}) \text{ [N]}$$

Luminosity:

$$L = 4\pi R^2 \sigma T^4 \text{ [erg s}^{-1}\text{]}$$

Luminosity (Faber-Jackson):

$$L_{\text{FJ}} \propto \sigma_v^\alpha \text{ [erg s}^{-1}\text{]}$$

Luminosity (Tully-Fisher):

$$\begin{aligned} L_{\text{TF}} &= \propto v_{\text{max}}^\alpha \text{ [erg s}^{-1}\text{]} \\ &= \left(\frac{M}{L} \right)^{-2} \left(\frac{1}{G^2 \langle I \rangle} \right) v_{\text{max}}^4 \text{ [erg s}^{-1}\text{]} \end{aligned}$$

Luminosity distance:

$$\begin{aligned} d_L &\equiv \sqrt{\frac{L}{4\pi F}} \text{ [Mpc]} \\ d_L(z \ll 1) &\approx \frac{c}{H_0} z \left(1 + \frac{q_0}{2} z \right) \text{ [Mpc]} \\ d_L(\kappa = 0) &\approx \frac{c}{H_0} z \left(1 + \frac{1 - q_0}{2} z \right) \text{ [Mpc]} \end{aligned}$$

Luminosity distance-redshift relation:

$$d_L = d(1 + z) \text{ [Mpc]}$$

Magnetic field:

$$\vec{B} = \mu \vec{H} \text{ [G]}$$

Magnetic monopole density:

$$n_M(t_{\text{GUT}}) \sim \frac{1}{(2ct_{\text{GUT}})^3} \sim 10^{82} \text{ [m}^{-3}\text{]}$$

Magnitude (absolute):

$$M \equiv -2.5 \log_{10} \left(\frac{L}{L_0} \right) \text{ [mag]}$$

Magnitude (apparent):

$$m \equiv -2.5 \log_{10} \left(\frac{F}{F_0} \right) \text{ [mag]}$$

Matter density-scale factor relation:

$$\rho_m = \rho_{m,0} a^{-3} \text{ [g cm}^{-3}\text{]}$$

Maxwell-Boltzmann equation:

$$\begin{aligned} n &= g \left(\frac{mk_B T}{2\pi \hbar^2} \right)^{3/2} \exp \left(-\frac{E}{k_B T} \right) \text{ [m}^{-3}\text{]} \\ &= g \left(\frac{mk_B T}{2\pi \hbar^2} \right)^{3/2} \exp \left(-\frac{h\nu}{k_B T} \right) \text{ [m}^{-3}\text{]} \end{aligned}$$

Mean free path:

$$\ell_{\text{mfp}} = \frac{1}{n\sigma} \text{ [pc]}$$

Mean molecular weight:

$$\mu \equiv \frac{\rho}{nm_{\text{H}}} \text{ [dimensionless]}$$

Metallicity index:

$$[\text{X}/\text{H}] \equiv \log \left(\frac{n(\text{X})}{n(\text{H})} \right)_* - \log \left(\frac{n(\text{X})}{n(\text{H})} \right)_{\odot} \text{ [dimensionless]}$$

Neutrino velocity:

$$v_{\nu} \sim 150(1+z) \left(\frac{m_{\nu}}{1 \text{ eV}} \right)^{-1} \text{ [km s}^{-1}\text{]}$$

Newton:

$$1 \text{ N} \equiv \frac{1 \text{ J}}{\text{m}}$$

Number density:

$$n = \frac{\rho}{\mu m_{\text{H}}} \text{ [cm}^{-3}\text{]}$$

Number density of Lyman- α forest lines:

$$N_{\alpha}(z) \propto (1+z)$$

Oort constants:

$$A \equiv -\frac{1}{2} \left[\left(\frac{dV}{dR} \right)_{R_0} - \frac{V_0}{R_0} \right] \text{ [km s}^{-1} \text{ kpc}^{-1}\text{]},$$
$$B \equiv -\frac{1}{2} \left[\left(\frac{dV}{dR} \right)_{R_0} + \frac{V_0}{R_0} \right] \text{ [km s}^{-1} \text{ kpc}^{-1}\text{]}$$

Opacity:

$$\kappa_{\nu} = \frac{8\pi}{3} \frac{e_e^2}{\mu_e m_u}$$
$$= 0.20(1+X) \text{ [cm}^2 \text{ g}^{-1}\text{]}$$

Opacity (Rosseland mean):

$$\frac{1}{\kappa} = \frac{\pi}{acT^3} \int_0^{\infty} \frac{\partial B}{\partial T} d\nu \text{ [cm}^2 \text{ g}^{-1}\text{]}$$
$$\kappa_{\text{sc}} = 0.20(1+X) \text{ [cm}^2 \text{ g}^{-1}\text{]}$$

Optical depth:

$$\tau = \int_0^s \alpha_{\nu} ds \text{ [dimensionless]}$$

Pascal:

$$1 \text{ P} \equiv 1 \text{ N m}^{-2} \equiv 1 \text{ kg m}^{-1} \text{ s}^{-2} \equiv 1 \text{ J m}^{-3}$$

Peculiar velocity:

$$v_{\text{pec}} = c \left(\frac{z - z_{\text{cos}}}{1+z} \right) \text{ [km s}^{-1}\text{]}$$

Pericenter:

$$r_1 = a(1-e) \text{ [AU]}$$

Photon energy:

$$E_{\gamma} = h\nu \text{ [eV]}$$

Photon occupancy number:

$$n_{\gamma} \equiv \frac{c^2}{2h\nu^3} I_{\nu} \text{ [dimensionless]}$$
$$\bar{n}_{\gamma} \equiv \frac{c^2}{2h\nu^3} \bar{I}_{\nu} = \frac{c^3}{8\pi h\nu^3} u_{\nu} \text{ [dimensionless]}$$

Photon scattering rate (ionized):

$$\Gamma = \frac{c}{\ell_{\text{mfp}}} = n_e \sigma_e c = n_b \sigma_e c = \left(\frac{n_{b,0}}{a^3} \right) \sigma_e c = \frac{4.4 \times 10^{-21} a^3}{\text{s}^{-1}} \text{ [s}^{-1}\text{]}$$

Planck function:

$$B_\nu(T) = \frac{2h\nu^3}{c^2} \frac{1}{\exp(h\nu/k_B T) - 1} \text{ [erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}]$$

Polarization fraction:

$$p(\lambda^2) = \frac{P}{I} \text{ [dimensionless]}$$

Polarization fraction (max value):

$$0 < p_{\max} \leq 0.09 \left[\frac{E(B - V)}{\text{mag}} \right] \approx 0.03 \left[\frac{A_V}{\text{mag}} \right] \text{ [dimensionless]}$$

$$0 < p_V \lesssim 0.03 \tau_V \text{ [dimensionless]}$$

Polarized angle:

$$\chi = \frac{1}{2} \arctan \left(\frac{U}{Q} \right) \text{ [rad]}$$

Polarized intensity:

$$P = \sqrt{Q^2 + U^2} = pI \text{ [Jy]}$$

Polytropic constant:

$$K = \frac{p}{\rho^\gamma} \text{ [dimensionless]}$$

Polytropic equation of state:

$$P = K \rho^\gamma \text{ [P]}$$

Polytropic index:

$$n = \frac{1}{\gamma - 1} \text{ [dimensionless]}$$

Power spectrum:

$$P(k) = 2\pi \int_0^\infty x^2 \frac{\sin(kx)}{kx} \xi(x) dx \text{ [dimensionless]}$$

Precession rate:

$$\Omega_p = \frac{\psi_p}{T_r} = \frac{\Delta\psi - 2\pi}{T_r} \text{ [rad yr}^{-1}]$$

Pressure scale height:

$$\frac{1}{H_P} \equiv -\frac{1}{P} \frac{dP}{dr} \text{ [m}^{-1}]$$

Proper distance:

$$d_p(t_0) = c \int_t^{t_0} \frac{dt}{a(t)} \text{ [Mpc]}$$

$$d_p(t_0, z \ll 1) \approx \frac{c}{H_0} z \left(1 - \frac{1+q_0}{2} z \right) \text{ [Mpc]}$$

Proper motion:

$$\mu = \frac{v_t}{d} \text{ [arcsec yr]}$$

Proper surface area:

$$A_p(t_0) = 4\pi S_\kappa(r)^2 \text{ [Mpc}^2]$$

R parameter:

$$R_V \equiv \frac{A_V}{A_B - A_V} \equiv \frac{A_V}{E(B - V)} \text{ [dimensionless]}$$

Radial period:

$$\begin{aligned} T_r &= 2 \int_{r_1}^{r_2} \frac{dr}{\sqrt{2[E - \Phi(r)] - \frac{L^2}{r^2}}} \text{ [yr]} \\ &= T_\psi = \frac{a^2}{L} \sqrt{1 - e^2} = 2\pi \sqrt{\frac{a^3}{GM}} \text{ [yr]} \end{aligned}$$

Radial velocity:

$$v_r = \Delta \mathbf{V} \cdot \left(\frac{\sin \ell}{-\cos \ell} \right) = (\Omega - \Omega_0) R_0 \sin \ell \text{ [km s}^{-1}\text{]}$$

Radiation density-scale factor relation:

$$\rho_r = \rho_{r,0} a^{-4} \text{ [g cm}^{-3}\text{]}$$

Radiation energy density:

$$u = aT^4 = \frac{4\sigma}{c} T^4 \text{ [erg cm}^{-3}\text{]}$$

Radiation energy density (specific):

$$(u_\nu)_{\text{LTE}} = \frac{4\pi}{c} B_\nu(T) = \frac{8\pi h \nu^3}{c^3} \frac{1}{e^{h\nu/k_B T} - 1} \text{ [erg cm}^{-3} \text{ Hz}^{-1}\text{]}$$

Radiation pressure acceleration:

$$\begin{aligned} g_{\text{rad}} &= -\frac{1}{\rho} \frac{dP_{\text{rad}}}{dr} \text{ [m s}^{-2}\text{]} \\ &= \frac{\kappa F_{\text{rad}}}{c} = \frac{\kappa L_r}{4\pi r^2 c} \text{ [m s}^{-2}\text{]} \end{aligned}$$

Radiative energy transport:

$$\frac{\partial T}{\partial r} = -\frac{3}{16\pi a c} \frac{\kappa \rho L}{r^2 T^3} \text{ [K cm}^{-1}\text{]}$$

Radiative transfer equation:

$$\frac{dI_\nu}{ds} = j_\nu - \alpha_\nu I_\nu \text{ [erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1} \text{ pc}^{-1}\text{]}$$

Random walk:

$$d = \sqrt{N} \ell \text{ [m]}$$

Rate of grain photoelectric heating:

$$\Gamma_{\text{PE}} \approx 4.0 \times 10^{-26} \chi_{\text{FUV}} Z'_d e^{-\tau_d} \text{ [erg s}^{-1}\text{]}$$

Rayleigh-Jeans limit:

$$I_\nu = \frac{2k_B T \nu^2}{c^2} \text{ [erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}\text{]}$$

Redshift of thermal coupling:

$$z_t \approx 140 \left(\frac{\Omega_b h^2}{0.022} \right)^{2/5} \text{ [dimensionless]}$$

Redshift-wavelength-scalefactor relation:

$$\frac{1}{1+z} = \frac{\lambda}{\lambda_0} = a \text{ [dimensionless]}$$

Refractive index:

$$m = \sqrt{\sigma} \text{ [dimensionless]}$$

Relaxation time:

$$\begin{aligned} t_{\text{relax}} &\approx \frac{R}{6v} \frac{N}{\ln(N/2)} \text{ [yr]} \\ t_{\text{relax}} &= \frac{t_{\text{cross}}}{6} \frac{N}{\ln(N/2)} \text{ [yr]} \end{aligned}$$

Robertson-Walker metric:

$$ds^2 = cdt^2 - a(t)^2 \left(\frac{dx^2}{1 - \kappa x^2/R^2} + x^2 \Omega^2 \right)$$

Rotational velocity (at Solar position):

$$V_0 = \sqrt{\frac{GM(< R)}{R_0}} \text{ [m s}^{-1}\text{]}$$

Saha equation:

$$\frac{n_{i+1}}{n_i} = T^{3/2} e^{(-1/T)} \text{ [dimensionless]}$$

Scale factor evolution (radiation-dominated):

$$a_m(t) \propto t^{1/2} \text{ [dimensionless]}$$

Scale factor evolution (matter-dominated):

$$a_m(t) \propto t^{2/3} \text{ [dimensionless]}$$

Scale factor evolution (Λ -dominated):

$$a_{\Lambda}(t) \propto e^{Ht} \text{ [dimensionless]}$$

Scale factor-redshift relation:

$$a = \frac{1}{1+z} \text{ [dimensionless]}$$

Schmidt-Kennicutt law:

$$\frac{\Sigma_{\text{SFR}}}{\text{M}_{\odot} \text{ year}^{-1} \text{ kpc}^{-2}} = (2.5 \pm 0.7) \times 10^{-4} \left(\frac{\Sigma_{\text{gas}}}{\text{M}_{\odot} \text{ pc}^{-2}} \right)^{1.4 \pm 0.15}$$

Schwarzschild radius:

$$r_S \equiv \frac{2GM}{c^2} = 2.95 \left(\frac{M}{\text{M}_{\odot}} \right) \text{ [km]}$$

Serkowski Law:

$$p(\lambda) \approx p_{\text{max}} \exp \left[-K \ln^2 \left(\frac{\lambda}{\lambda_{\text{max}}} \right) \right] \text{ [dimensionless]}$$

Siemen:

$$1 \text{ S} \equiv \text{kg}^{-1} \text{ m}^{-2} \text{ s}^3 \text{ A}^2$$

Solid angle:

$$\Omega = \frac{A}{r^2} \text{ [sr]}$$

Sonic mass:

$$M_s \approx \frac{c_s^2 \ell_s}{G} \text{ [M}_{\odot}]$$

Sound speed:

$$c_s = \sqrt{\frac{\partial P}{\partial \rho}} \text{ [m s}^{-1}]$$
$$c_s \sim \sqrt{\frac{k_B T}{\mu m}} \text{ [m s}^{-1}]$$

Volume (specific):

$$V \equiv \frac{1}{\rho} \text{ [g}^{-1} \text{ cm}^3]$$

Spin temperature:

$$T_s^{-1} = \frac{T_{\gamma}^{-1} + x_{\alpha} T_{\alpha}^{-1} + x_c T_K^{-1}}{1 + x_{\alpha} + x_c} \text{ [K}^{-1}]$$

Spin temperature of 21 cm:

$$\frac{n_1}{n_2} = \frac{g_1}{g_2} \exp \left(-\frac{0.068 \text{ K}}{T_s} \right) \text{ [dimensionless]},$$

Star formation history:

$$\psi(z) = 0.015 \frac{(1+z)^{2.7}}{1 + [1+z/2.9]^{5.6}} \text{ [M}_{\odot} \text{ year}^{-1} \text{ Mpc}^{-3}].$$

Star formation rate:

$$\text{SFR} = -\frac{dM_{\text{gas}}}{dt} \text{ [M}_{\odot} \text{ yr}^{-1}]$$

Star formation rate per unit area:

$$\Sigma_{\text{SFR}} \propto \Sigma_{\text{gas}}^{1.4} \text{ [M}_{\odot} \text{ year}^{-1} \text{ pc}^{-2}]$$

Stefan-Boltzmann constant:

$$\sigma = \frac{2\pi^5 k^4}{15c^2 h^3} \text{ [W m}^{-2} \text{ K}^{-4}]$$

Stefan-Boltzmann Law:

$$F = \sigma T^4 \text{ [W m}^{-2}]$$

Strömgren radius:

$$r_s = 30 \left(\frac{N_{48}}{n_{\text{H}} n_e} \right)^{1/3} \text{ [pc]}$$

Sunyaev-Zeldovich temperature fluctuation:

$$\frac{\Delta T_{\text{SZE}}}{T_{\text{CMB}}} = f(x)y = f(x) \int n_e \frac{k_B T_e}{m_e c^2} \sigma_T dl \quad [\text{dimensionless}]$$

$$\frac{\Delta T_{\text{SZ}}}{T_{\text{CMB}}} = -\tau_e \left(\frac{v_{\text{pec}}}{c} \right) \quad [\text{dimensionless}]$$

Surface brightness (bulge):

$$\mu_{\text{bulge}} = \mu_e + 8.3268 \left[\left(\frac{R}{R_e} \right)^{1/4} - 1 \right] \quad [\text{mag arcsec}^{-2}]$$

Surface brightness (disk):

$$\mu_{\text{disk}} = \mu_0 + 1.09 \left(\frac{R}{h_R} \right) \quad [\text{mag arcsec}^{-2}]$$

Tangential velocity:

$$v_t = \Delta \mathbf{V} \cdot \left(\frac{\cos \ell}{\sin \ell} \right) = (\Omega - \Omega_0) R_0 \cos \ell - \Omega D \quad [\text{km s}^{-1}]$$

Temperature gradient (adiabatic condition):

$$\left| \frac{dT}{dr} \right|_{\text{act}} > \left| \frac{dT}{dr} \right|_{\text{ad}} \quad [\text{K m}^{-1}]$$

or

$$\left| \frac{dT}{dP} \right|_{\text{act}} > \left| \frac{dT}{dP} \right|_{\text{ad}} \quad [\text{K P}^{-1}]$$

Temperature gradient (adiabatic transport):

$$\left(\frac{dT}{dr} \right)_{\text{ad}} = \left(1 - \frac{1}{\gamma} \right) \frac{\mu m_{\text{H}}}{k} \frac{GM_r}{r^2} \quad [\text{K m}^{-1}]$$

Temperature gradient (radiative):

$$\begin{aligned} \nabla_{\text{rad}} &\equiv \left(\frac{d \ln T}{d \ln P} \right)_{\text{rad}} \quad [\text{K P}^{-1}] \\ &= \frac{3}{16\pi acG} \frac{\kappa L P}{m T^4} \quad [\text{K P}^{-1}] \\ &= -\frac{3}{4ac} \frac{\kappa \rho}{T^3} \frac{L_r}{4\pi r^2} \quad [\text{K m}^{-1}] \end{aligned}$$

Temperature-redshift-scalefactor relation:

$$T = (1+z)T_0 = \frac{T_0}{a} \quad [\text{K}]$$

Thomson cross section:

$$\sigma_T = \frac{8\pi}{3} r_{e,0}^2 \quad [\text{m}^2]$$

Time in units of the initial mass doubling time:

$$\tau = \frac{t}{(M_0/\dot{M}_0)} \quad [\text{dimensionless}]$$

Tired Light Hypothesis energy scaling:

$$E = E_0 e^{(-r/R_0)} \quad [\text{eV}]$$

Tully-Fisher relation:

$$L_{\text{TF}} \propto v_{\text{max}}^4 \quad [\text{erg s}^{-1}]$$

Two-point correlation function:

$$\xi(r) = \left(\frac{r}{r_0} \right)^{-1.7} \quad [\text{dimensionless}]$$

Uncertainty Principle: see *Heisenberg Uncertainty Principle*

Velocity (circular):

$$v_c = \sqrt{\frac{GM}{r}} \quad [\text{m s}^{-1}]$$

Velocity FWHM (Gaussian):

$$(\Delta v)_{\text{FWHM}} = \sqrt{8 \ln 2} \sigma_v = 2\sqrt{\ln 2} b \quad [\text{km s}^{-1}]$$

Velocity (radial):

$$v_r = \left(\frac{\Delta\lambda}{\lambda_0} \right) c \text{ [m s}^{-1}\text{]}$$

Velocity (tangential):

$$v_t = d\mu = 4.74 \left(\frac{d}{1 \text{ pc}} \right) \left(\frac{\mu}{1'' \text{ year}^{-1}} \right) \text{ [m s}^{-1}\text{]}$$

Virial parameter:

$$\alpha_{\text{vir}} \sim \frac{\sigma_v R}{GM} \text{ [dimensionless]}$$

Virial theorem:

$$E_{\text{pot}} = -2E_{\text{kin}} \text{ [J]}$$

$$E_{\text{tot}} = \frac{1}{2}E_{\text{pot}} \text{ [J]}$$

Voigt profile:

$$\phi_{\nu}^{\text{Voigt}} \equiv \frac{1}{\sqrt{2\pi}} \int \frac{e^{-v^2/2\sigma_v^2}}{\sigma_v} p_v(v) \frac{4\gamma_{u\ell}}{16\pi^2[\nu - (1 - v/c)\nu_{u\ell}]^2 + \gamma_{u\ell}^2} dv \text{ [erg s}^{-1}\text{ cm}^{-2}\text{ sr}^{-1}\text{]}$$

Wavelength-redshift relation:

$$\lambda_0 = \frac{1}{a(t)}\lambda = (1+z)\lambda \text{ [m]}$$

Wien's Law:

$$B_{\nu}(T) = \frac{2h\nu^3}{c^2} \frac{1}{\exp(h\nu/k_B T) - 1} \text{ [erg s}^{-1}\text{ cm}^{-2}\text{ Hz}^{-1}\text{ sr}^{-1}\text{]}$$

$$B_{\nu}(T) \approx \frac{2h\nu^3}{c^2} \frac{1}{\exp(h\nu/k_B T)} \text{ [erg s}^{-1}\text{ cm}^{-2}\text{ Hz}^{-1}\text{ sr}^{-1}\text{]}$$

$$B_{\nu}(T) \approx \frac{2h\nu^3}{c^2} \exp\left(\frac{-h\nu}{k_B T}\right) \text{ [erg s}^{-1}\text{ cm}^{-2}\text{ Hz}^{-1}\text{ sr}^{-1}\text{]}$$

1.5 General Notes

- $k_B(300\text{ K}) \sim \frac{1}{40}\text{ eV}$
- $k_B(11,000\text{ K}) \sim 1\text{ eV}$
- $k_B(10^7\text{ K}) \sim 1\text{ keV}$
- Systems in thermal equilibrium satisfy normal populations of $(n_1/g_1) > (n_2/g_2)$; inverted populations satisfy $(n_1/g_1) < (n_2/g_2)$ (e.g., masers).
- If a particle scatters with a rate greater than the expansion rate of the Universe, that particle remains in equilibrium.
- Ordinary (i.e., baryonic) matter contributes at most 5% of the Universe's critical density.
- Multipole moment $\ell = 1$ and $\ell = 2$ are the dipole and quadrupole moment, respectively.
- The greatest baryon contribution to the density comes not from stars in galaxies, but rather from gas in groups of galaxies; in these groups, $\Omega \sim 0.02$.
- The ratio of the neutrino temperature T_ν to the CMB temperature T_{CMB} is $\frac{T_\nu}{T_{\text{CMB}}} = \left(\frac{4}{11}\right)^{1/3}$.
- We expect photons to decouple from matter when the Universe is already well into the matter-dominated era (i.e., not in the radiation-dominated era).
- When the Universe was only one second old, the mean-free-path of a photon was about the size of an atom.
- Nuclear binding energies are typically in the MeV range, which explains why Big Bang nucleosynthesis occurs at temperatures a bit less than 1 MeV even though nuclear masses are in the GeV range.
- The electron-positron energy threshold is $\sim 10^{10}\text{ K}$, above which there exist thermal populations of both electrons and neutrinos to make the reaction $p + e^- \leftrightarrow n + \nu$ go equally well in either direction.
- Thermodynamically, nucleons with greater binding energies are more energetically favourable.
- Simplest way to form helium is via deuterium fusion rather than the improbable coincidence of 2 protons and 2 neutrons all arriving at the same place simultaneously to make ${}^4\text{He}$ in one go.
- Nucleosynthesis starts at about 10^{10} K when the Universe was about 1 s old, and effectively ends when it has cooled by a factor of 10, and is about 100 times older.
- Primordial elemental abundances: 75% H, 25% He, trace Li.
- A normal population of states in LTE satisfies $(n_1/g_1)/(n_2/g_2) > 1$ via the Maxwell-Boltzmann equation; inverted populations such as masers satisfy $(n_1/g_1)/(n_2/g_2) < 1$.
- The solid angle subtended by a complete sphere is $4\pi\text{ sr}$.
- Just as the period of a Cepheid tells you its luminosity, the rise and fall time of a type Ia SN tells you its peak luminosity.
- The average Type Ia has a peak luminosity of $L = 4 \times 10^9 L_\odot$ which is 100,000 times brighter than even the brightest Cepheid variable.
- The Hubble distance d_H is the distance between the Earth and any astrophysical object receding away from us at the speed of light.
- At $z = 10^9$, the Jeans mass is $M_J \sim M_\odot$; at $z = 10^6$, the Jeans mass is $M_J \sim M_{\text{gal}}$.
- Apparent magnitudes $0 < m < 6$ are typically visible to the naked eye.
- The absolute magnitude of a light source M is defined as the apparent magnitude m that it would have if it were at a luminosity distance of $d_L = 10\text{ pc}$.
- The apparent magnitude is really nothing more than a logarithmic measure of the flux, and the absolute magnitude is a logarithmic measure of the luminosity.

- When space is positively curved, the proper surface area is $A_p(t_0) < 4\pi r^2$, and the photons from distance r are spread over a smaller area than they would be in flat space. When space is negatively curved, $A_p(t_0) > 4\pi r^2$, and photons are spread over a larger area than they would be in flat space.
- The Benchmark Model has a deceleration parameter of $q_0 \approx 0.55$.
- The angular distribution of the CMB temperature reflects the matter inhomogeneities at the redshift of decoupling of radiation and matter.
- The CMB is linearly polarized at the 10% level.
- CMB polarization probes the epoch of last scattering directly as opposed to the temperature fluctuations which may evolve between last scattering and the present. Moreover, different sources of temperature anisotropies (scalar, vector, and tensor) give different patterns in the polarization: both in its intrinsic structure and in its correlation with the temperature fluctuations themselves.
- The sound speed in the photon-dominated fluid of the early Universe is given by $c_s \approx c/\sqrt{3}$. Thus, the sound horizon is about a factor of $\sqrt{3}$ smaller than the event horizon at this time.
- At recombination, the free electrons recombined with the hydrogen and helium nuclei, after which there are essentially no more free electrons which couple to the photon field. Hence, after recombination the baryon fluid lacks the pressure support of the photons, and the sound speed drops to zero – the sound waves do no longer propagate, but get frozen in.
- In a universe which is dominated by dark matter the expected CMB fluctuations on small angular scales are considerably smaller than in a purely baryonic universe.
- The number density of each of the three flavors of neutrinos (ν_e , ν_μ , and ν_τ) has been calculated to be 3/11 times the number density of CMB photons. This means that at any moment, about twenty million cosmic neutrinos are zipping through your body.
- Due to the steep slope of the IMF, most of the stellar mass is contained in low-mass stars; however, since the luminosity of main-sequence stars depends strongly on mass, approximately as $L \propto M^3$, most of the luminosity comes from high-mass stars.
- Cooling by the primordial gas is efficient only above $T \gtrsim 2 \times 10^4$ K since metal lines cannot contribute to the cooling.
- Whereas in enriched gas molecular hydrogen is formed on dust particles, the primordial gas had no dust so H_2 must form in the gas phase itself, rendering its abundance very small.
- Even a tiny neutral fraction, $X_{HI} \sim 10^{-4}$, gives rise to complete GP absorption due to the large absorption cross section of $Ly\alpha$.
- The key to the detectability of the 21 cm signal hinges on the spin temperature T_s . Only if this temperature deviates from the background temperature, will a signal be observable.
- The spin temperature becomes strongly coupled to the gas temperature when $x_{tot} \equiv x_c + x_\alpha \gtrsim 1$ and relaxes to T_γ when $x_{tot} \ll 1$.
- The ratio of black hole mass and bulge mass is approximately 1/300. In other words, 0.3% of the baryon mass that was used to make the stellar population in the bulge of these galaxies was transformed into a central black hole.
- It is not hard to show that L is actually the length of the vector $\vec{r} \times \dot{\vec{r}}$, and hence that $r^2 \dot{\phi} \equiv L$ is just a restatement of the conservation of angular momentum. Geometrically, L is equal to twice the rate at which the radius vector sweeps out area.
- The density at which grains and gas become well-coupled is around $10^4 - 10^5 \text{ cm}^{-3}$, which is higher than the typical density in a GMC.
- The energy associated with the 21 cm transition is $\ll 1$ K, so even in cold regions atomic hydrogen can be excited.
- Since cosmic rays are relativistic particles, they have much lower interaction cross sections, and thus are able to penetrate into regions where light cannot.

- Sightlines through diffuse gas in the MW have extinction parameters of $R_V \approx 3.1$ as an average value.
- At submillimeter frequencies and above, real materials have only a negligible response to an applied magnetic field – this is because the magnetization of materials is the result of aligned electron spins and electron orbital currents, and an electron spin (or orbit) can change direction only on time scales longer than the period for the electron spin (or orbit) to precess in the local (microscopic) magnetic fields within atoms and solids. These fields are at most $B_i \lesssim 10$ kG, and the precession frequencies are $\omega_p \approx \mu_B B_i / \hbar \lesssim 10^{10} \text{ s}^{-1}$.
- Ray-tracing arguments would lead us to expect the extinction cross section to be equal to the geometric cross section, but diffraction around the target leads to additional small-angle scattering, with the total extinction cross section equal to twice the geometric cross section.
- Molecular hydrogen (H_2) has a binding energy of 4.5 eV.
- The ionization potential of 13.6 eV for atomic hydrogen isn't very different from the dissociation potential of 4.5 eV for molecular hydrogen.
- $\gamma = 4/3$ is the critical value required to have a hydrostatic object.
- Each deuterium burned provides 5.5 MeV of energy, comparable the 7 MeV per hydrogen provided by burning hydrogen, but there are only 2×10^{-5} D nuclei per H nuclei. Thus, at fixed luminosity the “main sequence” lifetime for D burning is shorter than that for H burning by a factor of $2 \times 10^{-5} \times 5.5/7 = 1.6 \times 10^{-5}$.
- Typical accretion rates are of order a few times $10^{-6} M_\odot \text{ yr}^{-1}$, so a $1 M_\odot$ star takes a few times 10^5 yr to form. Thus stars may burn deuterium for most of the time they are accreting.
- The highly forbidden spin-flip transition of 21 cm emission is intrinsically so rare (Einstein A coefficient $A_{ul} = 2.85 \times 10^{-15} \text{ s}^{-1}$) that very long paths are needed for the 21 cm line to be detectable.
- Each subshell has a maximum number of electrons which it can hold: s : 2 electrons, p : 6 electrons, d : 10 electrons, and f : 14 electrons.
- The s subshell is the lowest energy subshell and the f subshell is the highest energy subshell.
- Most of the interstellar gas in the Milky Way is neutral, and $\sim 78\%$ of the neutral hydrogen is atomic, or HI.
- A star is said to be in thermal equilibrium when the total rate of nuclear energy generation is almost perfectly equal to its surface luminosity.
- If the nuclear luminosity is equal to the surface luminosity, the star is in thermal equilibrium and its evolution proceeds on a nuclear timescale. Conversely, if these are not equal, the star is out of thermal equilibrium and its evolution proceeds on a thermal timescale, which usually is much shorter than the nuclear timescale.
- The strongest transitions are electric dipole transitions. These are transitions satisfying selection rules.
- there is a hierarchy in the transition probabilities: very roughly speaking, inter-system lines are $\sim 10^6$ times weaker than permitted transitions, and forbidden lines are $\sim 10^2 - 10^6$ times weaker than inter-system transitions.
- Despite being very “weak,” forbidden transitions are important in astrophysics for the simple reason that every atom and ion has excited states that can only decay via forbidden transitions. At high densities, such excited states would be depopulated by collisions, but at the very low densities of interstellar space, collisions are sufficiently infrequent that there is time for forbidden radiative transitions to take place.
- Even at $T = 10^8 \text{ K}$ Compton scattering reduces the opacity by only 20% of that given by the Thomson scattering equation.
- Degeneracy increases the mean free path considerably, since the quantum cells of phase space are filled up such that collisions in which the momentum is changed become rather improbable.

- In the deep interior of a star, if the actual temperature gradient is just *slightly* larger than the adiabatic temperature gradient, this may be sufficient to carry nearly all of the luminosity by convection. Consequently, it is often the case that either radiation or convection dominates the energy transport in the deep interior of stars, while the other energy transport mechanism contributes very little to the total energy outflow. The particular mechanism in operation is determined by the temperature gradient. However, near the surface of the star, the situation is much more complicated: both radiation and convection can carry significant amounts of energy simultaneously.
- One solar neutrino unit (SNU) corresponds to 10^{-36} captures per second and per target nucleus.
- What created the apparent solar neutrino deficit is the fact that neutrinos can change their flavour, both while travelling through vacuum and more efficiently in the presence of electrons in the solar interior. These so-called **neutrino oscillations** are possible only if neutrinos have mass.
- It is extremely difficult for a star to have more than about 10 percent of the magnitude of its gravitational energy stored in the form of rotational energy.
- Most, but not all, large-amplitude variables pulsate in the fundamental mode.
- Non-radial pulsation tends to produce smaller amplitudes than radial modes in the brightness and colour variation.
- Linear adiabatic theory (LAT) shows that the relative amplitude of pulsation tends to be largest in the outer layers of the star, where the density is lowest.
- Linear non-adiabatic theory (LNAT) of stellar pulsations gives periods which are slightly more realistic than the linear adiabatic theory (LAT), though they are about the same in most cases.
- Non-linear non-adiabatic theory accurately predicts the amplitudes of Cepheids and RR Lyrae stars, which pulsate in one or two radial modes. It has not been able to explain stars like Delta Scuti stars, which can and do pulsate in many radial and non-radial modes.
- The Cepheid instability strip extends from the top of the HR diagram to the bottom, at approximately constant temperature. That is because the instability mechanism occurs at a particular temperature – the temperature at which there is a helium ionization zone in the outer layers of the star.
- Aside from the stars of smallest mass ($M < 0.25 M_{\odot}$), we can roughly distinguish between two types of radiative/convective models: (1) radiative core + convective envelope (lower MS) and (2) convective core + radiative envelope (upper MS.) The transition from one type to the other occurs near $M = 1 M_{\odot}$.
- For an ideal monatomic gas, $\gamma = 5/3$ and convection will occur in some region of a star when $d \ln P / d \ln T < 2.5$.
- The base of the Sun's present-day convection zone is at $0.714 R_{\odot}$.
- Many Herbig Ae stars are still contracting to the main sequence, and are thus either still surrounded by remnants of their stellar cocoons, or have developed massive stellar winds.
- Many metallic-line A-type (Am) stars show the **anomalous luminosity effect** in which the luminosity criteria in certain spectral regions (in particular, the region from $4077 - 4200 \text{\AA}$) may indicate a giant or even a supergiant luminosity, whereas other regions indicate a dwarf luminosity or even lower.
- All T dwarfs are brown dwarfs.