Planning (VLA) observations

Loránt Sjouwerman, NRAO



Sixteenth Synthesis Imaging Workshop 16-23 May 2018















Outline

- General advice on planning any (ground based) observation
- AUI telescopes: the GBT, ALMA, VLBA, and in particular the VLA
 - VLA, VLBA and ALMA are radio frequency interferometers, GBT is a single radio dish
 - VLA is NRAO operated, VLBA is LBO operated (soon NRAO?) and GBT is GBO operated
 - ALMA is a joint endeavor (i.e. somewhat special with its own rules); for ALMA see next talk
- Specific advice for planning VLA observations (I): Proposing
 - Planning starts before proposing/applying for time
- Introduction to the web-based tool to apply for AUI telescope time
 - Proposal Submission Tool (PST) found at https://my.nrao.edu/
- Specific advice for planning VLA observations (II): Scheduling
 - Operations and logistics: dynamic scheduling and scheduling priorities
 - Creating observing schedules is about determining the optimum trade-offs (for YOUR science case)
- Introduction to the web-based tool to schedule VLA observations
 - Observation Preparation Tool (OPT) through PST portal or via https://obs.vla.nrao.edu/opt/

Planning observations




Planning any (ground based) observation

- Location of the telescope
 - Determines what part of the sky is visible
 - Declination range, e.g., for the VLA: -44° to +90°
 - Determines the dry/wet seasons and other weather and atmospheric related issues (e.g., troposphere and ionosphere)

Determines the level of interfering human activities (light pollution, radio frequency interference)

Particular telescope may have closures

 (e.g., maintenance) or instrumentation
 may be unavailable for periods of time,
 usually well announced





Planning any (ground based) observation

- Scientific and technical justification
 - Operating telescopes costs money; observing time is precious!
 - Time is available for well justified (scientific) projects on a **competitive** basis
 - Scientific justification (next slide) should clearly outline the broader scientific relevance and define goals that outline why only (new) telescope time using this particular instrument on a specific target will achieve that goal and contribute to the general knowledge of the research topic
 - Technical justification is to demonstrate that the choice of instrumentation, the observing time and the proposed methodology will yield the result using the requested resources in the most optimal way (i.e., within reason)
 - Projects are ranked by committees based on scientific and technical merit but even a top-notch scientific case may end up without observing time if the justification is poorly argued or technically not feasible
 - Conditional to constraints known to the TAC (time/resources available, day/night and other operational conditions, competition)

Scientific and technical justification

• Why?

- Science context and motivation: what makes this subject scientifically interesting for the field?
- Outline the broader impact of the science topic and the importance for astrophysics

What?

 Specific science questions: what missing but obtainable knowledge is furthering the field?

How?

 Specific science goals: how are these observations, resulting observables, going to answer the questions above and how will the science goals be achieved by this program?



Scientific and technical justification

- Must appeal to non-expert astrophysicist reviewer (try to avoid all jargon!)
- Who?
 - Target selection criteria: why are these object chosen for the observations, can the sample be smaller, can it instead be done with similar sources at less popular LST ranges?
- When and where?
 - Should this wait for a better opportunity, hosted by another telescope, that is, why now and why this telescope?
- What is needed?
 - Resources such as instrumental setups, observing time, operational conditions, data reduction requirements
 - Show it can be done!






AUI operated telescopes





- Robert C. Byrd Green Bank Telescope (GBT) operated by GBO
 - Fully steerable single dish antenna in WV
 - 100 meter diameter
 - Unblocked view
 - About 100 MHzto 100 GHz
- Not an interferometer (often used as an array element)
- Separate single dish workshops elsewhere






- Atacama Large Millimeter Array (ALMA) operated by JAO
 - 66 (50 movable and 4+12 stationary) antennas in Chile
 - 12 & 7 meter diameter
 - Baselines up to 16 km
 - About 50 GHz to
 I THz (1000 GHz, 300 μm)
- Shared instrument
 (East-Asia, Europe, North America)
- Separate talk here





- Very Long Baseline Array
 (VLBA) operated by LBO
 - 10 stationary antennas
 spread across the US
 - 25 meter diameter
 - Baselines up to 8500 km
 - About 300 MHz to 90 GHz
- Frequently combined with others (US, Europe, Asia, etc)





- Karl G. Jansky Very Large Array (VLA) operated by NRAO
 - 27 (+1) movable antennas near Socorro NM!
 - 25 meter diameter
 - Baselines up to36 km
 - About 70 MHzto 50 GHz
- Site tour on Sunday...





- Movable antennas: compact ("D" configuration) through spread out ("A")
 - Baseline coverage 35 m to ~I km (D), 3 km (C), II km (B), 36 km (A)
 Highest angular resolution in A, best surface brightness sensitivity in D for a given frequency
 - Configuration change every ~4 months
 "Any" array configuration includes regular and "move" time configurations
- Every antenna currently has 8 permanent receiver feeds
 - Continuous frequency coverage from 1 to 50 GHz (working on < 1GHz)
 - Continuum bandwidth of 2 or up to 8 GHz dual polarization (L: IGHz)
 - Switching between feeds takes 10-20 seconds, setups 10-40 seconds
- Extremely flexible WIDAR correlator
 - Up to 64 independent "spectral windows", 31 kHz to 128 MHz (< 8 GHz)
 - Frequency channels of 2 MHz down to 122 Hz (single pol, without recirculation)



- Movable antennas: compact ("D" configuration) through spread out ("A")
 - Baseline coverage 35 m to ~I km (D), 3 km (C), I I km (B), 36 km (A)



- Movable antennas: compact ("D" configuration) through spread out ("A")
 - Baseline coverage 35 m to ~I km (D), 3 km (C), II km (B), 36 km (A)
 Highest angular resolution in A, best surface brightness sensitivity in D for a given frequency
 - Configuration change every ~4 months
 "Any" array configuration includes regular and "move" time configurations
- Every antenna currently has 8 permanent receiver feeds
 - Continuous frequency coverage from I to 50 GHz (working on < IGHz)
 - Continuum bandwidth of 2 or up to 8 GHz dual polarization (L: IGHz)
 - Switching between feeds takes 10-20 seconds, setups 10-40 seconds
- Extremely flexible WIDAR correlator
 - Up to 64 independent "spectral windows", 31 kHz to 128 MHz (< 8 GHz)
 - Frequency channels of 2 MHz down to 122 Hz (single pol, without recirculation)






NRAO

- Movable antennas: compact ("D" configuration) through spread out ("A")
 - Baseline coverage 35 m to ~I km (D), 3 km (C), I I km (B), 36 km (A)
 Highest angular resolution in A, best surface brightness sensitivity in D for a given frequency
 - Configuration change every ~4 months
 "Any" array configuration includes regular and "move" time configurations
- Every antenna currently has 8 permanent receiver feeds
 - Continuous frequency coverage from I to 50 GHz (working on < IGHz)
 - Continuum bandwidth of 2 or up to 8 GHz dual polarization (L: IGHz)
 - Switching between feeds takes 10-20 seconds, setups 10-40 seconds
- Extremely flexible WIDAR correlator
 - Up to 64 independent "spectral windows", 31 kHz to 128 MHz (< 8 GHz)
 - Frequency channels of 2 MHz down to 122 Hz (single pol, without recirculation)



Frequency bands



Proposing for VLA observations



- Two proposal rounds per year
 - Deadlines near February Ist and August Ist
 - Both for Regular and Large proposals
- DDT: exploratory/target of opportunity
 - Small amount of time (typically couple of hours)
 - Must argue why the normal deadline could not be met
- Scientific justification should be clear and to the point
 - See a previous slide
- Technical justification should address and resolve all potential issues
 - Includes setup, sources, data size and reduction, analysis, logistics, etc.



- Read the Call for Proposals (CfP)
 - Changes, new opportunities, special instructions
- Read the on-line documentation!
 - Instructions and restrictions change
 - Suggestions and hints for higher success rate
- Helpdesk (https://help.nrao.edu) is available
 - Do not wait until just before the deadline, response may be delayed!
 - Best to start asking for help about 2-3 weeks before the deadline
 - i.e., just after the CfP is issued



- VLA receiver band
 - Specify the frequency to observe and why
 - Field-of-view at this frequency
 - Need mosaicking?
- VLA array configuration (A-D):
 - What angular resolution for the smallest details?
 - What angular scale for the largest structure?
 - Need multi-configuration?
 - B and C configurations less oversubscribed, can those be used instead?
- Need for subarrays? Use less antennas in "move" time?



- VLA frequency setup
 - Standard continuum observations?
 - In C, X, Ku bands, use 8-bit or 3-bit?
 - Multi-frequency synthesis?
 - Spectral line, with continuum or complex correlator configuration?
 - Bandwidth per spectral window
 - up to 64 independently configurable
 - Channel separation within each spectral window
 - Polarization within each spectral window
 - Other details like rest frequencies, velocities/redshifts
- Data rate for the setup within the limits?




- Observing time request
 - Use exposure calculator to estimate sensitivity, image RMS
 - https://obs.vla.nrao.edu/ect
 - Signal-to-noise needed for science, self-calibration?
 - Dynamic range limited or detection experiment?
 - Include overhead for calibration, special processing needs
 - (u,v) coverage considerations (snapshot, full polarization tracks)
 - Specific LST ranges that the sources are above the elevation limit (8°)
 - How many blocks of observing time versus available?
- Is the total project data volume realistic?
 - Data reduction resources (cpu, disks, people, clock time) available?





- Observing time request
 - Use exposure calculator to e
 - https://obs.vla.nrao.edu/ect
 - Signal-to-noise needed for selections
 - Dynamic range limited or de
 - Include overhead for calibration
 - (u,v) coverage considerations (s)
 - Specific LST ranges that the soul
 - How many blocks of observ.
- Is the total project data volume real
 - Data reduction resources (cpu,






- Observing time request
 - Use exposure calculator to estimate sensitivity, image RMS
 - https://obs.vla.nrao.edu/ect
 - Signal-to-noise needed for science, self-calibration?
 - Dynamic range limited or detection experiment?
 - Include overhead for calibration, special processing needs
 - (u,v) coverage considerations (snapshot, full polarization tracks)
 - Specific LST ranges that the sources are above the elevation limit (8°)
 - How many blocks of observing time versus available?
- Is the total project data volume realistic?
 - Data reduction resources (cpu, disks, people, clock time) available?







- Is source logistically observable?
 - Weather, e.g., decreased time for high frequencies in late summer



High-frequency observing in January mostly okay, even at LST 22-24h Mostly poor high-frequency observing conditions in August LST 6-22h

- Is source logistically observable?
 - Weather, e.g., decreased time for high frequencies in late summer
 - Time of day, e.g., maintenance during working hours, less RFI at night
 - Available time function of LST, e.g., Galactic LST at 18h in high demand
 - Solar activity, twilight observing, proximity to the Sun





Available time as function of LST: daytime used for maintenance and tests

Proposal Submission Tool




- NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)
 - Things change, please read the CfP and on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, allow for response time
- Create new proposal and select instrument: VLA, VLBA, GBT (& GMVA)
- Follow the tree on the left to construct the proposal
- Submit (and wait)
- Disposition letters 3-4 months later





• NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)




- NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)
 - Things change, please read the CfP and on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, allow for response time
- Create new proposal and select instrument: VLA, VLBA, GBT (& GMVA)
- Follow the tree on the left to construct the proposal
- Submit (and wait)
- Disposition letters 3-4 months later






- NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)
 - Things change, please read the CfP and on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, allow for response time





- NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)
 - Things change, please read the CfP and on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, allow for response time
- Create new proposal and select instrument: VLA, VLBA, GBT (& GMVA)



- NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)
 - Things change, please read the CfP and on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, allow for response time
- Create new proposal and select instrument: VLA, VLBA, GBT (& GMVA)

 NRAO: Proposals Mozilla Firefox (on gygax)






- NRAO portal at https://my.nrao.edu (ALMA portal: almascience.org depends on region)
 - Things change, please read the CfP and on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, allow for response time
- Create new proposal and select instrument: VLA, VLBA, GBT (& GMVA)

Follow the tree on the left to construct the proposal

- Submit (and wait)
- Disposition letters3-4 months later






Scheduling of VLA observations



- VLA is dynamically scheduled and uses scheduling priorities
 - A: most likely to be completed
 - C: filler, i.e., when time available and schedule appropriate
- Typically not known when a particular observation takes place
 - For a given array configuration and LST, select from the list of available observing schedules:
 - Highest priority at the top of the list
 - Scheduling priority
 - Science priority (i.e., competition from other Pls)
 - Deselected if science conditions (weather) unfavorable
 - Deselected if operational constraints unfavorable



- VLA is dynamically scheduled and uses scheduling priorities
 - A: most likely to be completed
 - C: filler, i.e., when time available and schedule appropriate






VLA LST hours

- VLA is dynamically scheduled and uses scheduling priorities
 - A: most likely to be completed
 - C: filler, i.e., when time available and schedule appropriate
- Typically not known when a particular observation takes place
 - For a given array configuration and LST, select from the list of available observing schedules:
 - Highest priority at the top of the list
 - Scheduling priority
 - Science priority (i.e., competition from other Pls)
 - Deselected if science conditions (e.g., weather) unfavorable
 - Deselected if operational constraints unfavorable



- VLA is dynamically scheduled and uses scheduling priorities
 - A: most likely to be completed



- Some PI control on trade-offs in the scheduling constraints for higher success rate, especially for scheduling priority B and C
- Read the documentation:
 - https://science.nrao.edu/facilities/vla/
 - Observational Status Summary
 - Guide to VLA observing
 - OPT manual
- Ask for help!
 - https://help.nrao.edu/




Observation Preparation Tool





Observation Preparation Tool (OPT)


- Use to create VLA observing schedules, a.k.a. "scheduling blocks (SBs)"
- NRAO portal at https://obs.vla.nrao.edu/opt/



Observation Preparation Tool (OPT)

- Use to create VLA observing schedules, a.k.a. "scheduling blocks (SBs)"
- NRAO portal at https://obs.vla.nrao.edu/opt/

Project is created in the OPT a few weeks before the array configuration




Observation Preparation Tool (OPT)

- Use to create VLA observing schedules, a.k.a. "scheduling blocks (SBs)"
- NRAO portal at https://obs.vla.nrao.edu/opt/
- Project is created in the OPT a few weeks before the array configuration
 - NRAO fills in read-only details, e.g., Pl, time, array, scheduling priority(!)
 - Proposers responsible for schedule and observing constraints (LST, API, ..)
 - Things change, please read the on-line documentation!
 - Helpdesk (https://help.nrao.edu/) is available, please use it
 - Schedule is checked for logistical issues, but not for science!
- Scheduling may require making trade-offs, depending on science goals
- Not all SBs will be completed (priority/competition, weather, constraints, operations)
- Observations are directly archived and pipelined, proprietary period
 - Proposers may be contacted about the pipelined data products

- Using the OPT, login and navigate, "look and feel"
- Create project SB (scheduling block) from scratch
- Hypothetical project to observe PN
 - Lines and continuum
- Experiencing the use of
 - SCT, source catalog tool
 - Search for calibrators, create a new source
 - RCT, resource (instrument setup) catalog tool
 - Default continuum setups, create a line setup
 - OPT, observation preparation (scan sequence) tool
 - Change (re)source, scan sequence, play with conditions


- Using the OPT, login and navigate, "look and feel"
- Create project SB (scheduling block) from scratch
- Hypothetical project to observe PN
 - Lines and continuum



- Using the OPT, login and navigate, "look and feel"
- Create project SB (scheduling block) from scratch
- Hypothetical project to observe PN
 - Lines and continuum
- Experiencing the use of
 - SCT, source catalog tool
 - Search for calibrators, create a new source
 - RCT, resource (instrument setup) catalog tool
 - Default continuum setups, create a line setup
 - OPT, observation preparation (scan sequence) tool
 - Change (re)source, scan sequence, play with conditions




Using the OPT, login and navigate, "look and feel"



- Using the OPT, login and navigate, "look and feel"
- Create project SB (scheduling block) from scratch
- Hypothetical project to observe PN
 - Lines and continuum
- Experiencing the use of
 - SCT, source catalog tool
 - Search for calibrators, create a new source
 - RCT, resource (instrument setup) catalog tool
 - Default continuum setups, create a line setup
 - OPT, observation preparation (scan sequence) tool
 - Change (re)source, scan sequence, play with conditions



Using the OPT, login and navigate, "look and feel"




- Using the OPT, login and navigate, "look and feel"
- Create project SB (scheduling block) from scratch
- Hypothetical project to observe PN
 - Lines and continuum
- Experiencing the use of
 - SCT, source catalog tool
 - Search for calibrators, create a new source
 - RCT, resource (instrument setup) catalog tool
 - Default continuum setups, create a line setup
 - OPT, observation preparation (scan sequence) tool
 - Change (re)source, scan sequence, play with conditions





Using the OPT, login and navigate, "look and feel"



- Using the OPT, login and navigate, "look and feel"
- Create project SB (scheduling block) from scratch
- Hypothetical project to observe PN
 - Lines and continuum
- Experiencing the use of
 - SCT, source catalog tool
 - Search for calibrators, create a new source
 - RCT, resource (instrument setup) catalog tool
 - Default continuum setups, create a line setup
 - OPT, observation preparation (scan sequence) tool
 - Change (re)source, scan sequence, play with conditions

Questions?

